

# Effects of antidepressants on parameters, malondialdehyde, and diphenyl-2-picryl-hydrazyl levels in mice spermatozoa

Ladan Bandegi<sup>1</sup> M.Sc., Morteza Anvari<sup>2</sup> Ph.D., Mahmood Vakili<sup>3</sup> Ph.D., Arezoo Khoradmehr<sup>2</sup> M.Sc., Aghdas Mirjalili<sup>1</sup> M.Sc., Ali Reza Talebi<sup>2</sup> Ph.D.

1. Department of Biology and Anatomy, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
2. Research and Clinical Center for Infertility, Yazd Reproductive Sciences Institute, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
3. Department of Community Medicine, School of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

## Corresponding Author:

Ali Reza Talebi, Research and Clinical Center for Infertility, Bouali Ave., Safaeyeh, Yazd, Iran.  
**Email:** prof\_talebi@ssu.ac.ir  
**Tel:** (+98) 9131532174

Received: 21 December 2016  
 Revised: 21 December 2018  
 Accepted: 13 March 2018

## Abstract

**Background:** Prescribing antidepressant drugs is becoming common. These drugs are known to affect sexual functions.

**Objective:** The study is aimed to assess the effects of amitriptyline and venlafaxine on sperm parameters and evaluate Malondialdehyde (MDA) and 1, 1-Diphenyl-2-picryl-hydrazyl values in BALB/ mice spermatozoa.

**Materials and Methods:** Forty adult male BALB/c mice were separated into five groups. Group I (control) received distilled water; group II amitriptyline (4 mg/kg); group III amitriptyline (4 mg/kg) +vitamin C (10 mg/kg); group IV venlafaxine (2 mg/kg); and group V received vitamin C (10 mg/kg) + venlafaxine (2 mg/kg). All drugs were administered by oral gavage for 35 days. After excision of caudal epididymis, it was located in 1 mL Ham's F10 medium at 37°C for 15 min and then analysis of sperm parameters was performed. To examine lipid peroxidation and total antioxidant capacity, the MDA and 1, 1-Diphenyl-2-picryl-hydrazyl were measured, respectively.

**Results:** The mean sperm parameters in the group treated with amitriptyline were significantly lower than in the other groups. MDA tests showed a significant difference between amitriptyline and control groups ( $p=0.007$ ).

**Conclusion:** The results of this study showed that amitriptyline consumption can weaken sperm parameters, which can be attributed to the increased production of ROS and toxicity resulting from amitriptyline consumption. Moreover, venlafaxine improved sperm parameters in mice and the lipid peroxidation in this group did not change compared to the control group.

**Key words:** Amitriptyline, Venlafaxine, MDA, Sperm chromatin.

*This article extracted from M.Sc. thesis. (Ladan Bandegi)*

## Introduction

Antidepressant drugs are widely used for the treatment of depression, anxiety-related diseases, and stress (1). The classification of antidepressant drugs is dependent on the drugs chemical structure and how the drugs function (2). Accordingly, antidepressant drugs fall into five families namely tricyclic antidepressants, selective serotonin reuptake inhibitors (SSRIs), serotonin-norepinephrine reuptake inhibitors, monoamine oxidase inhibitors, and atypical antidepressants (3). Reports indicate that antidepressant drugs have a toxic influence on body organs, particularly the male

reproductive system. Venlafaxine is one of the drugs belonging to the serotonin-norepinephrine reuptake inhibitor family. Venlafaxine, similar to tricyclic antidepressants, inhibits the reuptake of serotonin and norepinephrine, but the adverse effects are lower (4). The treatment of rats with venlafaxine has indicated that this drug modulates the oxidative stress induced by depression in the brain and medulla of rats (5).

Amitriptyline is a component of the tricyclic antidepressant family (6). Amitriptyline is used in the treatment of depression, fibromyalgia, chronic fatigue syndrome, migraine, irritable bowel syndrome, and atypical facial pain.

Amitriptyline acts by inhibiting the reuptake of serotonin and norepinephrine (6). Several studies have shown that amitriptyline has toxic effects that work through the enhancement of oxidative stress (7). A number of studies have demonstrated a high level of lipid peroxidation in the presence of amitriptyline (7, 8). Oxidative stress occurs as the result of the imbalance between reactive oxygen species (ROS) production and the activity of the antioxidant system (9, 10). The increased production of ROS may cause infertility (11). The targets of ROS in biological systems are nuclear DNA, proteins, and lipids. Owing to the existence of many polyunsaturated fatty acids in the spermatozoa membrane, it is very sensitive to ROS attacks (12).

Different factors increase testicular oxidative stress, including: exposure to pesticides, industrial chemical materials, and high doses of metals like iron, cadmium and lead, as well as several drugs such as antineoplastic drugs, antibacterials, antimalarials, calcium canal blockers, and antidepressants (10). Malondialdehyde (MDA) is one of the by-products of lipid peroxidation, that are produced by enzymatic or non-enzymatic processes (14). Increased lipid peroxidation and consequently enhanced MDA, may alter sperm functions, thus decreasing male fertility (15). MDA concentration and diphenyl-1-picrylhydrazyl (DPPH) represent lipid peroxidation and the total antioxidant capacity, respectively. In this study, we investigated the effects of amitriptyline and venlafaxine drugs on sperm parameters and the levels of MDA and DPPH of sperm in BALB/c mice.

Since not much attention has been paid to the effect of antidepressant drugs, especially the SNRL family of drugs, on sperm parameters and lipid peroxidation, the present study aims to examine the impact of amitriptyline and venlafaxine drugs on sperm parameters and on MDA and DPPH levels in BALB/C mice.

## Materials and methods

### Animals and treatments

In this experimental study, 40 adult male BALB/c mice, weighing 20-25 gr and age of 8 wk were obtained from animal house. The mice were kept for 12 hr in light/dark cycle with 22-25°C of temperature, and were fed with standard animal chow and water.

Venlafaxine and amitriptyline were obtained from Parsdarou and Abidi Pharmaceutical Company, Iran, respectively. Venlafaxine and vitamin C were dissolved in physiological saline and amitriptyline was dissolved distilled water. The animals were divided into five groups.

The group I (controls) was treated with distilled water, group II received amitriptyline (4 mg/kg) (2), group III amitriptyline (4 mg/kg)+vitamin C (10 mg/kg) (2, 16), group IV venlafaxine (2 mg/kg) (5), and group V vitamin C (10 mg/kg)+venlafaxine (2 mg/kg) (5, 16), to form the gavage. All groups were treated in mouse spermatozoa period (35 days), and after the treatment period, they were killed by the displacement of the neck vertebrae.

### Epididymal sperm aspiration and sperm analysis

Thirty-five days later, after the mice were killed and the caudal epididymis was brought out, it was placed in 1 mL of culture medium Hams F10, which has been previously placed in the CO<sub>2</sub>-bearing incubator. By slowly tearing up the tissue, sperms were extracted and placed in an incubator with 5% CO<sub>2</sub> at 37°C for 20 min. To analyze the sperms, viability was evaluated by stained eosin, and then motility and sperm concentration were evaluated via Makler chamber with ×100 magnification optical microscopes (Olympus Co. Tokyo, Japan).

Motility was classified into progressive (grade a and b: fast and slow) and non-progressive (grade c) spermatozoa. For evaluating normal morphology, stained

Papanicolaou was used and 200 spermatozoa were counted using a  $\times 100$  magnification light microscope (17). All the chemical agents and pigments were purchased from Merck & Co, Germany.

### Lipid peroxidation

The concentration of the lipid peroxidation product (MDA) or the thiobarbituric acid-reactive substance was determined according to the modified method of Uchiyama and Mihara. In brief, 0.375 ml HCL 20% and 0.375 ml TBA 0.6% were added to 0.1 ml semen plasma. The mixture was heated for 60 min in a boiling water bath. After cooling, 1.25 ml of n-butanol was added, and the mixture was shaken. Thereafter, the mixture was centrifuged at  $2000\times g$  for 5 min, and the absorbance of the upper solution was determined at 532 nm (18).

### DPPH measurement

The total antioxidant capacity was performed through the trapping of plasma-free radicals by reducing diphenyl-2-picrylhydrazyl. The DPPH stocks were prepared in methanol, which was buffered by acetic acid. The buffered methanol was prepared by mixing 40 mL of acetate buffer 0.1 M to 60 mL of methanol. The other solutions and chemicals were analytic grade. The reaction tubes were wrapped in aluminum foils and kept at  $30^{\circ}\text{C}$  for 30 min. All measurements were performed at weak light at the wavelength of 517 nm using a spectrophotometer (19).

### Ethical consideration

The experimental project was confirmed by the ethics committee of Shahid Sadoughi University of Medical Sciences, Yazd, Iran (IR.SSU.MEDICINE.REC.1394.274). The protocols of the Institutional Animal Care and Use Committee were followed during the course of the experiment in terms of the handling, maintenance, treatment, and killing of the animals.

### Statistical analysis

All data were analyzed by SPSS software (Statistical Package for the Social Sciences, version 18.0, SPSS Inc., Chicago, Illinois, USA). One-way ANOVA was applied to evaluate the data, and LSD post-test was performed to determine the difference between the two groups. Two-sided  $p < 0.05$  indicated a statistically significant difference between sperm evaluations.

## Results

### Sperm parameters

Table I indicates that in all sperm parameters including progressive motility, non-progressive motility, immotile, normal morphology, viability except concentration, there was a significant difference between groups II and IV ( $p=0.001$ ,  $0.007$ ,  $0.001$ ,  $0.005$ , and  $0.001$  respectively). Moreover, the motility parameters, viability and immotile spermatozoa were significantly different between groups II and III ( $p=0.009$ ,  $0.001$ , and  $0.001$  respectively).

There was a significant difference between groups I and IV in normal morphology, non-progressive motility, and viability ( $p=0.024$ ,  $0.038$  and  $0.019$  respectively). On the other hand, there was a significant difference between groups I and II with regard to progressive motility, immotile, sperm concentration and viability parameters ( $p=0.001$ ,  $0.001$ ,  $0.046$ , and  $0.001$  respectively).

### Assessment of Oxidative Stress

According to Table II, the amount of MDA in group II is higher than in the other groups, but it is lower in group V than in the other groups. The amount of MDA in drug groups with vitamin C is lower than the drug groups without vitamin C. However, there was a significant difference between groups IV and V ( $p=0.002$ ), groups I and II ( $p=0.007$ ) and was not significant between groups II and III.

In the DPPH test, the total antioxidant capacity amount in groups II, III, and IV decreased compared to the control group, but

it increased in group V compared to the other groups. However, these results were not statistically significant.

**Table I.** The results of sperm parameter analysis in study groups (n=8)

Variable	Group I	Group II	Group III	Group IV	Group V	p-value
Sperm Concentration (10 <sup>6</sup> )	22.41 ± 4.38	18.11 ± 3.72	16.86 ± 3.15	22.06 ± 2.45	22.61 ± 3.09	0.046 <sup>a</sup> 0.011 <sup>b</sup> <0.001 <sup>a</sup>
Progressive Motility (%)	44.12 ± 22.50	13.75 ± 4.49	30.87 ± 6.53	52.75 ± 5.39	48.50 ± 7.23	0.04 <sup>b</sup> 0.009 <sup>e</sup> <0.001 <sup>g</sup> 0.038 <sup>c</sup> 0.002 <sup>d</sup> 0.035 <sup>e</sup> 0.007 <sup>g</sup>
Non-progressive motility (%)	16.25 ± 10.84	13.37 ± 5.57	22.75 ± 8.94	25.50 ± 10.33	30.25 ± 7.42	<0.001 <sup>a</sup> 0.012 <sup>c</sup> 0.012 <sup>d</sup> 0.001 <sup>e</sup> <0.001 <sup>g</sup>
Immotile (%)	39.12 ± 23.35	71.75 ± 5.57	46.75 ± 9.91	21.25 ± 12.17	21.12 ± 4.91	0.024 <sup>c</sup> 0.005 <sup>g</sup> <0.001 <sup>a</sup> 0.019 <sup>c</sup>
Normal morphology (%)	43.71 ± 15.39	38.83 ± 11.39	47.00 ± 10.03	58.5 ± 10.47	54.37 ± 13.67	<0.001 <sup>a</sup> 0.017 <sup>d</sup> 0.001 <sup>e</sup> <0.001 <sup>g</sup>
Viability (%)	64.62 ± 22.58	31.25 ± 6.88	54.12 ± 10.10	80.25 ± 12.42	80.62 ± 5.15	

To analyze data we used standard s one way ANOVA test.

The mean difference was significant at the 0.05 level. All data are presented as mean ± SD)

a: Significant difference between group I and II

c: Significant difference between group I and IV

e: Significant difference between group II and III

g: Significant difference between group II and IV

b: Significant difference between group I and III

d: Significant difference between group I and V

f: Significant difference between group IV and V

**Table II.** Seminal plasma levels of total antioxidant capacity and malondialdehyde in study groups (n=8)

Variable	Group I	Group II	Group III	Group IV	Group V	p-value
MDA (nmol/ml)	0.0075 ± 0.002	0.0111 ± 0.003	0.0085 ± 0.002	0.0101 ± 0.002	0.0059 ± 0.0009	0.007 <sup>a</sup>
DPPH (µM)	991.92 ± 124.53	785.71 ± 58.961	838.40 ± 60.58	873.18 ± 71.14	935.65 ± 46.36	0.002 <sup>b</sup>

To analyze data we used standard s one way ANOVA test. The mean difference was significant at the 0.05 level. All data are presented as mean ± SD

a: Significant difference between group I and II

b: Significant difference between group IV and V

MDA: Malondialdehyde

DPPH: Diphenyl-1-picrylhy drazyl

## Discussion

The mains finding of the present study shows that amitriptyline consumption can weaken sperm parameters. Moreover, venlafaxine can improve sperm parameters in mice.

Different studies indicated that antidepressant drugs could have a negative effect on the male reproductive system. However, there is little knowledge of the effect of antidepressant drugs consumption on the ROS production. The purpose of this study is to investigate the influence of antidepressant

drugs on sperms parameters and ROS production. In addition, the influence of ROS production on the sperms parameters has been investigated. Our finding showed the toxicity of amitriptyline lead to decrease in the sperm parameters including sperm concentration, progressive motility, and viability. In our study, amitriptyline alone and with vitamin c showed a significant difference in sperm parameters. Actually, consumption of amitriptyline with vitamin c could improve progressive motility, non-progressive motility, viability and decrease immotile. In agreement with our study, Hassanane and co-workers

showed a decrease in sperm concentration as well as an increase in morphological abnormalities in the sperm of amitriptyline-treated mice (2).

It has been reported in the literature that the consumption 75-300 mg of trimipramin for 8 wks affects spermatogenesis which is in agreement with our results from amitriptyline-treated mice (20). The finding of the present study showed that venlafaxine than control group increased non-progressive, normal morphology, viability and decreased immotile.

Also, no significant change was observed in sperm concentration. Tanrikut and colleagues have analyzed the results of male infertility examination on two men who were receiving the citalopram, sertraline, and venlafaxine. The semen analysis of patient 1 who was receiving citalopram indicated oligospermia and 1% motility.

The semen analysis of patient 2 who was receiving sertraline revealed normal volume and no motility sperm (21). In another study conducted by Tanrikut, the consumption of paroxetine and SSRI for 5 wks had no adverse effects on key sperm parameters, However, it increased the DNA fragmentation significantly (22). The effects of SSRIs on sperm parameters have been investigated in two separate studies and it has been reported that SSRI exerted a negative influence on sperm parameters and quality, which is in contrast with our results from venlafaxine and venlafaxine-vitamin C groups (23, 24). In a similar series of experiments on patients who consumed SSRI for six months, Safarinejad observed a decrease in sperm parameters, a significant increase in DNA damage (25). These discrepancies between our results and the results of previous studies can be due to the difference in types of drugs and their dosage which have been used and the fact that the current study was performed on mice. In another study, Atli and colleagues showed that using sertraline decreases the sperm concentration and morphology. In addition, it

causes a decrease in GSH level, increase in MDA and changes the level of serum LH and testosterone (26).

Although the exact mechanism of the effects of antidepressant drugs on the reproductive system has not yet been understood well, it has been suggested that neuroendocrine factors may be involved in this mechanism (27). It has been shown that increase of serotonin in humans can affect ejaculation and sperm transition. For example, serotonin causes the inhibition of dopamine by increasing the prolactin level and stimulating the activity of prolactin-releasing factors (27). Increase in prolactin, causes the spermatogenesis to suppresses the defective sperm motility and changes the sperm quality (28).

Furthermore, Weydt and colleagues, in 2011, reported a few cases of gynecomastia due to the consumption of venlafaxine and fluoxetine, which were all associated with increased prolactin levels (29). In another study, SSRIs and SNRIs declined dopamine neurotransmissions with hyperprolactinemia (30). It is known that serotonergic antidepressants can potentiate 5Hydroxytryptophan receptor-mediated stimulation of prolactin secretion (31). It is unclear that how antidepressants affect spermatogenesis or sperm motility, however, it has been suggested that it may have an influence on sperm PH or viscosity, on nitric oxide concentration, or physiologic regulator of sperm motility (20).

The high levels of ROS affect the normal sperm quality (mobility, viability, function) by interacting with lipid membrane, proteins, nuclear and mitochondria DNA. Therefore, in our study to examine the production of ROS by drugs and their effects on the lipid peroxidation and also to evaluate total capacity of antioxidant, the amounts of MDA and DPPH were measured respectively.

Our results showed that the seminal plasma DPPH, showed no significant change.

However, the result of MDA test between group I and II, indicated a significant increase. These results are in accordance with Taziki and colleagues that showed amitriptyline led to a significant increase in lipid peroxide in the hepatocytes of rats and the amount of thiobarbituric acid-reactive substance (32). In 2010, Afify and colleagues reported that amitriptyline in high dose led to liver cytotoxicity and reduced testicular function (6).

Cordero and co-workers reported the mitochondrial cytotoxicity in patients who used amitriptyline (8). In this study, using amitriptyline by depressed patients, leads to increase in lipid peroxidation and reduction in coenzyme Q 10 (membrane antioxidant) and citrate synthesis activity (8). Our finding showed that MDA test between group I and IV was not significant but indicated a significant alteration between group IV and V. In other words, the vitamin C consumption in group V could decrease the MDA level compared to group IV.

According to Abdel-Wahab, venlafaxine is an effective antioxidant against lipid peroxidation damage and increases antioxidant defense line. Also in this study, the amount of MDA decreased in the depression group that was under treatment with venlafaxine (18). The level of lipid peroxidation in brain cortex, medulla, and erythrocyte was lower in the venlafaxine group compared to the depressed group in Eren's study which can be a confirmation to our results (5). El-Din and colleagues examined the influence of duloxetine on the biochemical and immunocytochemical alteration in the testes of rats, and reported increases in MDA level and decreases in testosterone, LH and FSH levels (33).

Also, in our study observed that MDA level in group V than in group IV, was a significant decline. By considering the lower level of MDA in group V than in group IV, it can be concluded that supplemented ascorbic acid in group V improved the effects of lipid

peroxidation caused by venlafaxine. In accordance with our results, Ayinde and colleagues reported that daily consumption of supplemented ascorbic acid can decrease stress oxidative and alteration spermatogenesis in albino rats (34). We observed that seminal plasma lipid peroxidation reduced in the venlafaxine group compared to the amitriptyline group; however, no significant change was observed between these two groups.

In the current study, the use of venlafaxine along with supplemental ascorbic acid significantly decreased lipid peroxidation. Moreover, considering the results of lipid peroxidation and of similar studies done so far on the two drugs, it seems that venlafaxine has less influence on male fertility than amitriptyline due to less ROS production and less influence of drugs on sperm parameters. This study shows that in the amitriptyline-vitamin C-treated mice, the sperm motility is improved and immotile is decreased, while in the venlafaxine-vitamin C-treated mice sperm viability is improved. Ascorbic acid plays an important role in the distinction process of spermatogenesis. The consumption of antioxidants causes conflicts with ROS. ROS might destroy testicular germ cells which leads to a decrease in sperm concentration and an increase in abnormal sperm (35). It should also be considered that with increases in lipid peroxidation the sperm motility is likely to be low (36).

Griveau and colleagues showed that ROS increases the lipid peroxidation which causes a decline in sperm mobility and then a reduction in unsaturated fatty acids (37). In accordance with Griveau's results, sperm concentration in the amitriptyline group decreased, which can be attributed to the high production of ROS in this group. In our study, a comparison between the amitriptyline and venlafaxine groups showed that lipid peroxidation has been increased, but it was not statistically significant. Simultaneous consumption of supplemental ascorbic acid

and amitriptyline decreased the lipid peroxidation; however, it did not result in any significant change between the two groups. These findings further support the idea of Das who reported a negative correlation between ascorbic acid and MDA relative to the defensive role of ascorbic acid against the process of lipid peroxidation (38).

Also, in this study, there was a significant positive correlation between the seminal plasma of ascorbic acid and the concentration, motility, and normal morphology of sperm which are in accordance with the results of Thiele (39). Hence, it is recommended for patients who are suffering from depression and are willing to have offspring to use antioxidants along with antidepressants drugs to reduce the effects of oxidative stress induced by drug consumption.

### Conclusion

Our findings show that the administration of amitriptyline in mice has more detrimental effects on sperm parameters and causes more peroxidation than venlafaxine dose.

### Acknowledgments

This study was supported by Department of Biology and Anatomy, Shahid Sadoughi University of Medical Sciences, Yazd, Iran. The authors would like to express their gratitude to the biological lab staffs in research and clinical center for Infertility and Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

### Conflict of interest

The authors declare that there is no conflict of interest.

### References

1. Novío S, Núñez MJ, Amigo G, Freire-Garabal M. Effects of fluoxetine on the oxidative status of

- peripheral blood leucocytes of restraint-stressed mice. *Basic Clin Pharmacol Toxicol* 2011; 109: 365-371.
2. Hassanane MS, Hafiz N, Radwan W, El-Ghor AA. Genotoxic evaluation for the tricyclic antidepressant drug, amitriptyline. *Drug Chem Toxicol* 2012; 35: 450-455.
3. Higgins A, Nash M, Lynch AM. Antidepressant-associated sexual dysfunction: impact, effects, and treatment. *Drug Healthc Patient Saf* 2010; 2: 141-150.
4. Marchand F, Alloui A, Chapuy E, Hernandez A, Pelissier T, Ardid D, et al. The antihyperalgesic effect of venlafaxine in diabetic rats does not involve the opioid system. *Neurosci Lett* 2003; 342: 105-108.
5. Eren I, Naziroğlu M, Demirdaş A, Çelik Ö, Uğuz AC, Altunbaşak A, et al. Venlafaxine modulates depression-induced oxidative stress in brain and medulla of rat. *Neurochem Res* 2007; 32: 497-505.
6. Afify M, Abd Elmaksoud MDE, Mosa T, Elshaer M, Kotb N. Differential effects of amitriptyline treatment on testicular and liver functions in adult male rats. *New York Sci J* 2010; 3: 10-18.
7. Moreno-Fernández AM, Cordero MD, Garrido-Maraver J, Alcocer-Gómez E, Casas-Barquero N, Carmona-López MI, et al. Oral treatment with amitriptyline induces coenzyme Q deficiency and oxidative stress in psychiatric patients. *J Psychiatr Res* 2012; 46: 341-345.
8. Cordero MD, Moreno-Fernández AM, Gomez-Skarmeta JL, de Miguel M, Garrido-Maraver J, Oropesa-Ávila M, et al. Coenzyme Q 10 and alpha-tocopherol protect against amitriptyline toxicity. *Toxicol Appl Pharmacol* 2009; 235: 329-337.
9. Viola G, Miolo G, Vedaldi D, Dall'Acqua F. In vitro studies of the phototoxic potential of the antidepressant drugs amitriptyline and imipramine. *Farmacologia* 2000; 55: 211-218.
10. Aprioku JS. Pharmacology of free radicals and the impact of reactive oxygen species on the testis. *J Reprod Infertil* 2013; 14: 158-172.
11. Bennet PJ, Moatti JP, Mansat A, Ribbes H, Cayrac JC, Pontonnier F, et al. Evidence for the activation of phospholipases during acrosome reaction of human sperm elicited by calcium ionophore A23187. *Biochim Biophys Acta* 1987; 919: 255-265.
12. Pasqualotto FF, Sharma RK, Nelson DR, Thomas AJ, Agarwal A. Relationship between oxidative stress, semen characteristics, and clinical diagnosis in men undergoing infertility investigation. *Fertil Steril* 2000; 73: 459-464.
13. Gomez E, Irvine D, Aitken R. Evaluation of a spectrophotometric assay for the measurement of malondialdehyde and 4-hydroxy-alkenals in human spermatozoa: Relationships with semen quality and sperm function. *Int J Androl* 1998; 21: 81-94.
14. Esterbauer H, Schaur RJ, Zollner H. Chemistry and biochemistry of 4-hydroxynonenal, malonaldehyde and related aldehydes. *Free Radic Biol Med* 1990; 11: 81-128.
15. Geva E, Bartoov B, Zabludovsky N, Lessing JB, Lerner-Geva L, Amit A. The effect of antioxidant treatment on human spermatozoa and fertilization rate in an in vitro fertilization program. *Fertil Steril* 1996; 66: 430-434.
16. Rudrama Devi K, Kusumlatha C, Dilip Reddy K, Kumari JK, Jael M. Ascorbic acid supplementation

- prevents the Adriamycin Induced Genotoxicity in male mice. *J Pharm Biomed Sci* 2012; 17: 1-4.
17. Talebi AR, Khalili MA, Nahangi H, Abbasi A, Anvari M. Evaluation of epididymal necrospemia following experimental chronic spinal cord injury in rat. *Int J Reprod Biomed* 2007; 5: 171-176.
  18. Abdel-Wahab BA, Salama RH. Venlafaxine protects against stress-induced oxidative DNA damage in hippocampus during antidepressant testing in mice. *Pharmacol Biochem Behav* 2011; 100: 59-65.
  19. Sharma OP, Bhat TK. DPPH antioxidant assay revisited. *Food Chem* 2009; 113: 1202-1205.
  20. Hendrick V, Gitlin M, Altshuler L, Korenman S. Antidepressant medications, mood and male fertility. *Psychoneuroendocrinology* 2000; 25: 37-51.
  21. Tanrikut C, Schlegel PN. Antidepressant-associated changes in semen parameters. *Urology* 2007; 69: 185.e5-7.
  22. Tanrikut C, Feldman AS, Altemus M, Paduch DA, Schlegel PN. Adverse effect of paroxetine on sperm. *Fertil Steril* 2010; 94: 1021-1026.
  23. Levin RM, Amsterdam JD, Winokur A, Wein AJ. Effects of psychotropic drugs on human sperm motility. *Fertil Steril* 1981; 36: 503-506.
  24. Lu S, Zhao Y, Hu J, Li X, Zhang H, You L, et al. Combined use of phosphodiesterase-5 inhibitors and selective serotonin reuptake inhibitors for temporary ejaculation failure in couple undergoing assisted reproductive technologies. *Fertil Steril* 2009; 91: 1806-1808.
  25. Safarinejad MR. Sperm DNA damage and semen quality impairment after treatment with selective serotonin reuptake inhibitors detected using semen analysis and sperm chromatin structure assay. *J Urol* 2008; 180: 2124-2128.
  26. Atli O, Baysal M, Aydogan-Kilic G, Kilic V, Ucarcan S, Karaduman B, et al. Sertraline-induced reproductive toxicity in male rats: evaluation of possible underlying mechanisms. *Asian J Androl* 2017; 19: 672-679.
  27. Tanaka E, Baba N, Toshida K, Suzuki K. Evidence for 5-HT<sub>2</sub> receptor involvement in the stimulation of preovulatory LH and prolactin release and ovulation in normal cycling rats. *Life Sci* 1993; 52: 669-676.
  28. Nargund VH. Effects of psychological stress on male fertility. *Nat Rev Urol* 2015; 12: 373-382.
  29. Weydt P, Schönfeldt-Lecuona CJ, Gahr M, Connemann BJ. Hypogonadism and gynecomastia with duloxetine. *Pharmacopsychiatry* 2011; 44: 77.
  30. Kaufman KR, Podolsky D, Greenman D, Madraswala R. Antidepressant-selective gynecomastia. *Ann Pharmacother* 2013; 47: e6.
  31. Meltzer H, Bastani B, Jayathilake K, Maes M. Fluoxetine, but not tricyclic antidepressants, potentiates the 5-hydroxytryptophan-mediated increase in plasma cortisol and prolactin secretion in subjects with major depression or with obsessive compulsive disorder. *Neuropsychopharmacology* 1997; 17: 1-11.
  32. Taziki S, Sattari MR, Egbal MA. Evaluation of amitriptyline-induced toxicity in freshly isolated rat hepatocytes and the protective role of taurine. *Life Sci J* 2013; 10: 314-320.
  33. El-Din SB, Abd-El Aty OA. Biochemical and immunocytochemical studies of the testicular changes after treatment with duloxetine hydrochloride and the possible protective effects of omega 3 in adult rat model of depression. *AAMJ* 2012; 10 (Suppl.): 264-298.
  34. Ayinde OC, Ogunnowo S, Ogedegbe RA. Influence of Vitamin C and Vitamin E on testicular zinc content and testicular toxicity in lead exposed albino rats. *BMC Pharmacol Toxicol* 2012; 13: 17-24.
  35. Vijayprasad S, Bb G, Bb N. Effect of vitamin C on male fertility in rats subjected to forced swimming stress. *J Clin Diagn Res* 2014; 8: HC05-HC8.
  36. Shabaniyan S, Farahbod F, Rafeian M, Ganji F, Adib A. The effects of Vitamin C on sperm quality parameters in laboratory rats following long-term exposure to cyclophosphamide. *J Adv Pharm Technol Res* 2017; 8: 73-79.
  37. Griveau JF, Dumont E, Renard P, Callegari JP, Le Lannou D. Reactive oxygen species, lipid peroxidation and enzymatic defence systems in human spermatozoa. *J Reprod Fertil* 1995; 103: 17-26.
  38. Das UB, Mallick M, Debnath JM, Ghosh D. Protective effect of ascorbic acid on cyclophosphamide-induced testicular gametogenic and androgenic disorders in male rats. *Asian J Androl* 2002; 4: 201-207.
  39. Thiele JJ, Freisleben HJ, Fuchs J, Ochsendorf FR. Ascorbic acid and urate in human seminal plasma: determination and interrelationships with chemiluminescence in washed semen. *Hum Reprod* 1995; 10: 110-115.