

# Hyperhomocysteinemia in Pakistani women suffering from unexplained subfertility

Seema Bibi<sup>1</sup> M.B.B.S., F.C.P.S., Mohammad Ali Pir<sup>2</sup> M.B.B.S., M.P.H., Roshan Ara Qazi<sup>2</sup> M.B.B.S., F.C.P.S., Misbah Bibi Qureshi<sup>3</sup> M.Phil.

1 Department of Obstetrics and Gynaecology, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan.

2 Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan.

3 University of Sindh, Jamshoro, Pakistan.

Received: 7 October 2009; accepted: 17 March 2010

## Abstract

**Background:** Hyperhomocysteinemia (hhcy) has been considered as a risk factor for several obstetrical complications such as early pregnancy loss, pre-eclampsia and IUGR. Recently its association with infertility has been underscored in IVF failures; however limited information is available about the relationship of hhcy and subfertility.

**Objective:** To find out the association between unexplained subfertility and hhcy in Pakistani women.

**Materials and methods:** This observational study was conducted in Department of Obstetrics and Gynaecology, Liaquat University Hospital Hyderabad from 1<sup>st</sup> April 2008 to 31<sup>st</sup> March 2009. Study group consisted of all those women who were subfertile for more than one year, have body mass index less than 25, regular menstrual cycle, normal pelvic examination findings and no past history of pelvic inflammatory disease. Exclusion criteria was male factor subfertility, endocrine and ovulatory dysfunction and tubal blockage. Evaluation was done by semen analysis, pelvic ultrasound scan, hysterosalpingography and hormonal assays. Fasting serum levels of homocysteine were determined using a fluorescence polarization immunoassay.

**Results:** In total, 61 subjects were enrolled in the study including 49 subfertile women and 12 healthy women. Among subfertile women, 39 (80%) were suffering from primary subfertility while 10 (20%) were complaining of secondary subfertility. Majority of the subjects were young, house wives and residents of Hyderabad city. Mean serum fasting homocysteine levels were significantly higher in women suffering from unexplained subfertility as compared to controls ( $12.8 \pm 5.1$  versus  $9.7 \pm 1.7$ ,  $p$ -value = 0.04).

**Conclusion:** Hyperhomocysteinemia was observed in women suffering from unexplained subfertility. However large scale clinical studies are required to confirm the association.

**Key words:** Hyperhomocysteinemia, Female infertility, Subfertility.

## Introduction

Homocysteine (hcy) is a sulphur containing amino acid which is not present in naturally

### Corresponding Author:

Seema Bibi, Department of Obstetrics and Gynaecology, Liaquat University of Medical and Health Sciences, Jamshoro, Pakistan.

**Email:** sobgsbib@cpsp.edu.pk

proteins. It is derived from demethylation of methionine, requiring folate, vitamin B<sub>6</sub> and B<sub>12</sub> as enzymatic co-factors. Malnutrition or malabsorption of folate and/or vitamin B<sub>6</sub> and B<sub>12</sub> or inherited enzymatic defects such as methylenetetrahydrofolate reductase (MTHFR) or Cystathionine B-synthetase (CBS) deficiency leads to raised hcy levels (1-3). Hyperhomocysteinemia (hhcy) has been associated with several ageing related atherosclerotic, thrombo-embolic and

neurodegenerative disorders (4-6). In the field of Obstetrics and Gynaecology, it has been underlined as risk factor for adverse pregnancy outcome like early pregnancy loss, neural tube defects, pre-eclampsia, abruptio placentae and intrauterine growth restriction (7, 8). Researchers also pointed towards the harmful effects of hhcy on female fertility. Exact role of hhcy in the earlier stages of reproductive physiology and in related diseases including subfertility is not clear. Defective follicular development, impaired chorionic villous vascularization, implantation failure and harsh uterine environment were the proposed reasons (9-12). However, limited data is available and mostly focused on women with IVF failures.

There are reports of folic acid, vitamin B6 and vitamin B<sub>12</sub> related hhcy in Pakistani population (13). Therefore this pilot study was planned to know the baseline data regarding the association of hhcy with subfertility in Pakistani women.

## Materials and methods

This observational study was conducted from 1<sup>st</sup> April 2008 to 31<sup>st</sup> March 2009 in the Department of Obstetrics and Gynaecology, Liaquat University (LUH) Hyderabad. It is the tertiary referral hospital, with 1385 beds, catering the need of both urban and rural population of Sindh province within the radius of approximately 250Km. During the study, about 45,000 women attended the out patient department for various obstetrical and gynecological problems, including 1152 subfertile women. The sample size was calculated by taking the frequency of unexplained subfertility as 7% in Pakistani population (14). About 61 participants were enrolled for the study, after informed consent. The cases were 49 subjects suffering from unexplained subfertility and 12 women were taken as controls. Control group consisted of healthy women with one or more successful pregnancies, without any obstetrical or medical problems like hypertension or Diabetes mellitus.

Study group consisted of all those women who were subfertile for more than one year, have Body mass index (BMI) less than 25, regular menstrual cycle, normal pelvic examination findings and no past history of pelvic inflammatory disease. Exclusion Criteria were male factor subfertility, endocrine and ovulatory dysfunction and tubal blockage. It was not possible to omit folic acid and vitamin B users from the study, as infertile women attending to our university hospital were initially managed by primary healthcare physicians who

prescribe these drugs routinely for every subfertile women. Evaluation was done by semen analysis, pelvic ultrasound scan, hystero-salpingography and hormonal assays i.e. serum follicle stimulating hormone, luteinizing hormone, thyroid stimulating hormone and serum prolactin. Venous blood samples were collected in fasting state and serum levels of homocysteine were determined using a Fluorescence polarization immunoassay on the IMx analyzer (Abbott Laboratory Pakistan Ltd) in the Research and Diagnostic Laboratory of Liaquat University of Medical and Health Sciences, Jamshoro. Serum levels of <12 μmol/ liter were considered normal, according to DACH LIGA homocysteine classification (15).

## Statistical analysis

Data was entered and analyzed in SPSS version 16. Student's 't' test was used to compare the mean ± standard deviation. P-value of ≤0.05 was considered statistically significant.

## Results

In total, 61 subjects were enrolled for the study, including 49 subfertile women and 12 healthy women. Among the subfertile women, 39 (80%) were suffering from primary subfertility while 10 (20%) were complaining of secondary subfertility. Demographic profile of both the groups is shown in table I.

Majority of the subjects were young, residents of Hyderabad city and house wives. Half of the subjects were literate. Mean fasting serum homocysteine level was significantly higher (p-value = 0.04) in cases (12.822±5.1823) compare with this level in controls (9.735 ± 1.7961).

**Table I.** Demographic profile (n=61).

Variables	Cases (n=49)	Controls (n=12)	p-value
Age *	26.9 ± 5.59	32.2 ± 8.24	0.01
BMI*	24.188±4.7	25.011±1.9	0.56
Duration of marriage*	6.6 ± 5.29	4.5 ± 1.88	0.19
Residence**			
Rural	12 (24.5%)	02 (16.7%)	0.71
Urban	37 (75.5%)	10 (83.3%)	
Education**			
Literate	25 (51%)	5 (41.7%)	0.74
Illiterate	24 (49%)	7 (58.3%)	
Occupation**			
House wife	48 (98%)	10 (83.3%)	0.09
Working	01 (2%)	02 (16.7%)	

BMI: Body mass index.

\* Results are expressed as mean ± standard deviation.

\*\* Results are expressed as frequency and percentage.

## **Discussion**

The study revealed raised serum fasting homocysteine levels in Pakistani women suffering from unexplained subfertility as compared to healthy subjects. Findings of our study support the preliminary work of D'Uva et al on the involvement of hcy metabolism in female reproduction (16). Their study revealed raised mean hcy levels of  $21.05 \pm 8.78$   $\mu\text{mol/liter}$  in 20 women with unexplained sterility as compared to controls. Mean Serum hcy levels of  $12.822$   $\mu\text{mol/liter}$  observed in 49 subfertile women in our study were comparatively less from the above cited study, which might be due to intake of folic acid by the subfertile women in the last three months. Reduction of serum hcy levels by intake of folate and vitamin B6 were also reported by the Korean study where serum hcy levels in infertile women, negatively correlated with total intake of vitamin B<sub>12</sub>, vitamin B<sub>6</sub> and folate (17).

Exact impact of hcy metabolism on female reproductive function is not clear. However several mechanisms were suggested to induce cellular dysfunction secondary to hhcy i.e. thrombophilia, reduced cell division, inflammatory cytokine production, altered nitric oxide metabolism, oxidative stress, apoptosis and defective methylation reaction (18). Studies have suggested that exposure of ovum to high hcy concentration may have deleterious effects on fertilization and early embryogenesis. Raised hcy levels in ovarian follicular fluid were associated with poor quality of embryo and may influence pregnancy outcome following natural or in vitro-fertilization (9, 10). Thrombosis of early decidual or chorionic vessels during early period of pregnancy might be responsible for implantation failure. Qublan and associates, in their study on the role of acquired / inherited thrombophilia in recurrent IVF implantation failure, found raised hcy levels in 60% of women with C677T MTHFR mutation (11).

In a recent study on apparently healthy Pakistani subjects, deficiencies of folate, vitamin B<sub>6</sub> and vitamin B<sub>12</sub> were found to be 39.7%, 52.8% and 6.8% respectively. Hhcy was found in 57.2% and it was negatively correlated with serum levels of vitamin B<sub>12</sub>, B<sub>6</sub> and folate (13). Prolonged deficiency may cause changes in ovulation or defective implantation leading to infertility (19). Deficiency of these vitamins may be attributed to poverty, high prevalence of intestinal parasitic infections and low intake of fresh fruits and vegetables (20).

Intervention studies have shown that supplementation with folate, vitamin B<sub>12</sub> and B<sub>6</sub> can lower hcy concentration (21, 22). The Nurses Health study II, on 18,555 participants found that regular use of multivitamins including B-vitamins and folic acid reduces the risk of ovulatory infertility (23). Overcoming micronutrient deficiency in women of reproductive age would be a good option for reducing the problem of subfertility at low cost and short time in a developing country like Pakistan.

Despite the limitation of small sample size, this study not only gave insight about the role of hcy in female reproduction but also points towards the hidden deficiency of vitamin B<sub>12</sub>, vitamin B<sub>6</sub> and folate in Pakistani women. On the basis of this pilot study, large scale community based studies can be planned to assess the prevalence and causes of vitamin B<sub>12</sub>, B<sub>6</sub> and folate deficiency and associated hhcy and its consequences on women of reproductive age group, particularly subfertility.

Mass supplementation by fortification of staple food with B-vitamins may prevent micronutrient deficiency related health problems, as many underdeveloped countries have adopted this strategy (24). There is also a need to create awareness in general population about the advantages of healthy and hygienic food eating habits.

## **Conclusion**

Serum fasting hcy levels were found to be raised in young, apparently healthy women who were suffering from subfertility as compared to controls. There is a need to plan further clinical studies on large scale to understand the association of hhcy with unexplained subfertility, along with hcy lowering effect of vitamin B<sub>6</sub>, vitamin B<sub>12</sub> and folate. Promotion of regular use of B-vitamin and folate by women of reproductive age will be cost effective strategy for the eradication of micronutrient deficiency related health problems including subfertility.

## **Acknowledgement**

We thank Ms. Afroz Nizamani and Dr. Shazia Makhdoom for data management and sample collection. The technical support of Mr. Adnan Khan of Research and Diagnostic Laboratory, Liaquat University of Medical & Health Sciences, Jamshoro, Pakistan is highly appreciated. We also thank Abbot Lab. (Pak) Ltd. for providing the equipments (IMx System).

## References

- Jacobson DW. Homocysteine and Vitamins in cardiovascular disease. *Clin chem* 1998; 44: 1833-1843.
- Refsum H, Nurk E, Smith D, Ueland PM, Gjesdal CG, Bjelland I, et al. The Hordaland Homocysteine Study. A community Based study of Homocysteine, Its Determinants, and Associations with Disease. *J Nutr* 2006; 136: S1731– 1740.
- Kang SS, Zhou J, Wong PWK, Kowalysyn J, Strokosch G. Intermediate Homocysteinemia: A Thermolabile Variant of Methylenetetrahydrofolate Reductase. *Am J Hum Genet* 1998; 43: 414-421.
- Medina MA, Amores-Sanchez MI. Homocysteine: an emergent cardiovascular risk factor? *Eur J Clin Invest* 2006; 30, 754-762.
- Herrmann W, Knapp JP. Hyperhomocysteinemia: a new risk factor for degenerative diseases. *Clin Lab* 2002; 48: 471-481.
- Markis M. Hyperhomocysteinemia and thrombosis. *Clin Lab Haem* 2000; 22: 133-143.
- Stegers – Theunissen RP, Van Iersel CA, Peer PG, Nelen WL, Steegers EA. Hyperhomocysteinemia, Pregnancy Complications and the Timing of investigation. *Obstetrics and Gynecology* 2004; 104:336-343.
- Nelen WL, Blom HJ, Steegers EA, Heijer MD, Thomas CMG, Eskes TK. Homocysteine and Folate levels as Risk Factors for Recurrent Early Pregnancy loss. *Obstet Gynecol* 2000; 95: 519-524.
- Boxmeer JC, Macklan NS, Lindemans J, Beckers NGM, Eijkemans MJC, Laven JSE, et al. IVF outcomes are associated with biomarkers of the homocysteine pathway in monofollicular fluid. *Hum Reprod* 2009; 24:1059-1066.
- Jerzak M, Putowski L, Baronowski W. Homocysteine level in ovarian follicular fluid or serum as a predictor of successful fertilization. *Ginacol Pol* 2003; 74: 949-952.
- Qublan HS, Eid SS, Ababneh HA, Amarin ZO, Smadi AZ, Al-Khafaji FF, Khader YS. Acquired and inherited thrombophilia: implication in recurrent IVF and embryo transfer failure. *Hum Reprod* 2006; 21: 2694-2698.
- Guzman MA, Navarro MA, Carnicer R, Sarria AF, Acin S, Arnal C, Muniesa P, Surra JC, Arbones – Mainer JM, Maeda N, Osada J. Cytathionine B. Synthase is essential for female reproductive function. *Hum Mol Genet* 2006; 15:3168-3176.
- Iqbal MP, Lindblad BS, Mehboobali N, Yousuf FA, Khan AH, Iqbal SP. Folic Acid and vitamin B6 deficiencies related hyperhomocysteinemia in apparently healthy Pakistani adults; is mass micronutrient supplementation indicated in this population? *JCPSP* 2009; 19: 308-312.
- Rahim R, Majid SS. Aetiological factors of infertility. *J Postgrade Med Inst* 2004; 18:166-71.
- Stanger O, Herrmann W, Pietrzik K, Fowler B, Giesel J, Dierkes J, et al. DACH-LIGA homocysteine (German, Austrian and Swiss homocysteine study): consensus paper on the rational clinical use of homocysteine, folic acid and B vitamins in cardiovascular and thrombotic diseases: guidelines and recommendations. *Clin Chem Lab Med* 2003; 41:1392 -1403.
- D' Uva M, Micco PD, Strina I, Alviggi C, Iannuzzo M, Ranieri A, et al. Hyperhomocysteinemia in Women with explained sterility or recurrent early pregnancy loss from southern Italy: a Preliminary report. *Thrombosis Journal* 2007; 5:10. doi:10.1186/1477-9560-5-10.
- Lim MY, Nam YS, Kim SS, Chang NS. Vitamin B status and Serum Homocysteine levels in infertile women. *Korean J Nutr* 2004; 37: 115-122.
- Forges T, Monnier – Barbarino P, Alberto JM, Guent – Rodriguez RM, Dava JL, Guent JL. Impact of folate and homocysteine metabolism on human reproduction health. *Hum Reprod Update* 2007; 13: 225-238.
- Bennett M. Vitamin B<sub>12</sub> deficiency, infertility and recurrent fetal loss. *J Reprod Med* 2001; 46:209-212.
- Siddiqui MI, Bilqees FM, Iliyas M, Perveen S. Prevalence of parasitic infections in a rural area of Karachi, Pakistan. *J Pak Med Assoc* 2002; 52: 315-320.
- Heijer MD, Brouwer IA, Bos GMJ, Blom HJ, Vander Put NMJ, Spaans AP, et al. Vitamin Supplementation Reduces Blood Homocysteine levels. A Controlled Trial in Patients with Venous Thrombosis and Helathy Volunteers. *Arterioscler Thromb Vasc Biol* 1998; 18: 356-361.
- Villa P, Perri C, Suriano R, Cucinelli F, Panunzi S, Ranieri M, et al. L-Folic Acid Supplementation in Healthy Post menopausal women: Effect on Homocysteine and Glycolipid Metabolism. *J CLIN Endocrinol Metab* 2005; 90: 4622-4629.
- Chavarro JE, Rich-Edwards JW, Rosner BA, Willet WC. Use of multivitamins and risk of ovulatory infertility. *Fertil Steril* 2008; 89:668-676.
- Steyn NP, Wolmorans P, Nel JH, Bourne HT. National fortification of staple foods can make a significant contribution to micronutrient intake of South African Adults. *Public Health Nutr* 2008; 11: 307-313.