

True unicornuate uterus- Pseudunicornuate uterus



Figure 1. Later HSG demonstrates apparently unicornuate uterus with normal left fallopian tube.



Figure 2. First hysterosalpingography (HSG) showed a apparently unicornuate uterus with normal fallopian tubes.

A 27 year old patient presented with primary infertility of 3 years' duration and also a history of myomectomy (5 years ago) was referred to our infertility clinic for investigation of infertility. The latest Hysterosalpingography (HSG) revealed an obstructed left fallopian tube with apparently a unicornuate uterus with luminal contour irregularity and normal left fallopian tube (Figure 1). Significant information in her past medical history revealed that she had another HSG two years before and her first hysterosalpingography (HSG) showed a apparently unicornuate uterus. Additional significant information in comparison with second HSG revealed that both fallopian tubes were opacified (Figure 2). In this case medical history also included hysteroscopic diagnosis of adhesion following open myomectomy at the age of 22. Comparison of previous graphies and hysteroscopic findings lead to a suggestion of pseudunicornuate uterus. Intrauterine adhesions develop after trauma to the basal layer of the endometrium. Unilateral excessive scarring of the uterus may lead to an obliteration of the uterine lumen resulting in an image that can mimic a unicornuate uterus (pseudunicornuate uterus) (1). A true unicornuate uterus should be excluded from pseudunicornuate uterus by a) horizontally oriented in its long axis due to deficient development of mullerian ducts b) smooth or regular contour c) with one tube. While pseudunicornuate uterus look like acquired lesion and cicatrisation leads to a usually irregular contour and uterus is more vertical in its long axis (2). Obtaining an accurate history, comparison of previous sonographic or laparoscopic findings, and awareness about this image of synechiae are the critical steps in differentiating a pseudunicornuate uterus from true unicornuate uterus.

Firoozeh Ahmadi M.D., Hadieh Haghighi B.Sc.

Department of Reproductive Imaging, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran.

Corresponding Author:

Firoozeh Ahmadi, Department of Reproductive Imaging, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran. P.O. Box: 19395-4644.

Email: f_ahmadi@royaninstitute.org

Tel/Fax: (+98) 21 23562446

Received: 10 August 2011

Accepted: 24 November 2011

References

1. Chavhan GB, Hira P, Rathod K, Zacharia TT, Chawla A, Badhe P, et al. Female genital tuberculosis: hysterosalpingographic appearances. *Br J Radiol* 2004; 77: 164-169.
2. Suleman A, Merchant A, Bharati H, Perna B, Badhe. Female Genital Tract Tuberculosis: A Review of Hysterosalpingographic Appearances Part2-The Uterus. *J Women's Imaging* 2004; 6: 153-159.

