

The effects of pyridaben pesticide on the DNA integrity of sperms and early in vitro embryonic development in mice

Ghodrat Ebadi Manas¹ Ph.D., Shapour Hasanzadeh² Ph.D., Golamreza Najafi² Ph.D., Kazem Parivar¹ Ph.D., Parichehr Yaghmaei¹ Ph.D.

1. Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran.
2. Department of Basic Veterinary Sciences, Histology and Embryology Sections, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

Corresponding Author:
Ghodrat Ebadi Manas,
Department of Biology, Science and Research Branch, Islamic Azad University, Tehran, Iran.
Email: ebadimanas@gmail.com
Tel/Fax: (+98) 9141460871

Received: 5 November 2012
Accepted: 20 February 2013

Abstract

Background: Pyridaben, a pyridazinone derivative, is a new acaricide and insecticide for control of mites and some insects such as white flies, aphids and thrips.

Objective: This study was designed to elucidate how pyridaben can affect the sperms' morphological parameters, its DNA integrity, and to estimate the effect of various quantities of pyridaben on in vitro fertilization rate.

Materials and Methods: In this study, 80 adult male Balb/C strain mice were used. Animals were divided into control and two test groups. Control group received distilled water. The test group was divided into two subgroups, viz, high dose (212 mg/kg/day) and low dose (53 mg/kg/day) and they received the pyridaben, orally for duration of 45 days. The spermatozoa were obtained from caudae epididymides on day 45 in all groups. Sperm viability, protamin compression (nuclear maturity), DNA double-strand breaks, and in vitro fertilizing (IVF) ability were examined.

Results: The pyridaben treatment provoked a significant decrease in sperm population and viability in epididymides. The data obtained from this experiment revealed that, the pyridaben brings about negative impact on the sperm maturation and DNA integrity in a time-dependent manner, which consequently caused a significant ($p < 0.05$) reduction in IVF capability. Embryo developing arrest was significantly ($p < 0.05$) higher in treated than the control group.

Conclusion: These results confirmed that, the pyridaben is able to induce DNA damage and chromatin abnormalities in spermatozoa which were evident by low IVF rate.

Key words: Pyridaben, In vitro fertilization, DNA damage, Mice.

This article extracted from Ph.D. thesis. (Ghodrat Ebadi Mans)

Introduction

Approximately 15-20% couples in their reproductive age suffering from infertility, out of which the male infertility contributes for about half of all this population (1). Among the variety of causes, environmental factors such as chemical agents and drugs seem to be the most important factor of infertility (2).

In recent years, reduction in fertility in male due to drop in sperm count and consequently decrease in reproductive capacity in human population has received enhanced interest. With reference to a report among men without a history of infertility, the mean sperm count declined from 113 million/ml in 1940 to 66 million/ml in 1990 (1). Furthermore, a significant decrease in mean seminal volume

from 3.40 ml to 2.75 ml was reported during the same period, indicating an even more pronounced decrease in the total sperm count (1).

A more recent study confirmed the drop in sperm counts over the past 20 years and, in addition, a decline in sperm motility as well as in the percentage of morphologically normal sperm (3). Pyridaben is the most recently introduced chemical for the control of all developmental stages of the spider mites (4, 5). Pyridaben has been studied for acute toxicity in different species. Major clinical signs observed were decreased spontaneous motor activity, abnormal gait, arched back posture, eye closing, priapism and bradypnea (6). Pyridaben is selective and stoichiometric inhibitor of complex I in the inner membrane of mitochondria (7).

When complex I is activated, mitochondrial nitric oxide synthase (mtNOS) shows considerable enzymatic activity and generate nitric oxide (NO) whereas inhabitation of complex I by pyridaben guides the mtNOS to lose ties with NO producing activity and to become a O₂ source (8, 9). Pyridaben was also highly toxic to midbrain organotypic slice and suggest the need for evaluation of the potential for these compounds to damage the dopaminergic system in animal and to determine whether there is significant human exposure to these pesticides (10).

Pyridaben is twice more potent than rotenone at inhibiting mitochondrial respiration. Pyridaben is a competitive inhibitor of DHR (dihydrorotenone) binding, indicating positive cooperativity (11, 12). The greater hydrophobicity of pyridaben may result in concentration of pyridaben in the membrane where the enzyme complex is located, resulting in enhanced inhibitory potency, therefore high potency of pyridaben causing cell death. Pyridaben causes significantly greater oxidative damage than a similar dose of rotenone (13).

Partial inhibition of complex I by pyridaben is sufficient to produce significant increase in reactive oxygen species (ROS) production (14-17). Pyridaben is a widely used miticide, binding to the PSST subunit of complex I and inhibits electron flow (18). Pyridaben has not been fully evaluated for its possible immunotoxic effect, whereas known that, it is a potent inducer of apoptosis in ST486 cells and in EW36 cells in combination with heat stress (19). In addition, inhabitation of electron flow at complexes I can generate excess ROS, and an oxidative stress condition, contributing to mitochondrial dysfunction (20-22).

It has been indicated that, pyridaben miticide inhabitation of NADH: ubiquinone oxidoreductase activity leading to the level of induced ornithine decarboxylase activity which causes anti-proliferative effect and thus anticancer action (23). The aim of this study was to evaluate the probable simultaneous effect of pyridaben on semen quality, sperms' DNA damages and in vitro fertilizing ability of sperms on time depending manner.

Materials and methods

Animals and treatment groups

This study was experimented on eighty adult Balb/C strain mice of 10 weeks age and

25±3g weight, maintained in laboratory animal housing facilities under controlled light conditions 12 hour light and 12 hour dark and temperature between 20-23°C. The animals were fed on standard mice pellet and watered ad libitum. In this study all experiments were conducted in accordance with the principles and procedures outlined by Urmia University guidance of ethical committee for research on laboratory animals.

The animals were divided in to control and test groups. The test subgroups nominated upon the dosage of pyridaben in the study as high dose (212 mg/kg) and low dose (53 mg/kg). The mice in control group received 0.2 ml distilled water. The high dose and low dose experiment groups received pyridaben at rate of 212mg/kg and 53 mg/kg through oral route (by gavage) respectively for the duration of 45 days.

Sperm collection

After running of treatment period, each mouse was sacrificed by decapitation according to recommendation of the institutional ethical committee. Spermatozoa were obtained from cauda epididimidis under a 20-time magnification provided by a stereo zoom microscope (model TL2, Olympus Co., Tokyo, Japan). Both cauda epididimides of each mouse was trimmed and minced in 1 ml HTF+4 mg/ml BSA medium pre-warmed to 37°C. After 20 minute incubation at 37°C in an atmosphere of 5% CO₂ and the grinded epididymal tissue was separated from the released spermatozoa.

Epididymal sperm count and viability assay

The cauda epididymis sperm reserves were determined using the standard hemocytometry (1). Sperm viability was analyzed using eosin-nigrosin staining technique. Ten microliters of each semen sample were placed on slide and stained with ten microliters of eosin-nigrosin. The live (non-stained), dead (red-stained heads), abnormal and morphologically immature sperms were evaluated (24).

Sperm chromatin integrity assay

Acidic aniline blue staining was used to detect chromatin defects of sperm nuclei related to their nucleoprotein content as associated with DNA. The Aniline blue (Ab) staining specifically reacts with lysine residues

in nuclear histones and reveals differences in the basic nuclear protein composition of the sperm. Histone-rich nuclei of immature sperm are rich in lysine and will consequently take up the blue stain. On the other hand, protamine rich nuclei of mature sperm are rich in arginine and cysteine and contain relatively low levels of lysine, which means they will not be stained by Ab (25). A drop of semen was spread on the glass slides and allowed to air-dry. All the smears were fixed in 3% buffered glutaraldehyde for 30 min. The slides were then stained with 5% aqueous aniline blue and mixed with 4% acetic acid (pH= 3.5) for 7 min then evaluated under light microscope.

DNA double strand breaks assay

A drop of semen was spread on the glass slides and allowed to air-dry. All the smears were fixed in methanol/acetic acid (3:1). The slides were then stained with 19% acridine-orange solution in phosphate citrate for 10 min in each slide. The sperms were evaluated with fluorescence microscope (Model GS7, Nikon, Japan) with a 100 oil immersion objective lens. Three types of staining patterns were identified; green sperms (double-stranded DNA), yellow and red sperms (single-stranded DNA) (26).

Collection of ovulated oocytes and in vitro fertilization

Superovulation was induced in 6-7 weeks old female mice by intraperitoneal injection of 10 IU pregnant mare serum gonadotropin (PMSG sigma, G4877) followed by intraperitoneal injection of 10 IU HCG (Sigma, C1063) 48 h later. At 12-14 hours post HCG injection, female mice were scarified by cervical dislocation. The oviducts were removed and the ampulla portion was put into a plastic dish containing HTF+ 4mg/kg BSA medium. The oocytes in cumulus masses were dissected out of the oviducts and introduced into the HTF+ 4mg/kg BSA medium. Microdrops of fertile sperm (1×10^6 sperm/ml) in HTF+ 4mg/kg BSA were prepared, and 10 to 15 oocytes were placed into each sperm microdrop (150 μ l).

The fertilization process was performed for four to six hours incubation at 37°C under 5% CO₂. The cumulus cell free fertilized oocytes were transferred to fresh drops of HTF+ 4mg/kg BSA medium for culture of embryos. All of the medium droplets were covered with

mineral oil (sigma M8410) and fertilized oocytes were evaluated by appearance of the pronuclei and polar bodies under the inverted microscope with magnification of 200 \times . After IVF, zygotes were washed 3 times with potassium simplex optimized medium (KSOM) and then transferred into fresh KSOM, cultured for an additional 5 days by incubation at 37°C under 5% CO₂. 24 hours after the zygotes culture, the two cell embryos rate as well as in vitro embryonic development were evaluated, during 5 days under phase-contrast microscopy to blastocyst stage.

Statistical analysis

Statistical analyses were performed on all data using ANOVA, by SPSS software version 16.0. All values were expressed as the mean \pm SE and $p < 0.05$ was considered to be statistically significant.

Results

Sperm count and morphology

Our results revealed that, in test group which received high dose of pyridaben, in comparison to the controls, the sperm count was reduced significantly ($p < 0.05$) after 45 days of administration (Table I). Aniline blue staining for sperm nucleus maturity revealed that, the ratio of nuclear immature sperms (light stained nucleus) was increased remarkably in pyridaben groups (low and high dose) in comparison to other test groups (Figure 1a). Furthermore the sperms with cytoplasmic droplets were observed as immature sperm. Our observations demonstrated that, the number of sperms with cytoplasmic droplets increased significantly ($p < 0.05$) in the 45th day in both pyridaben treated groups.

In the eosin-negrosin staining, those sperms with stained heads were considered as dead. Intensive sperm mortalities were observed in pyridaben groups mice with dose dependant mode (Figure 1b). It is to notify that, the severities of sperm parameters anomalies were more in high dose than the low dose exposed animals. The data for sperm count and morphology is presented in table I. The acridine-orange staining showed that, the number of sperms with double-strand DNA breaks are significantly ($p < 0.05$) intensified in the pyridaben exposed groups than the controls and this was more enhanced

after 45 days exposure (Figure1c). The numerical data for sperm DNA fragmentation and breaks is presented in table I.

The pyridaben influences the rate of fertilized oocytes

The in vitro fertilizations of oocytes by sperms which were collected from the pyridaben exposed groups animals were remarkably lower than the control group. The most considerable occurrence was that, in most cases the progression of 2-cell embryos

in to four and/or more cells embryos in pyridaben exposed groups stopped. Moreover, the majority of divisions were arrested in samples from the pyridaben exposed groups. By comparing the values for rate of fertilized oocytes, 2 cells and blastocysts between all of the groups indicated that the highest value belongs to the oocytes that exposed to sperms from the high dose of pyridaben group in the 45th day (Figure 1d). The data for in vitro fertilization rates is presented in table II.

Table I .Comparative parameters of sperm in control and experimental groups (Mean ± SE)

Different parameters	Control	Low dose	High dose
Sperm count (10 ⁶)	51.33 ± 1.45 ^a	36.67 ± 4.66 ^a	24.66 ± 3.48 ^b
Sperm viability (eosin-nigrosin) (%)	90.66 ± 1.45 ^a	71.00 ± 3.46 ^b	60.33 ± 2.72 ^b
Immature sperm (aniline blue) (%)	8.0 ± 0.58 ^a	14.66 ± 0.88 ^b	24.00 ± 4.04 ^b
DNA double strand sperm (acridine orange) (%)	9.67 ± 0.88 ^a	16.00 ± 0.57 ^b	26.00 ± 2.08 ^b

The different parameters in the control and test groups are analyzed through Tukey’s test. In each row the identical superscript indicates non-significant (P>0.05) difference between parameters.

Table II .Comparative parameters of embryos in control and experimental groups (Mean ± SE)

Different parameters	Control	Low dose	High dose
Fertilization oocyte (%)	88.34 ± 0.88 ^a	74.66 ± 3.53 ^b	48.00 ± 3.20 ^b
2- Cell embryo (%)	80.33 ± 1.45 ^a	66.65 ± 3.84 ^b	44.34 ± 5.45 ^b
Blastocyst (%)	81.34 ± 2.89 ^a	69.00 ± 1.15 ^b	40.34 ± 4.18 ^b

The different parameters in the control and test groups are analyzed through Tukey’s test. In each row the identical superscript indicates non-significant (P>0.05) difference between parameters




Figure1: (a) aniline-blue staining for spermatozoa indicate that 1- mature sperm with light nucleus and 2- immature sperm with blue nuclei. (b) eosin- nigrosin staining for spermatozoa indicate that 1- intensive sperm with light nuclei and 2- dead sperm with red nucleus (×1000). (c) acridine -orange staining for spermatozoa indicate that 1- mature sperm with light nuclei and 2- the number of sperms with double-strand DNA breaks (×1000). (d) 1- blastocyst after IVF in control group with 2- remarkably fragmented embryo (invert microscope ×1000).

Discussion

Pyridaben, a pyridazinone derivative, is a new acaricide and insecticide (4). The other compounds which are commonly used as pesticide including atrazine, benomyl, carbaryl, carbofuran, chlorpyrifos, endosulfan, dibromochloropropane (DBCP), 2, 4-dichlorophenoxy acetic acid, dimethoate, dioxin, hexachlorocyclohexane (HCH), lindane, malathion, mancozeb, methoxychlor, methyl parathion, organophosphate, imidaclopride and pyrethroid. The reproductive toxic effects of these compounds are widely studied (27, 28).

At present pyridaben is widely used as miticide and insecticide through all over the world, although this compound is toxic for different biological systems (4, 23, 29). Its noxious effects on the male and female reproductive systems are ill known. In the present study we showed that in pyridaben groups (high and low dose) sperm protamine-histone transition impairments, DNA missed integrity, and consequentially loss of motility increases. Ultimately our results showed that the quality of sperm content become lower and exhibit lower in vitro fertilization rate. In present study we used the effect of various pyridaben doses on maturation and DNA integrity and to identify in vitro fertilization ability of these sperms in different groups.

In the current study after 45 days, pyridaben group showed high morphologically immature sperms. A numbers of studies have suggested that the presence of spermatozoa with damaged DNA may be the result of an impaired chromatin packing or may be indicative of apoptosis (30, 31). Protamines may act as protective elements by sequestering metals capable of promoting the fragmentation of sperm DNA (32). Our special staining for protamine showed that by the time after pyridaben group, the intensive damage affected the sperms protamine-histone transition and this disorder may account in part for the extensive DNA damage observed in poorly packaged spermatozoa of the pyridaben group.

On the other hand the results from acridine-orange stainings were in good accordance with these findings and showed an elevated sperm DNA fragmentation and damage pyridaben rats. There are reports indicating that any disorder which resulted in a failure in epididymal sperm maturation, causes impaired sperm fertilizing ability (31,

33, 34). Development of spermatozoal ability to expose forward motility, undergoing capacitation, and penetrate the zona pellucida of the oocyte is examples of the several important properties, which the spermatozoa acquire during epididymal sperm passage (1). Our observations revealed that in pyridaben group morphologically immature sperms increased remarkably and the results from IVF correlated reversely with these findings. The ability of the embryo to survive appears to be negatively correlated with the level of DNA fragmentation in the germ line (35).

Previous studies showed that DNA-damaged sperms cannot fertilize the oocyte (31, 32). Our results in this study showed that by using the sperms from pyridaben rats some of the fertilized oocytes stopped to continue division in two cells and blastocyst embryo phase. Moreover, fertilization rate in treatment group compared with the control group was found remarkably low.

Conclusion

These results demonstrated that pyridaben is able to induce DNA damage and chromatin abnormalities in spermatozoa which could be contributed in observed low fertilizations rate.

Acknowledgments

This study was financially supported by the Research Council of Science and Research Branch, Islamic Azad University, Tehran. The authors would like to express their appreciations to staff members of the Department of Biology of this center for their modest cooperation and support of this research. We wish to thank Mr. Ali Karimi and staff of Histology and Embryology Laboratories of veterinary faculty for their very kind helps in laboratory works.

Conflict of interest

Authors do not have any conflict of interest.

References

1. Suzuki N, Sofikitis N. Protective effects of antioxidants on testicular functions of varicocele rats. *Yonago Acta medica* 1999; 42: 87-94.
2. Schlegel PN, Chagnac TS, Marshall FF. Antibiotics: potential hazards to male fertility. *Fertil Steril* 1991; 55: 235-242.
3. Agarwal A, Said TM. Role of sperm chromatin abnormalities and DNA damage in male infertility. *Hum Reprod Update* 2003; 9: 331-345.

4. Hirata K, Kawamura Y, Kuno M, Igarashi H. Development of a new acaricide pyridaben. *J Pest Sci* 1995; 20: 177-179.
5. Stumpf N, Nauen R. Cross-resistance, Inheritance, and biochemistry of mitochondrial electron transport inhibitor-acaricide resistance in *Tetranychus urticae*. *J Econ Entomol* 2001; 94: 1577-1583.
6. Hajime I, Satoru S. Summary of Toxicity Studies with pyridaben: Regulatory agricultural division. Nissan chemical industries Ltd.; 1994.
7. Gomez C, Bandez MJ, Navarro A. Pesticides and impairment of mitochondrial function in relation with the parkinsonian syndrome. *Front Biosci* 2007; 12: 1079-1093.
8. Parihar MS, Nazarewicz RR, Kincaid E, Bringold U, Ghafourifar P. Association of mitochondrial nitric oxide synthase activity with respiratory chain complex I. *Biochem Biophys Res Commun* 2008; 366: 23-28.
9. Parihar MS, Parihar A, Villamena FA, Vaccaro PS, Ghafourifar P. Inactivation of mitochondrial respiratory chain complex I lead mitochondrial nitric oxide synthase to become pro-oxidative. *Biochem Biophys Res Commun* 2008; 367: 761-767.
10. Sherer TB, Richardson JR, Testa CM, Seo BB, Panov AV, Yagi T, et al. Mechanism of toxicity of pesticides acting at complex I: relevance to environmental etiologies of Parkinson's disease. *J Neurochem* 2007; 100: 1469-1479.
11. Okun JG, Lümnen P, Brandt U. Three classes of inhibitors share a common binding domain in mitochondrial complex I (NADH: ubiquinone oxidoreductase). *J Biol Chem* 1999; 274: 2625-2630.
12. Schuler F, Casida JE. The insecticide target in the PSST subunit of complex I. *Pest Manag Sci* 2001; 57: 932-940.
13. Sherer TB, Betarbet R, Greenamyre JT. Environment, mitochondria, and Parkinson's disease. *Neuroscientist* 2002; 8: 192-197.
14. Barrientos A, Moraes CT. Titrating the effects of mitochondrial complex I impairment in the cell physiology. *J Biol Chem* 1999; 274: 16188-16197.
15. Kushnareva Y, Murphy AN, Andreyev A. Complex I-mediated reactive oxygen species generation: modulation by cytochrome c and NAD(P)⁺ oxidation-reduction state. *Biochem J* 2002; 368: 545-553.
16. Li N, Ragheb K, Lawler G, Sturgis J, Rajwa B, Melendez JA, et al. Mitochondrial complex I inhibitor rotenone induces apoptosis through enhancing mitochondrial reactive oxygen species production. *J Biol Chem* 2003; 278: 8516-8525.
17. Sipos I, Tretter L, Adam Vizi V. The production of reactive oxygen species in intact isolated nerve terminals is independent of the mitochondrial membrane potential. *Neurochem Res* 2003; 28: 1575-1581.
18. Schuler F, Yano T, Di Bernardo S, Yagi T, Yankovskaya V, Singer TP, et al. NADH-quinone oxidoreductase:PSST subunit couples electron transfer from iron-sulfur cluster N2 to quinone. *Proc Natl Acad Sci USA* 1999; 96: 4149-4153.
19. Bloom SE, Lemley AT, Muscarella DE. Potentiation of apoptosis by heat stress plus pesticide exposure in stress resistant human B-lymphoma cells and its attenuation through interaction with follicular dendritic cells: role for c-Jun N-terminal kinase signaling. *Toxicol Sci* 2006; 89: 214-223.
20. Garacia-Ruiza C, Colell A, Mari Morales A, Fernandez-Checa JC. Direct effect of ceramide on the mitochondrial electron transport chain leads to generation of reactive oxygen species. *J Biol Chem* 1997; 272: 11369-11377.
21. ST-Pierre j, Buckingham j A, Roebuck SJ, Brand MD. Topology of superoxide production from different sites in the mitochondrial electron transport chain. *J Biol Chem* 2002; 277: 44784-44790.
22. Chen, Q, Vazques EJ, Moghaddas S, Hoppel CL, Lesnefsky EJ. Production of reactive oxygen species by mitochondria: central role of complex III. *J Biol Chem* 2003; 278: 36027-36031.
23. Nianbai Fang and John E Casida. Anticancer action of cube insecticide. *Proc Natl Sci USA* 1998; 95: 3380-3384.
24. Toyoda Y, Chang MC. Fertilization of rat eggs in vitro by epididymal spermatozoa and the development of eggs following transfer. *J Reprod Fertil* 1974; 36: 9-22.
25. Nasr-Esfahani MH, Razavi S, Mardani M. Relation between different human sperm nuclear maturity tests and in vitro fertilization. *J Assist Reprod Genet* 2001; 18: 219-225.
26. Moustafa MH, Sharma RK, Thornton J, Mascha E, Abdel-Hafez MA, Thomas AJ, et al. Relationship between ROS reduction, apoptosis and DNA denaturation in spermatozoa from patients examined for infertility. *Hum Reprod* 2004; 19: 129-138.
27. Mathur N, Pandey G, Jain A GC. Review of the male reproductive toxicity. *J Herb Med Toxic* 2010; 4: 1-8.
28. Najafi GR, Razi M, Hoshyar A, Shahmohammadloo S, Feyzi S. The effect of chronic exposure with imidaclopride insecticide on fertility in mature male rats. *Int J Fertil Steril* 2010; 9: 9-16.
29. Gu QY, Chen WB, Wang LJ, Shen J, Zhang JP. Effects of sublethal dosage of abamectin and pyridaben on life table of laboratory populations of *Tetranychus turkestanii* (Acari: Tetranychidae). *Acta Entomologica Sinica* 2010; 53: 876-883.
30. Sakkas D, Mariethoz E, Manicardi G. Origin of DNA damage in ejaculated human spermatozoa. *Rev Reprod* 1999; 4: 31-37.
31. Duru NK, Morshedi M, Oehninger S. Effects of hydrogen peroxide on DNA and plasma membrane integrity of human spermatozoa. *Fertil Steril* 2000; 74: 1200-1207.
32. Bianchi P, Manicardi GC, Bizzaro D, Bianchi U, Sakkas D. Effect of DNA protamination on fluorochrome staining and in situ nicktranslation of murine and human mature spermatozoa. *Biol Reprod* 1993; 49: 1038-1043.
33. Fujishiro JA. Effect of varicocele on fertility potential: comparison between impregnating and non-impregnating groups. *Arch Androl* 1995; 35: 143-148.
34. Kessopoulou E, Tomlinson MJ, Barratt CL, Bolton AE, Cooke ID. Origin of reactive oxygen species in human semen spermatozoa or leukocytes. *J Reprod Fertil* 1992; 94: 463-470.
35. Evenson DP, Jost LK, Marshall D, Zinaman MJ, Clegg E, Purvis K, et al. Utility of the sperm chromatin structure assay as a diagnostic and prognostic tool in the human fertility clinic. *Hum Reprod* 1999; 14: 1039-1049.