

The prevalence of, and risk factors for, mycoplasma genitalium infection among infertile women in Ibadan: A cross-sectional study

Tinuade Adesola Ajani¹ FWACP, Timothy A. Olusesan Oluwasola² FWACS, Mustapha A. Ajani³ FWACP, Rasheed Ajani Bakare^{1, 4} FWACP

1. Department of Medical Microbiology, University College Hospital, Ibadan, Nigeria.
2. Department of Obstetrics and Gynaecology, University College Hospital, Ibadan and College of Medicine, University of Ibadan, Ibadan, Nigeria.
3. Department of Histopathology, Babcock University, Ilesha-Remo, Ogun State, Nigeria.
4. Department of Medical Microbiology, College of Medicine, University of Ibadan, Ibadan, Nigeria.

Corresponding Author:
Timothy A. Olusesan Oluwasola,
Department of Obstetrics and Gynaecology, University College Hospital, Ibadan, Nigeria.
Email: sesanoluwasola@yahoo.com
Tel: (+234) 8033384064

Received: 10 December 2016
Revised: 25 March 2017
Accepted: 2 August 2017

Abstract

Background: The association of mycoplasma genitalium (*M. genitalium*) with infertility has been documented. The infections are asymptomatic and difficult to diagnose. Understanding the associated risk factors will help in facilitating better screening measures for at-risk groups.

Objective: The aim was to determine the prevalence of, and risk factors for, *M. genitalium* infection among infertile women in Ibadan.

Materials and Methods: In this cross-sectional study, 402 women (267 infertile and 135 fertile) referred to 2 hospitals in Ibadan between March and November 2015 were enrolled. Information was obtained, using structured questionnaire, on sociodemographic and behavioral characteristics of the respondents while endocervical swabs were obtained for detection of *M. genitalium*. *MgPa* gene was diagnosed using the conventional Polymerase chain reaction. Bands corresponding to 495kb were documented as positive for *M. genitalium*.

Results: Among the infertile women, 43 (16.1%) had evidence of *M. genitalium* infection as against 3 (2.2%) of women without infertility ($p < 0.001$). Associated risk factors included having more than one lifetime sexual partner (OR=10.13, 95% CI: 3.76-33.97); husbands having other sexual partners (OR=12.88, 95% CI: 2.08-90.63); being a serial monogamist (OR=6, 95% CI: 4.35-8.27) and low socio-economic status (OR=2.80, 95% CI: 1.28-6.10). No relationship exists between the previous history of sexually transmitted infections and *M. genitalium*.

Conclusion: The risk factors for *M. genitalium* infection are similar to those peculiar to other sexually transmitted infections. Its routine screening should be incorporated into the current protocol for microbiological evaluation of infertile women.

Key words: Infertility, *Mycoplasma genitalium*, Prevalence, Risk factors, Screening.

Introduction

The current global infertility rate has been reported to be between 15-20% with the associated major economic burden on healthcare industry (1, 2). *Mycoplasma genitalium* (*M. genitalium*), an emerging sexually transmitted pathogen of the Mycoplasmatacea family, has been described as a major cause of various inflammatory conditions of the genital tract in women and non-gonococcal urethritis in men (3-5). The asymptomatic carrier state of *M. genitalium* is a major epidemiological concern as it enhances unnoticed transmission of the pathogen among sexual partners (6). In addition, challenges of culture and isolation of

M. genitalium contribute to under-investigation, misdiagnosis and inadequate treatment of patients with the pathogen (7-9).

In Kenya, Cohen and colleagues reported the presence of *M. genitalium* in the cervix and/or endometrium of 16% of women with histological diagnosis of acute endometritis compared with 2% in women without endometritis (10). In a related study, Simms and co-workers linked *M. genitalium* to pelvic inflammatory disease (PID), where 13% of women with clinically diagnosed PID were *M. genitalium* positive on Polymerase chain reaction (PCR) compared with 0% in the control group. Similarly, another study in the United States by Haggerty and colleagues also reported a prevalence of 14% among

patients with clinically diagnosed PID (11, 12). The evidences for *M. genitalium* cervicitis and PID suggested that the organism has potential to cause ascending infections thus resulting in infertility (11-14). Currently, in Nigeria, there is a dearth of studies on the overall prevalence and/or risk factors for acquiring *M. genitalium* in the general population and especially among patients with infertility.

The main objective of this study was to determine the prevalence of *M. genitalium* among infertile women in our environment and the possible associated risk factors as this will contribute significantly to establishing preventive strategies among “at-risk” population and consequently reduce the burden of infertility in Nigeria.

Materials and methods

This was a cross-sectional study involving 267 consenting women with a clinical diagnosis of infertility, and 135 women with proven fertility, conducted between March 1st, and November 30th, 2015. Using interviewer-administered questionnaire, we obtained information on socio-demographic characteristics and on risk factors associated with *M. genitalium* infection.

Inclusion criteria were women within the reproductive age group, women with confirmed diagnosis of infertility irrespective of the type and women who were attending the postnatal clinic which served as a proof of recent delivery of live babies. The women were also willing to allow endocervical samples to be taken for diagnosis of *M. genitalium* infection. On the other hand, the exclusion criteria were women who have been diagnosed to have sexually transmitted infections (STIs) or PIDs, those who were on antibiotics or who used antibiotics within the previous 6 months and also those who were not willing to provide consent.

Ethical standards were followed in the handling, storage, and disposal of specimens. Using sterile Copan eNat cervical swabs, endocervical swabs were collected under aseptic condition for each recruited woman, preserved with eNat preservation medium and

stored at -20°C until processing. Jona Bioscience Bacteria DNA preparation kit was used for DNA extraction in accordance with the manufacturer's instruction. The extracted DNA was amplified by conventional polymerase chain reaction, PCR (Eppendorf thermal cycler; Nexus series).

Primer sequences used for *M. genitalium* MgpA (major adhesion protein) gene primer were Mgp-F 5'AAG TGG AGC GAT CAT CAT TAC TAA C-3' and Mgp-R 5' CCG TGG TTA TCA TAC CTT CTG A- 3'. PCR set up comprised of 5 µl of DNA extract, 0.40 µl of primers (forward and reverse), 10.60 µl of PCR water and 4µl of Master mix (reaction buffer B, MgCl₂, DNTP, blue and yellow dye) all in a tube. All tubes were sealed and briefly centrifuged before amplification in PCR machine. After amplification, electrophoretic separation of PCR products was performed on 1.5% agarose gel stained with ethidium bromide, and visualized by Ultraviolet illumination (15).

About 10 µl of DNA ladder at 100bp gradient was placed at one end of wells of 1.5% agarose gel stained with 2µl ethidium bromide. Thereafter, 20 µl of DNA extract was added to each well on the agarose gel, placed into the electrophoretic tank and switched on at 100 volts for 60 min. The tray containing the agarose gel and bands were transferred to a bio-imaging system and results were read on a computer (Figure 1).

Bands corresponding to 495kb on the DNA ladder were documented as positive for *M. genitalium* and those that did not meet the criteria were recorded as negative.

Ethical consideration

Written informed consent was obtained from eligible participants. Ethical approval was obtained from the Joint Ethics Committee of the University of Ibadan and University College Hospital Ibadan before commencement (UI/EC/12/0201).

Statistical analysis

Descriptive and inferential statistical analysis of the data was done using the Statistical Package for the Social Sciences,

version 20.0, (SPSS Inc, Chicago, Illinois, USA). Variables of interest included levels of education and income, past history of infection with *Neisseria gonorrhoea* or any other STIs, personal or partner's use of male barrier contraceptive device (condom), the current number of sexual partners, number of lifetime sex partners and number of alcohol intake per day. Means and standard deviations were derived for quantitative variables while proportions were derived for qualitative variables. Association between categorical variables was determined using Chi-square test at statistical significance level of set 5%. Relationship between multiple variables were further determined using the multivariate logistic regression.

Results

Our study was aimed at determining the prevalence of, and associated risk factors for *M. genitalium* in Ibadan. It was a cross-sectional, hospital-based study involving 267 infertile women with mean age of 33.8±5.7 yr out of which majority, 156 (58.4%), had secondary infertility. In addition, 135 women with proven fertility were also recruited to serve as controls and their mean age is similar to that of the infertile group. Most of the respondents were married (95.1%), were Christians (62.2%) and had a tertiary level of education (58.1%) (Table I). Overall, *M. genitalium* infection was positive in 43 (16.1%) of the infertile group and 3 (2.2%) of the controls. There was a slightly higher prevalence of *M. genitalium* among those with secondary infertility 27 (17.3%) as against 16 (14.4%) of those with primary infertility although the difference is not statistically significant.

The relationship between commonly identified factors and acquisition of *M. genitalium* infection is described in table II. Family type especially serial monogamy, monthly income less than the National minimum wage of 18,000 naira (equivalent of 50 US Dollars) implying low socioeconomic status, daily alcohol consumption and having multiple sexual partners (having more than

one-lifetime sex partners and/or have husbands who have other sex partners) are significantly associated with acquiring the pathogen. About two-thirds, 62.5%, of participants whose partners have other sexual partners have a positive result for *M. genitalium* infection. However, there was no association between *M. genitalium* and age, past history of *Neisseria gonorrhoea* and other STIs and the use of condoms.

On logistic regression analysis of factors that are independently significant for *M. genitalium* infection, it was discovered that being single, being a serial monogamist, having spouses with more than one sexual partners and having more than one lifetime sexual partners remained significant (Table III). Participants whose husbands have other sexual partners are almost 13 times at risk of acquiring *M. genitalium* infection (OR=12.88, 95% CI: 2.08-90.63) while those respondents who have more than one-lifetime sexual partner are ten times at risk of the infection (OR=10.13, 95% CI: 3.76-33.97). In addition, being a serial monogamist increases the risk by 6 (OR=6.0, 95% CI: 4.35-8.27) and those with low socioeconomic status have about triple the risk of infection (OR=2.80, 95% CI: 1.28-6.10).

Table I. Socio-demographic factors of the participants with infertility

Variables	n (%)
Age Group (yr)	
20-24	9 (3.4)
25-29	52 (19.5)
30-34	90 (33.7)
35-39	71 (26.6)
≥ 40	45 (16.9)
Marital Status	
Married	254 (95.1)
Single	12 (4.5)
Divorced	1 (0.4)
Education	
Primary	19 (7.1)
Secondary	93 (34.8)
Tertiary	155 (58.1)
Religion	
Christianity	166 (62.2)
Islam	101 (37.8)
Type of infertility:	
Primary	111 (41.6)
Secondary	156 (58.4)
Previous history of abortion:	
Yes	118 (44.2)
No	149 (55.8)

*Descriptive analysis showing frequency and percentages

Table II: Factors associated with *Mycoplasma genitalium*

Variable	Mycoplasma genitalium		χ^2	p-value
	Yes	No		
Grouped Age (yr)			3.3	0.190
20-29	14 (23.0)	47 (77.0)		
30-39	21 (13.0)	140 (87.0)		
≥40	8 (17.8)	37 (82.2)		
Family type			8.3	0.040
Single	9 (19.6)	37 (80.4)		
Monogamous	31 (16.7)	155 (83.3)		
Polygamous	2 (5.9)	32 (94.1)		
Serial monogamist	1 (100.0)	0 (0.0)		
Monthly income (in naira)			8.3	0.004
<18,000	15 (29.4)	36 (70.6)		
≥18,000	28 (13.0)	188 (87.0)		
Past history of Gonorrhoea			3.7	0.159
Yes	19 (19.0)	81 (81.0)		
No	19 (12.8)	130 (87.2)		
Don't know	5 (27.8)	13 (72.2)		
Use of condom			2.1	0.143
Yes	9 (24.3)	28 (75.7)		
No	34 (14.8)	196 (85.2)		
Husband has other sex partners			14.3	0.001
Certainly yes	5 (62.5)	3 (37.5)		
Certainly no	27 (16.6)	136 (83.4)		
Not sure	11 (11.5)	85 (88.5)		
Lifetime sexual partners			29.9	<0.001
One	5 (3.8)	128 (96.2)		
>One	38 (28.4)	96 (71.6)		
Daily alcohol consumption (in bottles)			9.0	0.011
None	38 (15.0)	215 (85.0)		
1	4 (57.1)	3 (42.9)		
>1	1 (14.3)	6 (85.7)		


Data presented as n (%).

*Chi-square test statistics were used and p<0.05 was accepted as statistically significant

Table III: Logistic regression analysis

Variables	Odds Ratio	95% CI	p-value
Family type	Monogamous (Reference)		
Single	1.22	0.49-2.95	0.64
Polygamous	0.31	0.05-1.44	0.105
Serial monogamist	6.0	4.35-8.27	0.027
Monthly income, in naira (dollar)	≥18,000 (50)-Reference		
<18,000 (50)	2.80	1.28-6.10	0.004
Husband has other sex partners	Certainly no (Reference)		
Certainly yes	12.88	2.08-90.63	0.00012
Lifetime sexual partners	One (Reference)		
>One	10.13	3.76-33.97	<0.0001

*Odds ratio was used to explain the logistic regression analysis and p<0.05 was accepted as statistically significant

**Figure 1.** Gel electrophoresis of PCR amplicons for *Mycoplasma genitalium*

Discussion

Our study, which was aimed at identifying risk factors that are associated with the acquisition of *M. genitalium* among infertile women in Ibadan, Southwest Nigeria, found a prevalence of 16.1% among patients with infertility and 2.2% among women without infertility. This difference is statistically significant and implied that *M. genitalium* is associated with infertility. Previous studies from Nigeria had reported a prevalence of 4% among asymptomatic, infertile males in South-southern part of Nigeria and 6% among asymptomatic adolescents in South-eastern Nigeria but no studies had been done among infertile women to the best of our knowledge (14, 16).

In comparison with similar studies in other environments, our prevalence is slightly higher than that of Grzeško and his colleagues but lower than that of some other researchers (4, 17-19). The observed variations may be due to differences in the assay method employed as most studies using serological assays are limited in sensitivities because of cross-reactions between the antibodies of *M. genitalium* and *M. pneumoniae*. However, Rajkumar and his colleagues, using real-time PCR, also obtained higher prevalence rate in a pilot population although with patients who have primary infertility having a much higher rate than those with secondary infertility (20). Real-time PCR has been suggested to have more detection sensitivity than conventional PCR thereby implying that, in the absence of racial differences, the prevalence rate in our environment may actually be higher than we obtained (21-23).

The risk factors that are confirmed to be associated with *M. genitalium* infection from this study included: family type (single and serial monogamy) and multiple sexual partners in which the respondents have more than one lifetime sex partners and/or have husbands who equally have other sexual partners thus corroborating the findings of previous studies (4, 24-28). Thurman and his co-workers reported that *M. genitalium* was detected in 9.5% and 10.6% of women and men in heterosexual relationships reporting to an STI clinic (27). However, these subjects were five times more likely to be infected with

M. genitalium if their sexual partner was *M. genitalium* positive. Similarly, Tosh and his colleagues also reported that despite a high prevalence rate of 13.6% for *M. genitalium* among 383 young women, only one of the respondents who tested positive for the pathogen was sexually inexperienced (28). The increase in the probability of acquisition of infection among participants whose husbands have other sexual partners was likely driven by its association with sexual activity. This therefore confirms the plausible suggestion that transmission requires repeated exposure to an infected partner—which is often facilitated by the longer-term partnerships that are characteristic of cohabiting individuals.

It is interesting to note that being single is associated with the acquisition of *M. genitalium* although this is similar to the report from a study among sex-workers in China (29). Diverse reasons have been advanced for this finding ranging from cultural reasons which shift the blame of infertility on just one partner to that of some ladies opting to be single mothers in order to avoid the troubles and skirmishes of marriage. However, our study was unable to establish any association between past infection with *Neisseria gonorrhoea* or any other STI and *M. genitalium*. This result is similar to that of Hancock *et al* but in contrast with that from Vandepitte and co-workers in a study conducted among commercial sex workers (30, 31).

However, the lack of association between the previous infection with *Neisseria gonorrhoea* or other STIs and *M. genitalium* in this study may be due to factors such as recall bias, deliberate under-reporting or asymptomatic infections. Meanwhile, the lack of association between condom use and *M. genitalium* in our study is consistent with previous studies in Africa (31, 32). This is understandable as our study participants were being managed for infertility and as such were not expected to be using barrier contraceptive methods with their partners.

Conclusion

In summary, the prevalence of *M. genitalium* is significantly higher among infertile patients and may be an important potential pathogen for female infertility among

our study population. It is important to note that the prevalence rate may still be higher than obtained if there are opportunities to use real time PCR. The risk factors found to be associated with *M. genitalium* infection lend out further support for the sexual transmissibility of *M. genitalium* while being unmarried and having multiple sexual partners have been shown to be independent risk factors for its acquisition. Understanding these risk factors will enable targeted efforts at their prevention in order to reduce the burden of infertility in Nigeria. While infections like genital tuberculosis have been proven to have no relationship with unexplained infertility, it is apparent that further studies are required to determine the extent of involvement of *M. genitalium* in the cause and course of infertility among our women (33).

Acknowledgments

We acknowledge the contribution of Copan Italia in freely supplying the flocked cervical swabs.

Conflict of interest

The authors declared no conflict of interest

References

1. Zegers-Houchschild F, Adamson GD, de Mouson J, Ishiara O, Masour R, Nygren K et al. International committee for monitoring assisted reproductive technology (ICMART) and the world health organization (WHO) revised glossary of ART terminology, 2009. *Fertil Steril* 2009; 92: 1520-1524.
2. Durgesh DG, Bajaj JK, Damle SA, Jayanti MP, Sonali D, Shilpa CK. Study of *Chlamydia trachomatis* in infertile women. *Ind J Res* 2013; 2: 260-263.
3. McGowin CL, Anderson-Smiths C. *Mycoplasma genitalium*: An emerging cause of sexually transmitted disease in women. *PLoS Pathog* 2011; 7: e1001324.
4. Grzeško J, Elias M, Maczyńska B, Kasprzykowska U, Tłaczała M, Goluda M. Occurrence of *mycoplasma genitalium* in fertile and infertile women. *Fertil Steril* 2009; 91: 2376-2380.
5. Lis R, Rowhani-Rahbar A, Manhart LE. *Mycoplasma genitalium* Infection and female reproductive tract disease: A meta-analysis. *Clin Infect Dis* 2015; 61: 418-426.
6. Manhart LE, Holmes KK, Hughes JP, Houston LS, Totten PA. *Mycoplasma genitalium* among young adults in the United States: An emerging sexually transmitted infection. *Am J Public Health* 2007; 97: 1118-1125.

7. Plečko V, Zele Starčević L, Tripković V, Rezo Vranješ V, Skerlev M. Mycoplasma genitalium: Clinical Significance and Diagnosis. *Acta Dermatovenerol Croat* 2013; 21: 236-240.
8. Razin S. Mycoplasma. In Topley & Wilson's Microbiology and Microbial Infections. Wiley-Blackwell. Available at: <http://dx.doi.org/10.1002/9780470688618.taw0077>.
9. Jombo G, Enenebeaku M, Peters E, Itam H, Mbaawuaga E. Symptomatic genital Mycoplasma infections among ante-natal women in an urban community of northern Nigeria and the need to possibly widen the scope of present laboratory investigations. *Int J Infect Dis* 2008; 7: 1-7
10. Cohen CR, Manhart LE, Bukusi EA, Astete S, Brunharm RC, Holmes KK, et al. Association between Mycoplasma genitalium and acute endometritis. *Lancet* 2002; 359: 765-766.
11. Simms I, Eastick K, Mallinson H, Thomas K, Gokhale R, Hay P, et al. Associations between Mycoplasma genitalium, Chlamydia trachomatis and pelvic inflammatory disease. *J Clin Pathol* 2003; 56: 616-618.
12. Haggerty CL, Totten PA, Astete SG, Ness RB. Mycoplasma Genitalium among women With Nongonococcal, Nonchlamydial Pelvic Inflammatory Disease. *Infect Dis Obstet Gynecol* 2006; 2006: 30184.
13. Daley GM, Russell DB, Tabrizi SN, McBride J. Mycoplasma genitalium: a review. *Int J STD AIDS* 2014; 25: 475-487.
14. Ibadin KO, Osemwenkha AP, Ibeh IN. Urogenital tract infection in asymptomatic male patients with infertility in University of Benin teaching hospital, Benin City, Edo State. *Malaysian J Microbiol* 2012; 8: 289-292.
15. Anaedobe CG, Fowotade A, Omoruyi CE, Bakare RA. Prevalence, socio-demographic features and risk factors of Hepatitis B virus infection among pregnant women in Southwestern Nigeria. *Pan Afr Med J* 2015; 20: 406.
16. Chukwuka CP, Agbakoba NR, Emele FE, Oguejiofor C, Akujobi CN, Ezeagwuna DA, et al. Prevalence of genital Mycoplasmas in the vaginal tracts of adolescents in Nnewi, south-eastern, Nigeria. *World J Med Sci* 2013; 9: 248-253.
17. Mohseni Moghadam N, Kheirkhah B, Mirshekari TR, Fasihi Harandi M, Tafsiri E. Isolation and molecular identification of mycoplasma genitalium from the secretion of genital tract in infertile male and female. *Iran J Reprod Med* 2014; 12: 601-608.
18. Clausen HF, Fedder J, Drasbek M, Nielsen PK, Toft B, Ingerslev HJ et al. Serological investigation of Mycoplasma genitalium in infertile women. *Hum Reprod* 2001; 16: 1866-1874.
19. Tomusiak A, Heczko PB, Janeczko J, Adamski P, Pilarczyk-Żurek M, Strus M. Bacterial infections of the lower genital tract in fertile and infertile women from the southeastern Poland. *Ginekol Pol* 2013; 84: 352-358.
20. Rajkumar N, Sethi S, Dhaliwal L, Gupta N, Yadav R, Banga S, et al. Role of Mycoplasma genitalium in infertility and bad obstetric history in Indian non-pregnant women: a pilot study. *Int J Infect Dis* 2012; 16: e270.
21. Edberg A, Jurstrand M, Johansson E, Wikander E, Hoog A, Ahigvist T, et al. A comparative study of three different PCR assays for detection of Mycoplasma genitalium in urogenital specimens from men and women. *J Med Microbiol* 2008; 57: 304-309.
22. Müller EE, Venter JM, Magooa MP, Morrison C, Lewis DA, Mavedzenge SN. Development of a rotor-gene real-time PCR assay for the detection and quantification of mycoplasma genitalium. *J Microbiol Methods* 2012; 88: 311-315.
23. Svenstrup HF, Jensen JS, Björnelius E, Lidbrink P, Birkelund S, Christiansen G. Development of a quantitative real-time PCR assay for detection of mycoplasma genitalium. *J Clin Microbiol* 2005; 43: 3121-3128.
24. Andersen B, Sokolowski I, Østergaard L, Kjølseth Møller J, Olesen F, Jensen JS. Mycoplasma genitalium: prevalence and behavioural risk factors in the general population. *Sex Trans Infect* 2007; 83: 237-241.
25. Oakeshott P, Aghaizu A, Hay P, Reid F, Kerry S, Atherton H et al. Is mycoplasma genitalium in women the "New Chlamydia?" A community-based prospective cohort study. *Clin Infect Dis* 2010; 51: 1160-1166.
26. Svenstrup HF, Dave SS, Carder C, Grant P, Morris-Jones S, Kidd M, et al. A cross-sectional study of Mycoplasma genitalium infection and correlates in women undergoing population-based screening or clinic-based testing for Chlamydia infection in London. *BMJ Open* 2014; 4: e003947.
27. Thurman AR, Musatovova O, Perdue S, Shain RN, Baseman JG, Baseman JB. Mycoplasma genitalium symptoms, concordance and treatment in high risk sexual dyads. *Int J STD AIDS* 2010; 21: 177-183.
28. Tosh AK, Van Der Pol B, Fortenberry JD, Williams JA, Katz BP, Batteiger BE, et al. Mycoplasma genitalium among adolescent women and their partners. *J Adolesc Health* 2007; 40: 412-417.
29. Xiang Z, Yin YP, Shi MQ, Jiang N, Han Y, Wang HC, et al. Risk factors for Mycoplasma genitalium infection among female sex workers: a cross-sectional study in two cities in southwest China. *BMC Public Health* 2012; 12: 414.
30. Hancock EB, Manhart LE, Nelson SJ, Kerani R, Wroblewski JKH, Totten PA. Comprehensive assessment of sociodemographic and behavioral risk factors for mycoplasma genitalium infection in women. *Sex Transm Dis* 2010; 37: 777-783.
31. Vandepitte J, Muller E, Bukonya J, Nakubulwa S, Kyakuwa N, Buvé A, et al. Prevalence and correlates of Mycoplasma genitalium infection among female sex workers in Kampala, Uganda. *J Infect Dis* 2012; 205: 289-296.
32. Pépin J, Labbe AC, Khonde N, Deslandes S, Alary M, Dzokoto A, et al. Mycoplasma genitalium: an organism commonly associated with cervicitis among west African sex workers. *Sex Transm Infect* 2005; 81: 67-72.
33. Eftekhari M, Pourmasuni S, Sabeti P, Aflatoonian A, Sheikha H. Mycobacterium tuberculosis infection in women with unexplained infertility. *Int J Reprod BioMed* 2015; 13: 749-754.