



Research Article

Comparative gene expression analysis of *Fas* and related genes in preeclamptic and healthy women: A cross-sectional study

Zaima Ali^{1, 2} M.B.B.S., M Phil., Saba Khaliq¹ M.Sc., Ph.D., Saima Zaki³ M.B.B.S., FCPS, Hafiz Usman Ahmad¹ M.Sc., Khalid Pervaiz Lone¹ Ph.D.

¹Department of Physiology and Cell Biology, University of Health Sciences Lahore, Lahore, Pakistan.

²Department of Physiology, Lahore Medical and Dental College Lahore, Lahore, Pakistan.

³Department of Obstetrics and Gynecology, Jinnah Hospital Lahore, Lahore, Pakistan.

Corresponding Author:

Zaima Ali; Department of Physiology and Cell Biology, University of Health Sciences Lahore, Lahore, Pakistan.
Postal Code: 54000
Tel: (+92) 3244215272
Email: zaima.ali@hotmail.com

Received 8 December 2018

Revised 15 April 2019

Accepted 6 October 2019

Production and Hosting by
Knowledge E

© Ali *et al.* This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editor-in-Chief:

Aflatoonian Abbas M.D.

Abstract

Background: Preeclampsia is a hypertensive disorder of pregnancy affecting about 2-10% pregnancies worldwide. mRNA expression of tumor necrosis factor alpha (*TNF- α*), *Fas*, and *FasL* have been reported to be altered in placental bed in preeclamptic pregnancies. We hypothesized that the expression of these genes is also altered in peripheral blood mononuclear cells (PBMCs) in preeclampsia.

Objective: To compare the expression of Fas receptor and related genes in PBMCs of preeclamptic and normotensive pregnant women.

Materials and Methods: A cross-sectional comparative study comprising of 18 cases and 18 controls was designed. 5 ml of venous blood was drawn and collected considering aseptic measures. Buffy coat was separated by centrifugation and stored at -20°C. Favor Prep total RNA Isolation Kit (Favorgen, Taiwan) was used for RNA extraction. The mRNA expression of *TNF- α* , *Fas*, and *FasL* was measured by real-time polymerase chain reaction in PBMCs in preeclamptic and normal pregnancies.

Results: A significant increase in mRNA expression of *TNF- α* , *Fas*, and *FasL* ($p \leq 0.001$) was observed in PBMCs of preeclamptic pregnancies compared to the control group ($p \leq 0.001$). Moreover, a significant positive correlation was found between the *TNF- α* mRNA expression and *Fas* and *FasL* ($p \leq 0.001$).

Conclusion: The results lead to the conclusion that mRNA expression of *TNF- α* , *Fas*, and *FasL* in the maternal PBMCs is altered in preeclamptic pregnancies and might contribute to the pathogenesis of the disease.

Key words: Preeclampsia, *TNF- α* , *Fas*, Apoptosis.

OPEN ACCESS

1. Introduction

The outcome of a successful pregnancy is mainly dependent on placental sufficiency but any disorder resulting in poor placentation during pregnancy leads to preeclampsia (PE), restricted growth of the fetus and placental abruption (1, 2). PE is a gestational disorder, characterized by a rise in blood pressure and proteinuria commencing at more than 20wk of gestational age. It contributes a large share to the burden of maternal morbidity and mortality with an overall incidence of 2-10% worldwide (3). Normal placentation revolves around balanced angiogenesis and apoptosis. A number of studies have addressed disturbances in apoptosis in the placental bed in PE (4-6). *Tumor necrosis factor alpha (TNF- α)* secreted by macrophages and monocytes can induce apoptosis in vascular smooth muscle cells surrounding the spiral arteries as well as human trophoblast in the placental bed (7). Trophoblast cells secrete TNF- α and ischemia of the placenta results in increased release of this pro-inflammatory cytokine (8, 9). In normal pregnancy, there is a shift of maternal immune response from T helper 1 towards T helper 2 response (10). This shift does not occur in pregnancies complicated by PE with a resultant increase in the production of type 1 cytokines, e.g. TNF- α , IL-2, and Interferon gamma. All of these induce inflammation (11, 12).

Fas and FasL, members of TNF- α superfamily, are representative of the intrinsic pathway of apoptosis and are highly expressed on invasive extra-villous trophoblast cells as well as maternal lymphocytes. Abundant FasL expression on trophoblast was proclaimed to induce apoptosis of harmful activated maternal lymphocytes while the enhanced Fas expression on invasive trophoblast results in apoptotic death of these cells in the developing placenta (13). Many studies have reported the disturbance of the intrinsic pathway

of apoptosis with differential expression of TNF- α , Fas, and FasL in placental bed in pregnancies complicated with PE and intrauterine growth retardation (IUGR) (5, 14, 15). The emerging concept that release of placental remains into maternal blood modulates the gene expression of circulating blood cells has persuaded the researchers to study gene expression in circulating blood cells (16). Moreover, cells outside the placental bed like macrophages and monocytes have been reported to produce TNF- α (7). We hypothesized that peripheral blood mononuclear cells (PBMCs) might contribute as an extra placental source of mRNA expression of TNF- α , Fas, and FasL. Only preliminary data is available regarding this extra placental source (PBMCs) of these inflammatory and apoptotic mediators. The mRNA expression of these genes in PBMCs in PE has not been studied by real-time polymerase chain reaction (PCR) except for TNF- α (17). The selection of PBMCs as the experimental tissue using real-time PCR as the study method is novel regarding these three genes expression analysis in PE. To elucidate the role of mRNA expression of these genes, a cross-sectional comparative study was designed.

The aim of this study was to evaluate and compare the mRNA expression of TNF- α , Fas, and FasL in maternal blood PBMCs in PE and healthy pregnancies in Pakistani women.

2. Materials and Methods

2.1. Sample collection

The participants for this cross-sectional comparative study were collected from tertiary care hospital of Lahore from October 2016 to March 2017. Cases comprised of 18 pregnant women (between the ages of 18 and 40 yr) in the third trimester (28-40 wk), diagnosed as preeclamptic and 18 age-matched normal pregnant women at

the same gestational age, considered as controls. The criteria for PE were: new onset of systolic blood pressure >140 mmHg or diastolic blood pressure \geq 90 mmHg at >20 wk of gestation accompanied by 24 hr proteinuria \geq 300 mg (\geq 1+ on dipstick), in at least two random urine samples collected 4-6 hr apart. Cases were further grouped into early-onset PE (EOP, 28-32 wk) and late-onset PE (LOP, 32.1-40 wk) with nine women in each group. All the participants were non-smoker. Women with a history of diabetes mellitus, renal disease, arthritis, inflammatory bowel disease, chronic hypertension, cardiovascular illness (e.g. ischemic heart disease), or other chronic inflammatory disease were excluded. Demographic data were recorded along with complete medical, obstetric, and family history.

5 ml of venous blood was drawn and collected considering aseptic measures in EDTA-coated vacutainer. Buffy coat was separated by centrifugation and stored at -20°C within an hour of sample collection. Favor Prep total RNA Isolation Kit (Favorgen, Taiwan) was used for RNA extraction following the manufacturer's instructions. The concentrations of extracted RNA were measured using Nano-drop and stored at -80°C in RNase/DNase free water.

2.2. Quantitative real Time-PCR

Two μ g of total RNA of each sample was used to synthesize cDNA. RevertAid First-strand cDNA Synthesis kit (Thermo Scientific, USA) was used to synthesize cDNA following the manufacturer's instructions. Gene expression was quantified for 18 cases and 18 controls using synthesized cDNA and gene-specific-primers for Real-time PCR (Table I) on CFX 96 (Biorad, USA) using 2 X SYBR Green master mix (Fermentas, USA) according to the manufacturer's instructions. All reactions were carried in 20 μ l of the reaction mixture with 1 μ l of cDNA, 8 μ l of 2 X SYBR Green Real-Time PCR Master mix, 0.5 μ l of forward and reverse primers, and RNase-free water (Fermentas, USA). Real-time-PCR protocol was 94°C for 4 min, followed by 30 cycles of 94°C for 30 sec, annealing at 60°C for 30 sec, and extension at 70°C for 42 sec in a thermal cycle followed by melt curve analysis. Samples were assayed in duplicate and three housekeeping genes were used for data normalization. Relative gene expression analysis was performed using the $2^{-\Delta\Delta Ct}$ method. Confirmation of the products of the expected size was done by electrophoresis on agarose gel (Figure 1).

Table I. Sequence of primers used for quantitative real-time PCR

Gene	Sequence	PCR product length (bp)
Fas-FP	5'-CACCCCCAAACATGGAAATA-3'	186-bp
Fas-RP	5'-GGGTGGGGGAAAAATAAGAA-3'	
FasL-FP	5'-CTGGGGATGTTTCAGCTCTTC-3'	185-bp
FasL-RP	CTTCACTCCAGAAAGCAGGAC-3'	
TNFα-FP	5'-GGAGAAGGGTGACCGACTCA-3'	362-bp
TNFα-RP	5'-CTGCCAGACTCGGCAA-3'	
GAPDH-FP	5'-ACG GAT TTG GTC GTA TTG GG-3'	214-bp
GAPDH-RP	5'-CGC TCC TGG AAG ATG GTG AT-3'	
β-actin-FP	5'-TCC ACC TTC CAG CAG ATG TG-3'	75-bp
β-actin-RP	5'-GCA TTT GCG GTG GAC GAT-3'	
18s rRNA-FP	5'-AGA AAC GGC TAC CAC ATC CAA-3'	91-bp
18s rRNA-RP	5'-CCT GTA TTG TTA TTT GTC ACT ACC T-3'	




Figure 1. Confirmation of expected size products.

2.3. Ethical consideration

The study was performed in accordance with the ethical standards of the institutional ethical review (code: UHS/Education/126-16/2754) and with the Declaration of Helsinki 1964 and its later amendments. Informed written consent was obtained from all individual participants included in the study.

2.4. Statistical analysis

Statistical analysis was performed using Statistical Package for the Social Sciences, Version 22.0, Armonk, New York, USA (SPSS). The clinical parameters were expressed as means \pm SD (standard deviation). Expression of the studied genes was normalized against the mean of three housekeeping genes, that is, GAPDH, β -actin, and 18srRNA. Shapiro Wilk test was used to check the normality of data. The mRNA expression was compared using fold change values between two groups by students *t* test and One-way ANOVA with post-hoc Tukey's test for comparison between the multiple groups (EOP, LOP, and normotensive group). Differences in gene expressions were reported as fold change

with SD. Pearson's correlation was used to find the relationship between the expressions of different genes. $P \leq 0.05$ was considered statistically significant.

3. Results

3.1. Clinical characteristics

The study population consisted of 18 women with PE and 18 healthy normotensive controls. Maternal ages of the participants were 28.2 ± 5 and 25.4 ± 3 for the case and control groups, respectively ($p = 0.06$). Similarly, the difference between the gestational age at the time of sampling in the case group (32.3 ± 3) was not significant from the controls (32.2 ± 3) ($p = 0.91$).

3.2. Gene expression analysis

Gene expression between different groups was reported as fold change (Figures 2, 3). The PBMCs mRNA expressions of *Fas*, *FasL* and *TNF- α* were significantly different between the case and control groups with a marked increase in the case group ($p \leq 0.001$ for all). *FasL* showed a marked increase of

3.21-fold while the results for *TNF-α* and *Fas* were 3.08 and 2.98-fold, respectively (Table II).

For a detailed analysis of gene expression, the participants in the case group was divided into two subgroups, EOP and LOP, depending upon the time of onset of the disease. The difference in the mRNA expression of all the studied genes was not significant between the two subgroups (Table II).

3.3. Correlations between *TNF-α*, *Fas*, and *FasL* mRNA expression

mRNA expression of *TNF-α* showed a positive correlation with *Fas* expression in the case group calculated by Pearson’s correlation (Pearson’s rho 0.95, $p < 0.001$). Similarly, *FasL* was found to be positively correlated with *TNF-α* (Pearson’s rho 0.88, $p < 0.001$, Table III).

Table II. Comparison of mRNA expression between different groups

Gene	Normotensive (a)	Preeclampsia (b)	EOP (c)	LOP (d)	P-value
<i>Fas</i>	1.00	2.98 ± 0.55	2.99 ± 0.58	2.97 ± 0.55	a & b: <0.001* c & d: 0.48** a & c: <0.001** a & d: <0.001**
<i>FasL</i>	1.00	3.21 ± 0.43	3.37 ± 0.48	3.01 ± 0.28	a & b: <0.001* c & d: 0.34** a & c: <0.001** a & d: <0.001**
<i>TNF-α</i>	1.00	3.08 ± 0.62	3.15 ± 0.66	3.00 ± 0.61	a & b: <0.001* c & d: 0.78** a & c: <0.001** a & d: <0.001**

Data are presented as Mean ± SD. EPO: early onset preeclampsia; LOP: late onset preeclampsia
*P-value analyzed by students *t* test; **P-value analyzed by ANOVA with post-hoc Tukey’s test

Table III. Correlation between *TNF-α* and *Fas*, *FasL* expression

Gene		Pearson’s rho (<i>r</i>)	P-value
<i>TNF-α</i>	<i>Fas</i>	0.95	<0.001*
<i>TNF-α</i>	<i>FasL</i>	0.88	<0.001**

*P-value analyzed by Pearson’s correlation test between *TNF-α* and *Fas*; **P-value analyzed by Pearson’s correlation test between *TNF-α* and *FasL*




Figure 2. Comparison of PMBCs mRNA expression (fold change) between normotensive group and preeclampsia. Mean ± SD; * $p < 0.001$ analyzed by students’ *t* test




Figure 3. Comparison of PMBCs mRNA expression (fold change) between normotensive, early onset preeclampsia (EOP) and late onset preeclampsia (LOP). Mean \pm SD; * $p < 0.001$ EOP vs normotensive; ** $p < 0.001$ LOP vs normotensive, analyzed by ANOVA with post-hoc Tukey's test

4. Discussion

This study has reported elevated mRNA expression of *TNF- α* and its family members *Fas* and *FasL* in PBMCs in pregnancies complicated with PE as compared to normal pregnancies. The Fas-FasL, mediators of the intrinsic pathway of apoptosis, are deranged and show an incline in the expression in disorders characterized by disrupted immune tolerance or enhanced inflammation. Defective placentation with the disturbance in apoptosis in the placental bed has been well-studied. Researchers have reported differential Fas and Fas ligand expression in trophoblast in PE (5, 18). In 2011 Tomas and colleagues showed a disturbance of these with a decline in expression of FasL in the villous trophoblast in pregnancies complicated with PE (5). In contrast, Nishizawa and co-workers found an increase in FasL expression but no difference in Fas expression in placental tissue in PE and normal pregnancies (18).

In addition to trophoblast, Fas and FasL pathway is expressed and used by maternal lymphocytes. Altered expression of Fas has been reported in the lymphocytes in PE with a decrease in peripheral Gamma Delta T cells ($\gamma\delta$) and decidual Natural killer (NK) cells (19). This decrease in the expression

of Fas suggests decreased susceptibility of these cells to apoptosis, increasing the possibility of their contribution to the pathogenesis of preeclampsia (19). Moreover, utero placental ischemia results in excessive shedding of syncytiotrophoblast micro particles into the maternal circulation in PE. These syncytiotrophoblast micro particles (STBM) accentuate the systemic inflammation by enhancing the expression of proinflammatory cytokines by PBMCs, enlightening their role in the widespread inflammation (12). We hypothesized that the Fas and FasL pathway is altered in the PBMCs in PE and reported an increase in mRNA expression of both *Fas* and *FasL*. In contrast, Kuntz and colleagues have reported a decrease in surface expression of these two genes on leukocytes and lymphocytes in PE (20). The variation in the results could be attributed to the difference in the time of sample collection. In our study sample collection was done in the third trimester while they took the samples after delivery. The levels of different inflammatory markers change from late pregnancy to the postpartum period (21).

Defective placentation in PE results in oxidative stress and increased production of proinflammatory *TNF- α* with resultant activation

of leukocytes (22). In addition to increase in Fas and FasL in PE, we found a 3.08-fold increase in mRNA expression of *TNF- α* which is in accordance with the results reported by Chen and colleagues (17). Role of the excessive maternal systemic inflammatory response to pregnancy in the pathogenesis of PE is well established. There is a shift of cytokine profile in PE with an increase in proinflammatory cytokines as compared to the predominance of regulatory cytokine in normal pregnancy (23). A higher level of *TNF- α* , the proinflammatory cytokine, has been reported in the serum of pregnancies complicated with PE (24, 25). Extra placental sources like activated leukocytes or the endothelium might contribute to the increased levels of this inflammatory marker as there was no significant difference in the placental tissue (25). The higher mRNA expression of *TNF- α* in the PBMCs in PE in this study supports the involvement of tissues other than placenta in the pathogenesis of the disease. *TNF- α* enhances the production of eosinophil oxidants and toxicity toward human endothelial cells, activating the respiratory burst, and making the endothelium more vulnerable to oxidant damage (26). This study has also showed a significant positive correlation of *TNF- α* with Fas and FasL ($p \leq 0.001$). Our results are supported by Fluhr and co-workers who reported that *TNF- α* up regulates the expression of *Fas* in human endometrial stromal cells, sensitizing them to apoptosis (27). Moreover, *TNF- α* is known to promote Fas expression in trophoblast cells as evident by an in vitro study by Aschkenazi and colleagues (28). They worked on trophoblast cell cultures and found increased sensitivity of these cells to Fas-mediated apoptosis after treatment with *TNF- α* . Serum from preeclamptic pregnancies decreased the viability of trophoblast cells and increased their sensitivity to Fas-mediated apoptosis in first-trimester trophoblast cell line (29). Our findings of increased expression of FasL in PBMCs, enlighten the probability that these cells might contribute to enhanced apoptosis of trophoblast expressing Fas. Excessive apoptosis of cytotrophoblast in placenta

is reported by a number of researchers (30-32). Similarly, an increase in *Fas* expression by PBMCs in PE might result in apoptotic death of these cells by the *FasL*-expressing cytotrophoblast, affecting the survival of the fetal allograft escaping immune recognition.

5. Conclusion

The study concludes that mRNA expression of *TNF- α* , *Fas*, and *FasL* is altered in the maternal PBMCs in preeclamptic pregnancies and might contribute to the pathogenesis of the disease. Increased expression of *TNF- α* , *Fas*, *FasL* mRNA in PBMCs highlights the role of enhanced apoptosis and inflammation in PE. The regulation of *TNF- α* and *Fas*, *FasL* interactions in PBMCs need to be investigated in future approximations to better understand their role in the pathogenesis of PE. Longitudinal studies with multiple serial samples in different trimesters of pregnancy could enlighten the role of *Fas*, *FasL* pathway in the pathogenesis of PE. These processes can be potential targets for therapeutic agents to modify the course and improve the prognosis of the disease.

Acknowledgments

This research was partially funded from the research development fund by the University of Health Sciences Lahore, Lahore, Pakistan.

Conflict of Interest

The authors declare that there is no conflict of interest.

References

- [1] Brosens I, Pijnenborg R, Vercruyse L, Romero R. The "Great Obstetrical Syndromes" are associated with disorders of deep placentation. *Am J Obstet Gynecol* 2011; 204: 193–201.
- [2] Romero R, Kusanovic JP, Chaiworapongsa T, Hassan SS. Placental bed disorders in preterm labor, preterm PROM, spontaneous abortion and abruptio placentae. *Best Pract Res Clin Obstet Gynaecol* 2011; 25: 313–327.

- [3] Uzan J, Carbonnel M, Piconne O, Asmar R, Ayoubi JM. Pre-eclampsia: pathophysiology, diagnosis, and management. *Vasc Health Risk Manag* 2011; 7: 467–474.
- [4] Can M, Guven B, Bektas S, Arikan I. Oxidative stress and apoptosis in preeclampsia. *Tissue Cell* 2014; 46: 477–481.
- [5] Tomas SZ, Prusac IK, Roje D, Tadin I. Trophoblast apoptosis in placentas from pregnancies complicated by preeclampsia. *Gynecol Obstet Invest* 2011; 71: 250–255.
- [6] Barakonyi A, Miko E, Szereday L, Polgar PD, Nemeth T, Szekeres-Bartho J, et al. Cell death mechanisms and potentially cytotoxic natural immune cells in human pregnancies complicated by preeclampsia. *Reprod Sci* 2014; 21: 155–166.
- [7] Boyle JJ, Weissberg PL, Bennett MR. Tumor necrosis factor- α promotes macrophage-induced vascular smooth muscle cell apoptosis by direct and autocrine mechanisms. *Arterioscler Thromb Vasc Biol* 2003; 23: 1553–1558.
- [8] Cackovic M, Buhimschi CS, Zhao G, Funai EF, Norwitz ER, Kuczynski E, et al. Fractional excretion of tumor necrosis factor- α in women with severe preeclampsia. *Obstet Gynecol* 2008; 112: 93–100.
- [9] Zhou P, Luo X, Qi HB, Zong WJ, Zhang H, Liu DD, et al. The expression of pentraxin 3 and tumor necrosis factor- α is increased in preeclamptic placental tissue and maternal serum. *Inflamm Res* 2012; 61: 1005–1012.
- [10] Smith SD, Dunk CE, Aplin JD, Harris LK, Jones RL. Evidence for immune cell involvement in decidual spiral arteriole remodeling in early human pregnancy. *Am J Pathol* 2009; 174: 1959–1971.
- [11] Mihu D, Razvan C, Malutan A, Mihaela C. Evaluation of maternal systemic inflammatory response in preeclampsia. *Taiwan J Obstet Gynecol* 2015; 54: 160–166.
- [12] Germain SJ, Sacks GP, Soorana SR, Sargent IL, Redman CW. Systemic inflammatory priming in normal pregnancy and preeclampsia: the role of circulating syncytiotrophoblast microparticles. *J Immunol* 2007; 178: 5949–5956.
- [13] Kauma SW, Huff TF, Hayes N, Nilkaeo A. Placental Fas ligand expression is a mechanism for maternal immune tolerance to the fetus. *J Clin Endocrinol Metab* 1999; 84: 2188–2194.
- [14] Resic Karara J, Zekic Tomas S, Marusic J, Roje D, Kuzmic Prusac I. Fas and FasL expression in placentas complicated with intrauterine growth retardation with and without preeclampsia. *J Matern Fetal Neonatal Med* 2016; 29: 1154–1159.
- [15] Prusac IK, Zekic Tomas S, Roje D. Apoptosis, proliferation and Fas ligand expression in placental trophoblast from pregnancies complicated by HELLP syndrome or preeclampsia. *Acta Obstet Gynecol Scand* 2011; 90: 1157–1163.
- [16] Martinez-Fierro ML, Garza-Veloz I, Carrillo-Sanchez K, Martinez-Gaytan V, Cortes-Flores R, Ochoa-Torres MA, et al. Expression levels of seven candidate genes in human peripheral blood mononuclear cells and their association with preeclampsia. *Hypertens Pregnancy* 2014; 33: 191–203.
- [17] Chen G, Wilson R, Wang SH, Zheng HZ, Walker JJ, McKillop JH. Tumour necrosis factor- α (TNF- α) gene polymorphism and expression in pre-eclampsia. *Clin Exp Immunol* 1996; 104: 154–159.
- [18] Nishizawa H, Ota S, Suzuki M, Kato T, Sekiya T, Kurahashi H, et al. Comparative gene expression profiling of placentas from patients with severe pre-eclampsia and unexplained fetal growth restriction. *Reprod Biol Endocrinol* 2011; 9: 107–118.
- [19] Barakonyi A, Miko E, Szereday L, Polgar PD, Nemeth T, Szekeres-Bartho J, et al. Cell death mechanisms and potentially cytotoxic natural immune cells in human pregnancies complicated by preeclampsia. *Reprod Sci* 2014; 21: 155–166.
- [20] Kuntz TB, Christensen RD, Stegner J, Duff P, Koenig JM. Fas and Fas ligand expression in maternal blood and in umbilical cord blood in preeclampsia. *Pediatr Res* 2001; 50: 743–749.
- [21] Bränn E, Edvinsson Å, Rostedt Punga A, Sundström-Poromaa I, Skalkidou A. Inflammatory and anti-inflammatory markers in plasma: from late pregnancy to early postpartum. *Sci Rep* 2019; 9: 1863–1872.
- [22] Raghupathy R. Cytokines as key players in the pathophysiology of preeclampsia. *Med Princ Pract* 2013; 22 (Suppl.): 8–19.
- [23] Pinheiro MB, Martins-Filho OA, Mota AP, Alpoim PN, Godoi LC, Silveira AC, et al. Severe preeclampsia goes along with a cytokine network disturbance towards a systemic inflammatory state. *Cytokine* 2013; 62: 165–173.
- [24] Perucci LO, Gomes KB, Freitas LG, Godoi LC, Alpoim PN, Pinheiro MB, et al. Soluble endoglin, transforming growth factor-Beta 1 and soluble tumor necrosis factor alpha receptors in different clinical manifestations of preeclampsia. *PLoS One* 2014; 9: e97632.
- [25] Benyo DF, Smarason A, Redman CW, Sims C, Conrad KP. Expression of inflammatory cytokines in placentas from women with preeclampsia. *J Clin Endocrinol Metab* 2001; 86: 2505–2512.
- [26] Slungaard A, Vercellotti GM, Walker G, Nelson RD, Jacob HS. Tumor necrosis factor alpha/cachectin stimulates eosinophil oxidant production and toxicity towards human endothelium. *J Exp Med* 1990; 171: 2025–2041.
- [27] Fluhr H, Krenzer S, Stein GM, Stork B, Deperschmidt M, Wallwiener D, et al. Interferon- γ and tumor necrosis factor- α sensitize primarily resistant human endometrial stromal cells to Fas-mediated apoptosis. *J Cell Sci* 2007; 120: 4126–4133.
- [28] Aschkenazi S, Straszewski S, Verwer KM, Foellmer H, Rutherford T, Mor G. Differential regulation and function of the Fas/Fas ligand system in human trophoblast cells. *Biol Reprod* 2002; 66: 1853–1861.
- [29] Neale D, Demasio K, Illuzi J, Chaiworapongsa T, Romero R, Mor G. Maternal serum of women with pre-eclampsia reduces trophoblast cell viability: evidence for an increased sensitivity to Fas-mediated apoptosis. *J Matern Fetal Neonatal Med* 2003; 13: 39–44.
- [30] Dagdelen M, Temur M, Yılmaz Ö, Altındag T, Uslu T, Özbay PÖ. Placental bed apoptosis is increased in pregnant women with pre-eclampsia versus normotensive pregnant women. *J Obstet Gynaecol* 2016; 36: 974–979.
- [31] Mendilcioglu I, Karaveli S, Erdogan G, Simsek M, Taskin O, Ozekinci M. Apoptosis and expression of Bcl-2, Bax, p53, caspase-3, and Fas, Fas ligand in placentas complicated by preeclampsia. *Clin Exp Obstet Gynecol* 2011; 38: 38–42.
- [32] Longtine MS, Chen B, Odibo AO, Zhong Y, Nelson DM. Villous trophoblast apoptosis is elevated and restricted to cytotrophoblasts in pregnancies complicated by preeclampsia, IUGR, or preeclampsia with IUGR. *Placenta* 2012; 33: 352–359.