

Short communication

Perifollicular and subendometrial blood flow and ICSI cycle outcome

Elham Pourmatroud^{1, 2*} M.D., Batoool Hosein Rashidi^{1*} M.D., Maryam Rahmani³ M.D.

- 1 Vali-e-asr Reproductive Health Research Center, Tehran University of Medical Sciences, Tehran, Iran.
- 2 Department of Obstetrics and Gynecology (IVF), Ahvaz Joundishapoor University of Medical Sciences (AJUMS), Ahvaz, Iran.
- 3 Emam Khomini Hospital, Tehran University of Medical Sciences, Tahran, Iran.

Received: 25 November 2009; accepted: 28 April 2010

Abstract

Background: There is a fundamental correlation between follicles and endometrium in intracytoplasmic sperm injection (ICSI) cycles.

Objective: To assess the relation between perifollicular perfusion and sub endometrial parameters in Doppler ultrasonography and outcome of in ICSI cycles.

Materials and Methods: In this prospective descriptive pilot study, 10 patients were enrolled. Strict inclusion criteria were considered. Routine long protocol was used for ICSI. On the day of follicle retrieval, colour Doppler indices were determined. Sub endometrial pulsatility index (PI) and resistance index (RI) and perifollicular perfusion were assessed. After oocyte retrieval the count of metaphase 2 (M2) oocytes, embryo with grade A quality and the result of cycle were evaluated also.

Results: RI and PI indices had a positive correlation. Follicles with ≥ 18 mm diameter and follicles with $>75\%$ perfusion had a direct relation. Also, subendometrial RI had a significant relation with follicular status (p-value= 0.04), But there was not a significant triple correlation (between endometrium, follicles and outcome).

Conclusion: The mutual effects of vascularization status in two fundamental parts in ART, is still unclear. The evaluation with Doppler ultrasonography should focus on two compartments together as one functional part at the same time. It means even in presence of good markers in each part, the final decision must be taken by co-evaluation of follicles and endometrium.

Key words: Doppler sonography, Endometrium, Follicle, ICSI.

Introduction

Perifollicular perfusion and ovarian stromal blood flow are useful markers for optimal evaluation of follicles, with the former having a direct relationship with follicular oxygenation and oocyte maturation (1). The chance of obtaining a high-quality oocyte, and thus a good embryo, increases when perifollicular blood flow

which is preferably between 50% and 75% (2) and higher pregnancy rate will be expected when at least one embryo is from a highly-perfused follicle (3). In the past decade, several studies showed that endometrial thickness does not indicate better receptivity, nor does it have a correlation with the outcome of ART cycles (4, 5). The multi-layered endometrium is not always with pregnancy outcome, after embryo transfer. Because a good endometrial blood flow is necessary for implantation to take place in ART, Doppler sonography could be a helpful tool in the assessment of endometrial and sub-endometrial blood flow. The aim of our study was to evaluate two topics at the same time; perifollicular

Corresponding Author:

Elham Pourmatroud; Azadegan Street, Emam Khomini Hospital, Ahvaz, Iran.

Email: e.pourmatroud@yahoo.com

*These authors have contributed equally to this work.

perfusion as well as PI (pulsatility index), and RI (Resistance index) of subendometrial blood flow. We focused on those parameters because the association between perfollicular perfusion and follicular oxygenation and oocyte maturation has been approved. Furthermore, PI and RI are the most conventional and comprehensive characters for subendometrial blood flow description.

Materials and methods

Subject selection

This prospective-analytical pilot study was conducted in the infertility clinic (Vali-e-Asr Reproductive Health Research Centre) of a tertiary teaching hospital, affiliated to Tehran University of Medical Sciences, Tehran, Iran. Approval was obtained from institutional Research Ethics Board. Also, oral consent was taken from all participants. All subjects were 20-35 years old women who were in the first ART cycle. All participants were non-smokers, did not use any drug and had normal basal follicle stimulating hormone level (FSH<10 mIU/ml) and regular menstruation cycles.

Exclusion criteria were:

- 1) History of or present disease of uterus such as endometrial polyp, uterine myoma or congenital uterine anomaly.
- 2) History of or present disease of ovaries (like ovarian cyst) or fallopian tubes (especially hydrosalpinx) (7).
- 3) History of or present endometriosis which has been confirmed by laparoscopy.

Within a period of six months (July 2008-December 2008), ten patients met all the above-mentioned criteria and were enrolled into the study.

ICSI protocol

Routine OCP+long GnRH agonist protocol (Superfact, Avantis, Germany+Gonal F, Merk serono, Germany) was used for ovarian stimulation. Patients with only one 18mm follicle or excessive ovarian response with tendency of developing ovarian hyperstimulation syndrome (OHSS) were excluded from the study. Oocyte retrieval was done with vaginal sonographic guidance 36-40 hours after HCG (Pregnyl, Organon, Spain) administration. From each ovary, oocytes in metaphase II maturation phase were determined separately. Intra-cytoplasmic sperm injection was performed for fertilization of oocytes.

On the third day, eight-cell embryos of equal size and regular blastomeres (grade A) (8) calculated in each patient and a maximum of two grade A embryos were transferred vaginally and if there, the others were frozen. Luteal phase support was started from oocyte pickup day by micronized vaginal progesterone (400mg BID) and continued for 14 days. Pregnancy was then checked by the serum β hCG level and luteal support was continued in case of positive result. Six weeks after embryo transfer, vaginal ultrasonography was done in pregnant women to determinate the status of pregnancy.

Intervention

Ultrasonographic evaluation was done on the day of oocyte retrieval. In order to reduce diurnal variation of blood flow, patients evaluated on the morning (7.5-8.5AM) before going to the operation room (9). All 2D ultrasonographic measurements were performed by a single observer, using a 7.5 MHz transvaginal transducer (EV 9-4) with colour Doppler facility (Siemens Sonoline G605). At first, conventional vaginal 2D-ultrasonography was done and the number of follicles \geq 18mm in each ovary, maximum endometrial thickness (mm), and its morphologic appearance (triple-layer or non-triple layer) were determined.

After that, the power Doppler window was placed on the highest longitudinal plane of both ovaries (with normal quality of colour, colour gain - 3.4 pulse repetition frequencies of 600HZ and wall motion filter of 50HZ). When an adequate power Doppler signal was obtained, perfollicular perfusion was evaluated according to follicular circumference. Follicles with \geq 75% perfusion were determined in each ovary. If "flash" artefacts were seen (due to patient's breathing or intestinal movement), the evaluation was repeated until a constant result was achieved. The probe was then placed over the longitudinal section of the uterus, which presents the whole subendometrial area. The uterine subendometrial region was evaluated using a scanning angle of 90 degrees and colour setting. The form of the velocity wave was analyzed in at least three points during three to five cardiac cycles and mean RI and PI values were calculated.

Statistical analysis

Data were entered into SPSS version 11.5 computer software and subsequently analyzed using statistical tests. Student's t-test was used for quantitative parameters. Correlation was

assessed by Pearson method. P-value <0.05 was considered as significant (power study was 80%).

Results

The mean age of patients was 28.2±4.7 years (range=21-35years). Eight patients had primary and two had secondary infertility. Causes of infertility were: unexplained (3 cases), male factor (4 cases), ovulatory dysfunction (2 cases), and tubal factor (2 cases). One patient had two infertility factors. Other data of the subjects have been listed in table I.

Table I: Demographic data in study group.

Parameters	Mean±SD	Minimum Maximum
Duration of infertility (year)	6.4±4.1	2-14
Dose of gonadotropin (Amp)	25.1±12.6	14-55
Duration of stimulation (day)	9.6±1.1	7-11
Endometrial thickness (mm)	8.5±1.7	5.5-12
Sub endometrial PI	2.08±0.87	1.09-3.53
Sub endometrial RI	0.76±0.08	0.67-0.93
Follicles ≥18 mm RT	4.8±2.5	2-10
Follicles ≥18 mm LT	4.2±1.9	1-8
Follicles >75% perfusion RT	2.2±1.7	0-6
Follicles >75% perfusion LT	2±1.4	0-4
Picked up oocyte RT	5±2.9	1-11
Picked up oocyte LT	3.6±1.4	1-6
Metaphase 2 oocyte RT	3.5±2.5	1-1
Metaphase 2 oocyte LT	1.9±0.7	1-3
Total embryo numbers	4.6±2.7	2-11
Grade A embryo	1.9±5.2	0-4

Resistance index (RI) and pulsatility index (PI) had direct correlation (p-value=0.007). Neither endometrial thickness nor its appearance had any relationship with subendometrial blood flow, RI, or PI. Also, the dose of gonadotropins and duration of ovarian stimulation had no effect on these parameters. There was a significant correlation between bilateral ovarian follicles with ≥18 mm in diameter and the number of follicles with ≥75% perfollicular perfusion (p-value=0.004), but none had a relationship with the number of metaphase II oocytes or grade A embryos. Also, the total number of oocytes picked-up from the right ovary correlated with the number of metaphase II oocytes (p-value=0.006) and the total number of embryos (p-value=0.005). Surprisingly, the total dose of gonadotropins has a positive correlation with grade A embryos (p-value=0.01). Regarding the correlation between subendometrial blood flow

or perfollicular perfusion with other parameters, a significant negative relationship was found to exist between subendometrial RI and ≥75% perfused follicles in the right ovary (p-value=0.040) and a positive relationship was found to exist with the total oocytes retrieved from the left ovary (p-value=0.01). In this study, two patients became pregnant but none of the parameters were found to have significant relationship with this outcome.

Discussion

In this study, due to the presence of several confounding factors and strict inclusion criteria, we finally managed to enroll only ten patients. Unfortunately, the correlation between clinical outcome and colour Doppler ultrasonographic parameters was not efficacious with our low study population.

In order to increase the chance of pregnancy in ART cycles, better synchronization between ovarian follicles and the endometrium is required; with a suitable oocyte (metaphase II), endometrial receptivity must be around a specific time (implantation window). Therefore, it would be more helpful if utero-ovarian vascularity was assessed at the same time, especially in cases with monofollicular cycles (10) or in patients which need natural cycle, such as those with specific medical conditions.

On the other hand, although there is evidence suggesting a link between uterine and ovarian perfusion and outcome of ART cycles (11), however it is difficult, at present, to interpret the actual relationship. In our study, we found a new correlation between resistance index (RI) of subendometrial blood flow and ovarian response but we couldn't explain why the correlation was negative in one side. Probably if other parameters like flow index (FI) or vascularization index (VI) were used, our results would have been different. Also, another explanation would have been needed if the evaluation had been done on the day of HCG administration (12) or if it was repeated within two or three days (13).

In conclusion, we believe that the utilization of Doppler ultrasonography usage in ART cycle is in its primary stages. This report re-emphasizes that: the evaluation with Doppler ultrasonography should focus on two compartments together as one functional part at the same. It means even in presence of good markers in each part, the final decision must be taken by co-evaluation of follicles and endometrium.

References

1. Van Blerkom J. Can the developmental competence of early human embryos be predicted effectively in the clinical IVF laboratory. *Hum Reprod* 1997; 12:1610-1614.
2. Coulam CB, Good C, Rinehart JS. Color Doppler indices of follicular blood flow as predictor of pregnancy after IVF and embryo transfer. *Hum Reprod* 1999; 14: 1979-1982.
3. Robson SJ, Norman RJ. Power Doppler assessment of follicle vascularity at the time of oocyte retrieval in invitro fertilization cycle. *Fertil Steril* 2008; 90: 2179-2182.
4. Li TC, Nuttal L, Klentzeris L, Cook ID. How well does ultrasonographic measurements of endometrial thickness predict the result of histological dating? *Hum Reprod* 1992; 7: 1-5
5. Dietterich C, Check JH, Choe JK, Nazari A, Lurie D. Increased endometrial thickness on the day of HCG injection does not adversely affect pregnancy or implantation rates following IVF/ET. *Fertil Steril* 2002; 77: 781-786.
6. Ng EH, Chan CC, Tang OS, Ho PC. Comparison of endometrial and sub endometrial blood flows among patients with and without hydrosalpinx shown an scanning during IVF treatment. *Fertil Steril* 2006; 85: 333-338.
7. Hrishikesh D Pai. Embryo transfer technique. In: Kamini A Rao. Laboratory manual in assisted reproductive technology. Newdelhi India JAYPEE Brothers medical Publishers Ltd 2006; 202.
8. Ziadi J, Jurkovic D, Campbell S, Pittrof R, MC Gregor A, Tan SL. Description of circadian rhythm in uterine artery blood flow during the peri ovulatory period. *Hum Reprod* 1995; 10: 1642-1646.
9. Vlaisavljevic V, Reljic M, Gavric Iovrec V, Zazula D, Sergeant N. Measurement of perifollicular blood flow of the dominant preovulatory follicle using 3D power Doppler. *Ultrasound Obstet Gyn* 2003; 22: 520-526.
10. Ozturk O, Bhattacharya S, Saridogan E, Jauiaux E, Templeton A. Role of utero ovarian vascular impedance: predictor of ongoing pregnancy in an IVF-ET programme. *Repro biomed online* 2004; 9: 299-305.
11. Dechand H, Bessueille E, Bousquet PJ, Regftman N, Hamamah S, Hedon B. Optimal timing of ultrasonographic and Doppler evaluation of uterine receptivity to implantation. *Repro Biomed Online* 2008; 16: 368-375.
12. Chien LW, Lee WS, AU HK, Tzeng CR. Assessment of changes in utero ovarian arterial impedance during the peri implantation period by Doppler sonography in women undergoing assisted reproduction. *Ultrasound Obstet Gyn* 2004; 23: 496-500.
13. Ng EH, Chan CC, Tang OS, Yeung WS, Ho PC. The role of endometrial blood flow measured by 3D power Doppler ultrasound in the prediction of pregnancy during IVF fertilization treatment. *Eur J Obstet Gyn Reprod Boil* 2007; 135: 8-16.