

## Antispasmodic effects of *Citrus aurantium* flowers aqueous extract on uterus of non-pregnant rats

Akram Ahangarpour<sup>1</sup> Ph.D., Ali Akbar Oroojan<sup>2</sup> B.Sc., Ashraf Amirzargar<sup>2</sup> M.Sc., Maryam Ghanavati<sup>3</sup> B.Sc.

1 Department of Physiology, School of Medicine, Physiology and Diabetes Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

2 Department of Physiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

3 Department of Nursing, School of Nursing and Midwifery, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

Received: 21 July 2010; accepted: 5 January 2011

### Abstract

**Background:** *Citrus aurantium* is a small citrus tree, with scented white flowers. The *C. aurantium* is used in Asian herbal medicine primarily to treat digestive problems.

**Objective:** The goal of this study is to investigate the effect of *C. aurantium* flower's aqueous extract on uterine contraction in presence of some known uterus stimulants.

**Materials and Methods:** In experimental study 30 virgin Wistar rats 200-300gr were obtained. After laparotomy, a piece of Uterus was dissected out and mounted in an organ bath (10ml) containing De Jalon (29°C) and contracted by KCl (60mM), oxytocin (10mU/ml) and barium chloride (4mM) then the effect of *C. aurantium* flower's aqueous extract (1-8 mg/ml) on the uterine contractions was investigated. Uterus was separately incubated with propranolol (1μM), naloxone (1μM) and the role of β-adrenoceptors, opioid receptors were evaluated.

**Results:** Cumulative concentrations of the extract (1-8 mg/ml) decreased KCl, oxytocin and barium chloride induced uterine contractions, dose-dependently (p<0.001). *C. aurantium* flower's aqueous extract was unaffected on incubation the tissue with propranolol and naloxone.

**Conclusion:** It seems that the extract induced antispasmodic effect mainly via calcium influx blockade. However, neither β-adrenoceptors nor opioid receptors were involved. Since the extract has antispasmodic effect on uterus contraction therefore we can suggest that more study will be necessary to relief dysmenorrheal.

**Key words:** *Citrus aurantium*, Propranolol, Naloxone, Antispasmodic, Uterus, Rat.

### Introduction

*Citrus aurantium*, Seville or sour orange is a small citrus tree, with aroma white flowers. The most common use of *C. aurantium* is medicinal rather than culinary. The entire Raw fruit, is used in Asian herbal medicine primarily to Remedy digestive problems. There is thus a history of benign human consumption of *C. aurantium* fruit (1). This herb use for treatment of obesity and

suppress appetite (2). *C. aurantium* has pharmacological applications that its flowers have been used as a sedation and heart tonic in traditional medicine (3). In humans, rats, guinea pigs, cats, and/or dogs, extracts of *C. aurantium* have been evaluated for effects on the cardiovascular system that include changes in blood pressure, cardiovascular toxicity, contractility and excitability of the heart muscle, and/or adrenergic activity (4). *C. aurantium* extract can increase the L-type calcium current in guinea pig ventricular myocytes significantly in a concentration-dependent manner, and promote the opening of calcium channel (5). Studies of the flowers of this natural medicine have reported that

### Corresponding Author:

Akram Ahangarpour, Department of Physiology, Physiology and Diabetes Research Center, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran.

**Email:** ahang1002002@yahoo.com

some constituents exhibited antioxidant activity by flavonoids containing and the aqueous extract showed inotropic effect. The *C. aurantium* extract had an effect in the hypertension (1, 2). The new products that called “ephedra free” usually contain *C. aurantium* extracts and it has 3 to 6% of p-synephrine. p-synephrine acts such as cardiac debit increase, peripheric vasoconstriction, broncodilatation and CNS stimulation (6). Since in traditional medicine the use of this plant was recommended in pregnancy therefore the present study was an attempt to investigate the effect of *C. aurantium* flower aqueous extract on uterine contraction and its possible mechanism.

## Materials and methods

### Plant extraction

*C. aurantium* flower were purchased from local herbal shop in 2009 and were powdered by miller machine. Aqueous extract was accommodated by sokcele method using 30gr powder and 300ml distilled water for 8 hours. The complex was filtered with Whatman No1 filter paper. After that solution dried at 25°C to obtain powder and it was stored at Refrigerator until being used.

### Animals and tissue preparation

30 Adult female Wistar rats (200-300gr) were obtained from Ahwaz Jundishapur University of Medical Sciences animal house and kept at 12- h light/ dark cycle and at 20-24°C were allowed free access to tap water and commercial chow. They were divided in to 5 groups (KCl-induced, barium chloride- induced, propranolol incubation, naloxane incubation and oxytocin-induced).

In this study the rats were sacrificed by a sharp blow on the neck. After laparotomy uterus was removed and washed with cold and oxygenated with De Jalon solution and cut into 1.5cm long and then mounted in an organ bath containing De Jalon solution (10ml) between two stainless steel hooks vertically. One of the hooks was upper hook that connected to an isometric transducer (UF1 Harvard transducer, UK) and this transducer was connected to a recorder machine (Harvard Universal Oscillograph, UK) and another one was lower hook that fixed at the bottom of the organ bath. The De Jalon solution was contained NaCl (154), KCl (5.6), CaCl<sub>2</sub> (0.3), NaHCO<sub>3</sub> (1.7), MgCl<sub>2</sub> (1.4) and glucose (5.55) (in mM) with pH of 7.4 and temperature of 29°C and the bubble of air was present in the De Jalon continuously (6). The piece of uterus was then Keep under 1g initial tension and Permit to Parity for 1h during which bath solution was washed every 15 min.

### Drugs

Potassium chloride and barium chloride were purchased from Merck (Germany), naloxone (Tolid Daru Company, Tehran-Iran), propranolol (Tolid Daru Company, Tehran-Iran) and oxytocin (Weimer Pharma, Germany).

*C. aurantium* extract powder and all chemicals were dissolved in De Jalon and added to organ bath, degree of drugs solution was equivalent with laboratories temperature (29°C).

### Experimental protocols

After equilibrium period, the uterus was contracted by KCL 60mM (7, 8) and barium chloride 4mM (9) in the plateau of contraction the concentrations of extract (1, 2, 4 and 8 mg/ml) were added cumulatively. For the study uterus opioid receptors, naloxone (1µM) and for the study of β-adrenoceptors receptors, propranolol (1µM) (10) were added to organ bath about 30 min without any refreshing for blocking these receptors in uterus tissue and after that concentrations of extract (1, 2, 4 and 8 mg/ml) were added cumulatively. Uterus tissue contracted by oxytocin (10 mU/ml) (10) and when oxytocin doesn't have plateau the tissue was washed. After that extract was added to organ bath without any dose of oxytocin and then percent's differences between peak amplitude of oxytocin was calculated. These protocols repeated for 6 rats in each groups.

### Statistical analysis

The mean±SEM of contraction forces induced by KCl, barium chloride and OT were regarded as (0%) and calculated for each group. The results were statistically analyzed by one way ANOVA and post hoc LSD tests and (p-values <0.05) was significant. The (n) was the number of animals that used in each group.

## Results

### Effect of cumulatively concentration of *C. aurantium* flowers aqueous extract after making uterus contractions with KCL (60mM)

KCl (60mM) induced made contraction in uterus tissue, after 2min when the contractions of uterus arrived to the plateau, the concentrations of *C. aurantium* flowers aqueous extract (1, 2, 4 and 8 mg/ml) was added cumulatively to the organ bath with 10 min intervals and without refreshing bath solution. KCl (60mM) had significantly difference with all of the concentrations of extracts (p<0.001) also the relaxation effect of 1mg/ ml extract was significantly different with 4 and 8 mg/ml (p<0.01) of extract on contraction that made by KCL (Figure 1).

**Effect of *C. aurantium* flowers aqueous extract activity after tissue incubation with propranolol**

After induced uterus contraction by KCl (60mM) and without refreshing bath, propranolol (1µM) added to organ bath for 30 min without any refreshing for blocking β-adrenoceptors, then concentrations of extract (1, 2, 4 and 8 mg/ml) were added cumulatively to the organ bath with 10 min intervals and without refreshing.

There were significantly differences among propranolol (1µM) with 2 (p<0.01), 4 and 8 mg/ml (p<0.001) of extract. Relaxation effect of 1mg/ml extract had significant differences with 4 and 8 mg/ml (p<0.001) also this effect of 2mg/ml extract had significant differences with 4 (p<0.05) and 8 mg/ml (p<0.01) (Figure 1).

**Effect of *C. aurantium* flowers aqueous extract on barium chloride-induced contraction**

After barium chloride (4mM) induced uterus contraction, *C. aurantium* flowers aqueous extract (1, 2, 4 and 8 mg/ml) was added cumulatively. Contractions barium chloride (4mM) had significant difference with another groups (p<0.001). Relaxation effect of 1 and 2 mg/ml extract has significant difference with 8 mg/ml of extract (p<0.001). Also 1mg/ml of *C. aurantium* flowers aqueous extract had significant difference (p<0.05) with 4 mg/ml of extract (Figure 2).


**Effect of *C. aurantium* flowers aqueous extract activity after tissue incubation with naloxone**

KCl (60mM) induced made contraction in uterus tissue after 2min interval, naloxone (1µM) as a non- selective opioid receptor antagonist added to organ bath for 30 min without refreshing. Then 1, 2, 4 and 8 mg/ml of extract were added to organ bath cumulatively and the interval was 10min. The result showed a significant difference between naloxone (1µM) and all groups (p<0.001) and relaxation effect of 1 mg/ml extract had significant differences with 8 mg/ml (p<0.05) group (Figure 2).


**Effect of *C. aurantium* flowers aqueous extract activity on oxytocin-induced uterus contraction**

First oxytocin (10mU/ml) was added to the bath with 3min interval, then uterus tissue in the bath was washed with De Jalon. Then 1mg/ml concentration of *C. aurantium* flowers aqueous extract was added to the bath for 3 min and without any refreshing, oxytocin was used with more than 3min interval. This protocol has been used for 2, 4 and 8 mg/ml concentration of extract. The time was about 15 min. After that all of the uterus tissue


was washed. All of the concentrations of extract had shown relaxation effect with oxytocin (10mU/ml) group significantly (p<0.001). Also 1mg/ml of extract had significant difference with 2 (p<0.05), 4 and 8 mg/ml (p<0.001) and the difference between 2 mg/ml with 8 mg/ml (p<0.05) was significant (Figure 3).



**Figure 1.** Effect of cumulative concentration of *C. aurantium* aqueous extract on KCL (60 mM) induced uterus contractions and after incubation with propranolol (1 µM) (n=6, \*\*= p<0.01, \*\*\*= p<0.001). a= difference between KCL group with other groups. b= difference between 1mg/ml group with other groups. c= difference between 2mg/ml group with other groups. d= difference between 4mg/ml group with other groups. e= difference between 8mg/ml group with other groups.



**Figure 2.** Effect of cumulative concentration of *C. aurantium* aqueous extract on barium chloride (4 mM) induced uterus contractions and after incubation with naloxone (1 µM) (n=6, \*= p<0.05, \*\*\*= p<0.001). a= difference between barium chloride group with other groups. b= difference between 1mg/ml group with other groups. c= difference between 2mg/ml group with other groups. d= difference between 4mg/ml group with other groups. e= difference between 8mg/ml group with other groups.



**Figure 3.** Effect of non-cumulative concentration of *C. aurantium* aqueous extract on rat uterus contracted by oxytocin (10mU/ml) (n=6, \*= p<0.05, \*\*\*= p<0.001).

a= difference between oxytocin group with other groups.  
 b= difference between 1mg/ml group with other groups.  
 c= difference between 2mg/ml group with other groups.  
 d= difference between 4mg/ml group with other groups.  
 e= difference between 8mg/ml group with other groups.

## Discussion

Present results of this study showed that aqueous extract of *C. aurantium* flowers induces antispasmodic effect on the uterine muscle contraction caused by KCl, barium chloride or oxytocin. Also highest dose of the aqueous extract had most anticontractile effect on the uterus contraction.

Propranolol is  $\beta$ -adrenoceptors antagonism and in some studies it seems clear that activation of  $\beta$ -adrenoceptors in uterine smooth muscle results in inhibition of myometrial contraction (11), after incubation with propranolol (1 $\mu$ M), antispasmodic effect of *C. aurantium* flowers aqueous extract decreased the contractile tension of uterine smooth muscle that made by KCl 60 mM and propranolol (1 $\mu$ M) incubated in nonpregnant virgin's rat. According to these results we can suggest that *C. aurantium* flowers aqueous extract doesn't have any effect from  $\beta$ -adrenoceptors. In endometrial and myometrial regions of the uterus have been observed Opioid system expression (12).

Opioid receptors activation relaxes uterus (13). Naloxone is a nonselective opioid receptors antagonist. In these results *C. aurantium* flowers aqueous extract activity had relaxant effect on uterus after incubation by naloxone (1 $\mu$ M). Therefore we can suggest that extract activity was

not mediated via these receptors. Uterine smooth muscle contraction is mediated mainly via increased intracellular  $Ca^{2+}$  and is accomplished by excitation-contraction coupling mechanisms. A high  $K^+$  medium could depolarize the cellular membrane of uterine smooth muscle (14).

Moreover, it is well-known that  $K^+$  induced contraction in smooth muscles is due to an increase in  $Ca^{2+}$  influx through voltage-operated  $Ca^{2+}$  channels. It has been suggested that any substance that inhibits high  $K^+$  induced contractions of smooth muscles is a blocker of  $Ca^{2+}$  influx (15). KCl induced uterine smooth muscle contraction results mainly from calcium influx via voltage-sensitive calcium channels (16).

The present findings suggest that aqueous extract of *C. aurantium* flowers inhibits KCl induced uterine contractions acting on voltage-sensitive calcium channels, because this contraction is dependent on extracellular calcium. Oxytocin (OT) induces its effects via OT receptors (OTRs), which are members of the heptahelical family of G protein-coupled receptors. It is now well established that OT binding to its myometrial receptors leads to an increase in intracellular free calcium via the generation of inositol trisphosphate (17).

The oxytocin stimulated increase in  $[Ca^{2+}]_i$  is rapid and declines rather quickly in comparison to that seen in some other cell types, presumably as a result of active pumping of calcium out of the cell and back into the endoplasmic reticulum by the calcium transport ATPases (18). In this study, the effect of aqueous extract of *C. aurantium* flowers on contraction of uterus that has been made by oxytocin-induced was investigated. This investigation had done in a De Jalon solution and this solution had calcium concentration in itself.

This extract reduce spontaneous motility and superimposing contractions with oxytocin-induced and this relaxation effect was dose-dependently that 8mg/ml of this extract had most relaxation effect in uterus muscle. Oxytocin also activates the L-type VDCCs (voltage dependent calcium channels) (17, 18).

In our results, we demonstrated that aqueous extract of *C. aurantium* flowers reduced the oxytocin-induced contractions in De Jalon solution with calcium. Therefore we can suggest that the effect of this extract is via blockade L-type VDCCs that redound to relax the contracted of uterus tissue.

Also we can say that voltage sensitive calcium channels have specific role in these contractions that decreased by using the concentrations of this extract. The effect of Barium chloride is inducing contraction in smooth muscle via blocking the potassium channels (19). Also, it has been reported that  $Ba^{2+}$  increases  $Ca^{2+}$  release from intracellular pools in smooth muscles and uterus (9). Our result showed that *C. aurantium* flowers extract reduced the contractions that induced by barium chloride dose- dependently, and this result was from prevention of releasing calcium or calcium efflux.

In conclusion our result indicated that *C. aurantium* flowers aqueous extract induces antispasmodic effect on isolated rat uterus and this extract decreased the uterus contractions without involving  $\beta$ - adrenoceptors, and opioid receptors. These results indicate that *C. aurantium* flowers aqueous extract induces antispasmodic effect on rat uterus mainly through blockage of the VDCCs.

The precise mechanism of extract activity can be the study of  $Ca^{2+}$  alterations in the presence of the extract in animals or human myometrium. The antispasmodic effect of the plant's extract observed in the present study, support the clinical efficacy and use of *C. aurantium* flowers in the treatment of dysmenorrhoea and other uterine spasmodic disorders.

This process appears to be the most relevant physiologically and should be concentrated on in the future research.

### Acknowledgment

This research was financially supported by the University Student Research Center of Ahvaz Jundishapour University of Medical Sciences, Ahvaz, Iran.

### References

1. Fugh- Berman A, Myers A. Citrus aurantium, an Ingredient of Dietary Supplements Marketed for Weight Loss: Current Status of Clinical and Basic Research. *Exp Biol Med (Maywood)* 2004; 229: 698-704.
2. Monsef-Esfahani HR, Amanzade Y, Alhani Z, Hajimehdipour H, Faramarzi MA. GC/ MS analysis of Citrus aurantium L. hydrolate and its comparison with the commercial samples. *Iran J Pharm Res* 2004; 3: 177-179.
3. Calapai G, Firenzuoli F, Saitta A, Squadrito F, Arlotta MR, Costantino G, et al. Antiobesity and cardiovascular toxic effects of Citrus aurantium extracts in the rat: A preliminary report. *Fitoterapia* 1999; 70: 586-592.
4. Fang F, Dong M, Zhu H. Effect of Citrus aurantium extract on L-type calcium currents in ventricular myocytes of single guinea pigs. *Human Yi Ke Da Xue Xue Bao* 2003; 4: 353-356.
5. Arbo MD, Franco MT, Larentis ER, Garcia SC, Sebben VC, Leal MB, et al. Screening for in-vivo (anti)estrogenic activity of ephedrine and p-syneprine and their natural sources Ephedra sinica Stapf. *Arch Toxicol* 2009; 83: 95-99.
6. Ahangarpour A, Oroojan AA. The effects of Cassia italica leaves aqueous extract on non-pregnant uterus contraction in rats. *Iran J Reprod Med* 2010; 4: 179-184.
7. Artech ME, Ausina P, Delgado J, Fernandez JJ, Nort M, Candenas ML, et al. Inhibitory effects of okadaic acid on rat uterine contractile responses to different spasmogens. *Fundam Cilm Pharmacol* 1997; 11: 47-56.
8. Yeh YY, Liu L. Cholesterol-lowering effect of garlic extracts and organosulfur compounds: human and animal studies. *J nutr* 2001; 131: 989- 993.
9. Gharib Naseri MK, Mazlomi H, Goshairesh M, Vakilizadeh G, Heidari A. Antispasmodic Effect of Zataria multiflora Boiss. Leaf Extract on the Rat Uterus. *Iran J Pharm Res* 2006; 2: 131-136.
10. Gharib Naseri MK, Mohammadian M, Gharib Naseri Z. Antispasmodic effect of Physalis alkekengi fruit extract on rat uterus. *Iran J Reprod Med* 2008; 4: 193-198.
11. Tolszczuk M, Pelletier G. Autoradiographic localization of beta-adrenoreceptors in rat uterus. *J Histochem Cytochem* 1988; 36: 1475-1479.
12. Zhu Y, Pintar JE. Expression of Opioid Receptors and Ligands in Pregnant Mouse Uterus and Placenta1. *Biol Reprod* 1998; 59: 925-932.
13. Ohia SE, Laniyonu AA. Naloxone-insensitive inhibitory and excitatory effects of opioid agonists in the rat isolated uterus. *J Pharm Pharmacol* 1989; 41: 168-172.
14. Zhang WW, Li Y, Wang XQ, Tian F, Cao H, Wang MW, et al. Effects of magnolol and honokiol derived from traditional Chinese herbal remedies on gastrointestinal movement. *World J Gastroenterol* 2005; 11: 4414-4418.
15. Chiwororo WD, Ojewole JA. Spasmolytic effect of psidium guajava Linn (Myrtaceae) leaf aqueous extract on rat isolated uterine horns. *J Smooth Muscle Res* 2009; 1: 31-38.
16. Ponce-Monter H, Perez S, Zavala MA, Perez C, Meckes M, Maclas A, et al. Relaxant effect of xanthomicrol and 3a-angeloyloxy-2a-hydroxy 13,14 zdehydrocaticvic acid from Brickellia paniculata on rat uterus. *Biol Pharm Bull* 2006; 29: 1501-1503.
17. Zhou XB, Lutz S, Steffens F, Korth M, Wieland T. Oxytocin receptors differentially signal via Gq and Gi proteins in pregnant and nonpregnant rat uterine myocytes: implications for myometrial contractility. *Mol Endocrinol* 2007; 21: 740-752.

18. Sanborn BM. Hormones and calcium: mechanisms controlling uterine smooth muscle contractile activity. The Litchfield Lecture. *Exp Physiol* 2001; 86: 223- 237.
19. Huang Y. BaCl<sub>2</sub>- and 4-aminopyridine-evoked phasic contractions in the rat vas deferens. *Br J Pharmacol* 1995; 115: 845-851.