

Short communication

Evaluation the frequency of factor V Leiden mutation in pregnant women with preeclampsia syndrome in an Iranian population

Samieh Karimi¹ M.D., Majid Yavarian² D.M.T., Ph.D., Azadeh Azinfar¹ M.D., Minoos Rajaei¹ M.D., Maryam Azizi Kootanaee¹ M.D.

1 Department of Obstetrics and Gynecology, Hormozgan Fertility and Infertility Research Center, Shariati Hospital, Hormozgan University of Medical Sciences, Bandar Abbas, Iran.

2 Department of Human Molecular and Clinical Genetics, Shiraz University of Medical Sciences, Shiraz, Iran.

Received: 13 November 2010; accepted: 18 May 2011

Abstract

Background: Role of genetic factors in etiology of preeclampsia is not confirmed yet.

Objective: Gene defect frequency varies in different geographic areas as well as ethnic groups. In this study, the role of factor V Leiden mutation in the pathogenesis of preeclampsia syndrome among the pregnant population of northern shore of Persian Gulf in Iran, were considered.

Materials and Methods: Between Jan. 2008 and Dec. 2009, in a nested case control study, pregnant women with preeclampsia (N=198) as cases and healthy (N=201) as controls were enrolled in the study. DNA were extracted from 10 CC peripheral blood and analyzed for presence of factor V Leiden mutation in these subjects. The maternal and neonatal outcomes of pregnancy according to the distribution of factor V Leiden were also compared among cases.

Results: In total, 17(8.6%) of cases and 2(1%) of controls showed the factor V Leiden mutation. The incidence of factor V Leiden was typically higher in preeclamptic women than control group (OR: 9.34 %95 CI: 2.12-41.01). There was no difference in incidence rate of preterm delivery < 37 weeks (OR: 1.23 %95 CI: 0.38-4.02), very early preterm delivery < 32 weeks (OR: 1.00 %95 CI: 0.12-8.46), intra uterine fetal growth restriction (IUGR) (OR: 1.32 %95 CI: 0.15-11.30), and the rate of cesarean section (OR: 0.88 %95 CI: 0.29-2.62) among cases based on the prevalence of factor V Leiden mutation.

Conclusion: The pregnant women with factor V Leiden mutation are prone for preeclampsia syndrome during pregnancy, but this risk factor was not correlated to pregnancy complications in the studied women.

Key words: Preeclampsia, Factor V Leiden mutation, Obstetric complication, Iran.

Introduction

The preeclampsia syndrome is a precarious condition with a challenge in medical treatment

Corresponding Author:

Azadeh Azinfar, Hormozgan Fertility and Infertility Research Center, Department of Obstetrics and Gynecology, Shariati Hospital, Hormozgan University of Medical Sciences, Bandar Abbas, Iran.

Email: aazinfar@yahoo.com

(1, 2). Two different kinds of treatment have suggested facing with this disastrous condition. One of the treatments is considering expectant management and another is placenta delivery. Considering expectant management especially for preeclampsia syndrome with occurrence before 24 weeks of gestation has reported to be associated with high maternal and fetus morbidity and mortality even though in tertiary referral centers.

On the other hand, however, placenta delivery is a rescue medical intervention for mother but it threatens the neonatal survival despite spending enough medical care in well- developed medical centers due to early preterm delivery and prematurity, consequently (3-6).

So, it seems to be efficient to find the ways to prevent the occurrence of preeclampsia syndrome during pregnancy especially in early stages. Some new data support from an increased tendency for coagulation in the pathogenesis of this syndrome. The mutation in the amino acid 506 (R506Q) of factor V gene that is known as Leiden is the most common genetic disorder is assigned pregnant women in a prethrombotic state. In this mutation, Adenine in nucleoside 1691 is substituted with guanine, which results to produce an abnormal activated protein C with decreased anticoagulant activity in subjects with this inherited thrombophilic defect (7, 8).

Recently it was found that women with this variant allele are at increased risk for pregnancy complications such as: preeclampsia, early onset of preeclampsia, very early preterm delivery, intra uterine growth restriction, low birth weight, small for gestational age, early and late recurrent fetal loss, still birth, placental abruption and susceptibility to recurrence of preeclampsia (9-17); however other studies have not support this hypothesis (18-22).

Therefore, it seems to be important to investigate the role of prethrombotic factors in the etiology of preeclampsia syndrome and also anticoagulant therapy like heparin therapy to prevent the occurrence of this syndrome in vulnerable women. For this reason, we designed this study to find whether the frequency of Leiden mutation is higher in patients with preeclampsia syndrome and this factor is responsible for higher obstetrics complications in these subjects in an Iranian pregnant population.

Materials and methods

Subjects

Between Jan 2008 and Dec 2009, a nested case control study was conducted to assay the correlation between factor V Leiden mutation and preeclampsia syndrome in pregnant women who were attended to Shariati women hospital for the

prenatal care or delivery in Bandar Abbas, Iran. In this study, 198 pregnant women with preeclampsia were enrolled as cases and 201 healthy pregnancies were considered as controls that were matched according to the age, ethnicity, and number of gravidity.

Cases with multiple pregnancy, molar pregnancy, diabetes mellitus, chronic hypertension, renal diseases, and systemic lupus erythematosus, past history of hypertension, recurrent abortion or thromboembolic events were excluded from the study. Such exclusion criteria were also set for 198 matched controls.

After completing demographic data sheet as well as exclusion criteria form for each participant, the systolic and diastolic blood pressure were measured as twice at least 6 hour apart by an appropriate-sized cuff. Then, 10 cc blood samples were obtained from all subjects for DNA analysis to assay the presence of factor V Leiden mutation in these two groups.

All women were observed for neonatal and maternal outcomes including gestational age at delivery, preterm delivery <37 weeks, very early preterm delivery <32 weeks, early onset of preeclampsia <30 weeks, intra uterine growth restriction, intra uterine fetal death and also cesarean section or normal vaginal delivery at the end of pregnancy. Finally, the pregnancy outcomes were compared among cases in according to the FV Leiden mutation.

In this study, the criteria from the American college of obstetrics and gynecologists were used to detection preeclampsia syndrome in studied population (23). In brief, systolic blood pressure of 140 mm hg or higher or a diastolic blood pressure of 90 mm hg or higher, association of 300 mg protein in 24-hour urine or $\geq +1$ on two random urine sample dipsticks at least 6 hour apart and also no more than 1 week apart that typically developed after 20 weeks gestation in women without a past history of chronic hypertension was considered as preeclampsia.

Preeclampsia associated with heavy proteinuria $\geq 5\text{gr}/24\text{hour}$ or $\geq +3$ on two dipsticks, oliguria $\leq 500\text{cc}/24\text{hour}$, platelet count $\leq 100'000$, elevated liver enzymes with persistent epigastric or right upper quadrant pain, pulmonary edema, blood pressure $\geq 160/110$, severe headache or visual disturbance recognized as severe preeclampsia.

Eclampsia is defined as occurrence of seizure without any identifiable etiology in preeclamptic women and Hellp syndrome as hemolysis, elevated liver enzymes and low platelet counts in studied patients.

Pregnant women with Hellp syndrome or eclampsia was classified as severe preeclampsia. In this study, we recorded IUGR as child birth weight <10 the percentile for sex and gestational age based twice sonography 2 weeks apart. The gestational age was measured using last menstrual period (LMP) and if it was unknown, we used ultra sonography instead. The ethics committee of Hormozgan University of Medical Sciences approved the study and signed informed consent was obtained from all subjects to participate at the study.

Molecular Study

To determine of FV Leiden mutation, Genomic DNA extraction was performed following protocol described by Miller *et al* (24). PCR amplification was carried out in 25 µl reaction volume containing 0.5 units Taq polymerase, 200 µM dNTP, 500 µM of each of previously described primers. After initial denaturation, 35 cycles at 95°C for 30s, 57°C 30s, and 72°C for 20s and followed extension by 72 for 10 min were performed. About 10 µl of PCR product digest with MNI I restriction enzymes (25) and result was analyzed by agarose gel electrophoresis containing ethidium bromide to find Wild Genotype: G/G, Mutant Genotype (Heterozygote): G/A and Mutant Genotype (Homozygote): A/A. (G=guanine, A=Adenine)

Statistical analysis

The SPSS software (version13, Inc, Chicago, Illinois, USA) was used to analysis the collected data. The chi-squared test was applied to determine the correlation between factor V Leiden mutation and preeclampsia in this study and also we performed t-test to compare quantitative variables between cases and controls. The correlation between Leiden mutation and the incidence rates of IUGR and IUFD were examined by Fisher's Exact Test. A p-value<0.05 was considered significant statistically and odds ratio (OR) with %95 confidence interval was calculated for pregnancy complication variables on Iranian

population curve and we observed IUFD as not detectable fetal heart rate in at least

Results

Three women who met the exclusion criteria were drawn from the study cases. The age, race, and number of gravidity were not different among cases and controls. Duration of pregnancy was lower in patients than control group that it was expectable. The demographic data and clinical characteristics of study patients are presented in table I. The prevalence of factor V Leiden was higher in cases 17 (8.6%) than control group 2 (1%) that it was statistically significant (OR: 9.34 %95 CI: 2.12-41.01).

Table II shows the frequency of normal genotype and genotypes with mutant allele in cases and controls. Of the 46 (23.2%) studied women as severe preeclampsia, 3 patients had Hellp syndrome and 6 of them were referred to the hospital with eclampsia. Leiden mutation was only found in 3 patients with severe preeclampsia. There wasn't any association between severity of preeclampsia and factor V Leiden mutation (OR: 0.68 %95 CI: 0.18-2.50).

The maternal and neonatal outcomes among the carriers of factor V Leiden are summarized in table III. The frequency of preterm delivery<37 weeks (OR: 1.23 %95 CI: 0.38-4.02) and very early preterm delivery<32 weeks (OR: 0.83 %95 CI: 0.10-7.03) were not higher in women with Leiden mutation than other cases without the mutation. Nine cases delivered IUGR that the majority of them 7 (77.8%) had severe preeclampsia. Three cases with wild type factor V were found with IUFD. The incidence rates of IUGR and IUFD among cases were not correlated to the presence of factor V gene mutation (Table III).

Of the 198 cases studied, 135 (68.2%) had normal vaginal delivery and 63 (31.8%) underwent cesarean section. In patients with cesarean section, 5 (2.5%) had factor V mutation and rest of them were negative for the mutant allele 58 (29.3%). The mode of delivery between cases based on Leiden mutation was not statistically different (Table III). In the present study, the correlation between Leiden mutation and severity of proteinuria as an indicator of the presence of factor V Leiden was examined but it was not significant (Table III).

Table I. Demographic data of cases and controls.

	Cases n=198	Controls n=201	p-value
Maternal age (years)	25.98±5.78 (15-40)	25.98±5.54 (14-40)	0.99
Gestational age at delivery (weeks)	36.22±2.70 (22-39)	37.52±0.96 (34-42)	0.00
Gravidity (mean)	2.28±1.74 (1-10)	2.27±1.45 (1-9)	0.95
Systolic blood pressure (mm hg)	147.67±11.02	109.43±10.56	0.00
Diastolic blood pressure (mm hg)	96.87±8.47	72.81±7.44	0.00

Table II. Frequency of normal genotype and genotypes with mutant allele in studied patients.

Studied population genotype	G/G (n%)	G/A (n%)	A/A (n%)
Preeclampsia women (n=198)	181 (91.4%)	17 (8.6%)	0 (0%)
Controls (n=201)	199 (99%)	2 (1%)	0 (0%)

Wild genotype: G/G

Mutant genotype (Heterozygote): G/A

Mutant genotype (Homozygote): A/A

Table III. Maternal and neonatal outcomes according to the presence of FV Leiden mutation among cases.

FVL Mutation	Yes (n%)	No (n%)	p-value	Odds ratio (%95 CI)
Gestational age at delivery (weeks)	36.12±2.69	36.23±2.71	0.87	-
Preterm delivery < 37 weeks	4 (2%)	36 (18.2%)	0.72	1.23 (0.38-4.02)
Very early preterm delivery <32 weeks	1 (0.5%)	9 (4.5%)	0.87	0.83 (0.10-7.03)
IUGR	1 (0.5%)	8 (4%)	0.56	1.32 (0.15-11.30)
IUFD	0 (0%)	3 (1.5%)	0.76	-
Cesarean section	5 (2.5%)	58 (29.3%)	0.82	0.88 (0.29-2.62)
Early onset of preeclampsia < 30 weeks	0 (0%)	9 (4.5%)	0.34	-
Sever preeclampsia	3 (1.5%)	43 (21.7%)	0.56	0.68 (0.18-2.50)
Proteinuria ≥ +3 or more	2 (1%)	19 (9.6%)	0.87	1.13 (0.24-5.35)

Discussion

The role of genetic factors in the etiology of preeclampsia is not confirmed yet. However in recent decades, several studies have shown that pregnant women who are carriers of factor V Leiden mutation have faced with an increased risk of preeclampsia in their pregnancy but other studies have not support this hypothesis. Some studies reported the prevalence of factor V Leiden mutation varies among a nation to other nation and it's observed the lowest prevalence of the mutation among Asian nations especially in Indonesian and Japanese population (26-29).

It is notable to mention that there are few studies that have assessed the frequency of Leiden mutation among Iranians. In one study on Kurdish

population in western Iran, it was reported the prevalence of 2.97% in 434 healthy subjects and in another study in South of Iran by Karimi *et al* it was obtained 4.1% among 198 volunteers healthy and 6.42% in Kuwaitis of Iranian origin by Dashti *et al* (30-32).

The present study investigated whether the factor V Leiden mutation is able to predispose pregnant women to preeclampsia and also to alter the pregnancy outcome in these subjects in Iran where is located in the western Mediterranean region and to our knowledge, there are only one study with a very small number of cases (33) that have assessed this correlation in this region of the world.

The overall prevalence of Leiden mutation in preeclamptic women was obtained 8.6%, which

was comparable with the reported prevalence of 6% (2/33), 8.9% (4/21) and 10% (4/40) in Davalos, Dashti, Dizon-Townson and Prasmusinto *et al* studies, respectively (34-36). In this study, there was statistically a significant correlation between Leiden mutation and the presence of preeclampsia syndrome in Iranian pregnant women.

This result is in agreement with findings from the several large meta-analysis studies that have explored to determine any significant association between the factor V Leiden mutation and the presence of preeclampsia during pregnancy. In a meta-analysis consist of cohort studies, Dudding *et al* (37) reported that it seems the maternal factor V Leiden mutation is able to increase the risk of preeclampsia until fifty percent levels (50%) and Kosmas *et al* (38) assessed results from 19 studies that have investigated to find this correlation and identified women with this mutation are at increased risk for preeclampsia 2.5 fold higher than women without the mutation.

In a similar study by Facchinetti *et al* (11), they genotyped 172 cases with a previous history of preeclampsia for common heritable thrombophilic genes mutations and observed women with a thrombophilic gene mutation are susceptible to achieve preeclampsia during their next pregnancy more than 2.5 times higher than controls. In another study by Dalmaç *et al* (39) in Brazilian population, they considered the correlation between methylenetetrahydrofolate reductase (MTHFR), Prothrombin Mutation (FII), Plasminogen activator inhibitor (PAI-1) and also factor V Leiden mutation and mild/sever preeclampsia in 75 pregnant women.

They inconcluded from the study the presence of these mutations alone are not able to predispose pregnant women for preeclampsia but a combination of them appears to be correlated with the development of preeclampsia syndrome in pregnancy.

In contrast to our findings, many studies didn't find any significant correlation between factor V Leiden mutation and increased rate of preeclampsia among pregnant women with this variant allele. Davalos *et al* (34) reported no significant association between factor V Leiden mutation and preeclampsia in 33 Mexican women that compared to 62 normotensive pregnant women as controls. Salomon *et al* (19) study showed no

increased rate of preeclampsia in 191 women with common prothrombotic genes compared to 446 controls (4.3% Vs. 5.25 p=0.59).

Two recent large nested case-control studies couldn't reveal such association, too. Hiltunen *et al* (6) studied 100000 pregnant women in Finland and found however the presence of factor V Leiden mutation increased the risk of preeclampsia 1.7 fold higher than controls but this association was not statistically significant. In the Montreal preeclampsia study that included 5332 women by Kahn *et al* (40), it was reported no difference in the prevalence of factor V Leiden mutation among preeclamptic women and controls.

In another interesting survey, two different ethnical nations (Indonesian and Germany/Croatians women) were assessed to clarify any correlation between factor V Leiden mutation and preeclampsia in a case-control study by Prasmusinto *et al* (36), while this correlation was significant in Germany/Croatians women, they couldn't find such association among Indonesian women because it was not found any women with factor V Leiden in Indonesian cases and control group. The findings from this study imply that the ethnicity plays an important role in the examination such correlation among different nations.

In the current study, the frequency of preterm delivery, very early preterm delivery, early onset of preeclampsia, IUGR, IUFD and also the mode of delivery were not different among cases in according to the distribution of factor V Leiden mutation. The data obtained from the some studies support this finding. In a survey by zahed *et al* (41), they found the factor V Leiden; prothrombin and methylenetetrahydrofolate reductase mutations are not predictable risk factors of disastrous pregnancy outcomes in Lebanese women.

Salomon *et al* and Dudding *et al* (15, 37) in their studies couldn't find evidence regarding the higher incidence of IUGR among women with the factor V Leiden mutation while it's reported the IUGR and IUFD were more prevalent in women with the Leiden mutation by Nurk *et al* (9) and in other similar studies (8, 11). Moreover it was observed the higher prevalence of very early preterm delivery (less than 32 weeks) in these subjects in Facchinetti *et al* (7) study contrasting our study.

We expected to observe the high rate of cesarean section in preeclampsia cases with the mutation based on some previous reports that recorded the higher rate of preeclampsia complications among these patients, but it was not significant when compared to cases without the mutation in our study.

The weakness of the study was the small number of studied patients and it was done on a population of Iranian pregnant women that maybe different from other nations genetically, hence other studies with large sample size are needed to examine such association in another ethnical nations.

In the end, it is important to mention however at the present, the role of heritable thrombophilia in the pathogenesis of preeclampsia is not definitely established but a few studies explored that heparin therapy is effective to reduce the adverse pregnancy outcomes in women carrying the V Leiden mutation (5, 42).

In short, the study showed a significant correlation between the occurrence of preeclampsia and mutation in the factor V Leiden that it's necessary to be confirmed by further studies in other areas especially in Iran. The authors emphasize to this important point that the obtained data may be to be different from other nations due to genetic diversity among different nations.

Acknowledgments

The authors would like to thank all of the pregnant women involved in the study and special thank the staff of Shiraz pastor clinical laboratory.

References

- Irminger-Finger I, Jastrow N, Irion O. Preeclampsia: A danger growing in disguise. *Int J Biochem Cell Biol* 2008; 40: 1979-1983.
- Saadat M, Nejad SM, Habibi G, Sheikhvatan M. Maternal and neonatal outcomes in women with preeclampsia. *Taiwan J Obstet Gynecol* 2007; 46: 255-259.
- Dildy GA 3rd, Belfort MA, Smulian JC. Preeclampsia recurrence and prevention. *Semin Perinatol* 2007; 31: 135-141.
- Bolte AC, van Geijn HP, Dekker GA. Management and monitoring of severe preeclampsia. *Eur J Obstet Gynecol Reprod Biol* 2001; 96: 8-20.
- Gaugler-Senden IP, Huijssoon AG, Visser W, Steegers EA, de Groot CJ. Maternal and perinatal outcome of preeclampsia with an onset before 24 weeks' gestation: Audit in a tertiary referral center. *Eur J Obstet Gynecol Reprod Biol* 2006; 128: 216-221.
- MacKay AP, Berg CJ, Atrash HK. Pregnancy-related mortality from preeclampsia and eclampsia. *Obstet Gynecol* 2001; 97: 533-538.
- Lachmeijer AM, Dekker GA, Pals G, Aarnoudse JG, ten Kate LP, Arngriímsson R. Searching for preeclampsia genes: The current position. *Eur J Obstet Gynecol Reprod Biol* 2002; 105: 94-113.
- Franchini M, Lippi G. Factor V Leiden and hemophilia. *Thromb Res* 2010; 125: 119-123.
- Paidas MJ, Ku D-HW, Langhoff-Roos J, Arkel YS. Inherited thrombophilias and adverse pregnancy outcome: Screening and management. *Semin Perinatol* 2005; 29: 150-163.
- Hiltunen LM, Laivuori H, Rautanen A, Kaaja R, Kere J, Krusius T, et al. Blood group ab and factor V Leiden as risk factors for pre-eclampsia: A population-based nested case-control study. *Thromb Res* 2009; 124: 167-173.
- Facchinetti F, Marozio L, Frusca T, Grandone E, Venturini P, Tiscia GL, et al. Maternal thrombophilia and the risk of recurrence of preeclampsia. *Am J Obstet Gynecol* 2009; 200: 41-45.
- Howley HE, Walker M, Rodger MA. A systematic review of the association between factor V Leiden or prothrombin gene variant and intrauterine growth restriction. *Am J Obstet Gynecol* 2005; 192: 694-708.
- Nurk E, Tell GS, Refsum H, Ueland PM, Vollset SE. Factor V Leiden, pregnancy complications and adverse outcomes: The hordaland homocysteine study. *QJM* 2006; 99: 289-298.
- Mtiraoui N, Borgi L, Hizem S, Nsiri B, Finan RR, Gris J-C, et al. Prevalence of antiphospholipid antibodies, factor V g1691a (Leiden) and prothrombin g20210a mutations in early and late recurrent pregnancy loss. *Eur J Obstet Gynecol Reprod Biol* 2005; 119: 164-170.
- Hiltunen LM, Laivuori H, Rautanen A, Kaaja R, Kere J, Krusius T, et al. Factor V Leiden as risk factor for unexplained stillbirth- a population-based nested case-control study. *Thromb Res* 2010; 125: 505-510.
- Glueck CJ, Gogenini S, Munjal J, Tracy T, Prankoff J, Wang P. Factor V Leiden mutation: A treatable etiology for sporadic and recurrent pregnancy loss. *Fertil Steril* 2008; 89: 410-416.
- Hussein AS, Darwish H, Shelbayeh K. Association between factor V Leiden mutation and poor pregnancy outcomes among palestinian women. *Thromb Res* 2010; 126: 78-82.
- Livingston JC, Barton JR, Park V, Haddad B, Phillips O, Sibai BM. Maternal and fetal inherited thrombophilias are not related to the development of severe preeclampsia. *Am J Obstet Gynecol* 2001; 185: 153-157.
- Salomon O, Seligsohn U, Steinberg DM, Zalel Y, Lerner A, Rosenberg N, et al. The common prothrombotic factors in nulliparous women do not compromise blood flow in the fetomaternal circulation and are not associated with preeclampsia or intrauterine growth restriction. *Am J Obstet Gynecol* 2004; 191: 2002-2009.
- Von Tempelhoff GF, Heilmann L, Spanuth E, Kunzmann E, Hommel G. Incidence of the factor V Leiden-mutation, coagulation inhibitor deficiency, and elevated antiphospholipid-antibodies in patients with preeclampsia or hellp-syndrome. *Thromb Res* 2000; 100: 363-365.
- Kim YJ, Williamson RA, Murray JC, Andrews J, Pietscher JJ, Peraud PJ, et al. Genetic susceptibility to preeclampsia: Roles of cytosine-to-thymine substitution at

- nucleotide 677 of the gene for methylenetetrahydrofolate reductase, 68-base pair insertion at nucleotide 844 of the gene for cystathionine β -synthase, and factor V Leiden mutation. *Am J Obstet Gynecol* 2001; 184: 1211-1217.
22. Nath CA, Ananth CV, DeMarco C, Vintzileos AM, New Jersey-Placental Abruptio Study Investigators. Low birthweight in relation to placental abruptio and maternal thrombophilia status. *Am J Obstet Gynecol* 2008; 198: 291-295.
 23. ACOG Committee on Obstetrics Practice. ACOG practice bulletin. Diagnosis and management of preeclampsia and eclampsia. Number 33, January 2002. American college of obstetricians and gynecologists. *Int J Gynaecol Obstet* 2002; 77: 67-75.
 24. Lucotte G, Mercier G. Population genetics of factor V Leiden in Europe. *Blood Cells Mol Dis* 2001; 27: 362-367.
 25. Miller SA, Dykes DD, Polesky HF. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 1988; 16: 1215.
 26. Beauchamp NJ, Daly ME, Hampton KK, Cooper PC, Preston FE, Peake IR. High prevalence Prothrombin G20210A was reported to be more common of a mutation in the factor V gene within the UK population: relationship to activated protein C resistance and familial thrombosis. *Br J Haematol* 1994; 88: 219-222.
 27. Bauduer F, Lacombe D. Factor V Leiden, prothrombin 20210a, methylenetetrahydrofolate reductase 677t, and population genetics. *Mol Genet Metab* 2005; 86: 91-99.
 28. Limdi NA, Beasley TM, Allison DB, Rivers CA, Acton RT. Racial differences in the prevalence of factor V Leiden mutation among patients on chronic warfarin therapy. *Blood Cells Mol Dis* 2006; 37: 100-106.
 29. Kobashi G, Yamada H, Asano T, Nagano S, Hata A, Kishi R, et al. The factor V Leiden mutation is not a common cause of pregnancy-induced hypertension in Japan. *Semin Thromb Hemost* 1999; 25: 487-489.
 30. Rahimi Z, Vaisi-Raygani A, Mozafari H, Kharrazi H, Rezaei M, Nagel RL. Prevalence of factor V Leiden (G1691A) and prothrombin (G20210A) among Kurdish population from Western Iran. *J Thromb Thrombolysis* 2008; 25: 280-283.
 31. Karimi M, Panahandeh Shahraki G, Yavarian M, Afrasiabi A, Dehbozorgian J, Bordbar M, et al. Frequency of factor V Leiden and Prothrombin Polymorphism in South of Iran. *Iran J Med Sci* 2009; 34: 137-140.
 32. Dashti AA, Jadaon MM. Race differences in the prevalence of the factor V Leiden mutation in Kuwaiti nationals. *Mol Biol Rep* 2011; 38: 3623-3628.
 33. Sadat Hosseini M, Gharekhani P, Sadeghi M. Association of inherited thrombophilia and antiphospholipid syndrome with severe preeclampsia. *Iran J Reprod Med* 2007; 5: 183-186.
 34. Dizon-Townson DS, Nelson LM, Easton K, Ward K. The factor V Leiden mutation may predispose women to severe preeclampsia. *Am J Obstet Gynecol* 1996; 175: 902-905.
 35. Dávalos IP, Moran MC, Martínez-Abundis E, González-Ortiz M, Flores-Martínez SE, Machorro V, et al. Methylenetetrahydrofolate reductase c677t polymorphism and factor V Leiden variant in Mexican women with preeclampsia/eclampsia. *Blood Cells Mol Dis* 2005; 35: 66-69.
 36. Prasmusinto D, Skrablin S, Fimmers R, van der Ven K. Ethnic differences in the association of factor V Leiden mutation and the c677t methylenetetrahydrofolate reductase gene polymorphism with preeclampsia. *Eur J Obstet Gynecol Reprod Biol* 2004; 112: 162-169.
 37. Dudding T, Heron J, Thakkinstian A, Nurk E, Golding J, Pembrey M, et al. Factor V Leiden is associated with preeclampsia but not with fetal growth restriction: A genetic association study and meta-analysis. *J Thromb Haemost* 2008; 6: 1869-1875.
 38. Kosmas IP, Tatsioni A, Ioannidis JP. Association of Leiden mutation in factor V gene with hypertension in pregnancy and pre-eclampsia: A meta-analysis. *J Hypertens* 2003; 21: 1221-1228.
 39. Dalmáz CA, Santos KG, Botton MR, Tedoldi CL, Roisenberg I. Relationship between polymorphisms in thrombophilic genes and preeclampsia in a Brazilian population. *Blood Cells Mol Dis* 2006; 37: 107-110.
 40. Kahn SR, Platt R, McNamara H, Rozen R, Chen MF, Genest J Jr, et al. Inherited thrombophilia and preeclampsia within a multicenter cohort: The Montreal preeclampsia study. *Am J Obstet Gynecol* 2009; 200: 151-159.
 41. Zahed LF, Rayes RF, Mahfouz RA, Taher AT, Maarouf HH, Nassar AH. Prevalence of factor V Leiden, prothrombin and methylene tetrahydrofolate reductase mutations in women with adverse pregnancy outcomes in Lebanon. *Am J Obstet Gynecol* 2006; 195: 1114-1118.
 42. Wang Y, Walli AK, Schulze A, Blessing F, Fraunberger P, Thaler C, et al. Heparin-mediated extracorporeal low density lipoprotein precipitation as a possible therapeutic approach in preeclampsia. *Transfus and Apher Sci* 2006; 35: 103-110.

