

9th Yazd International Congress and Student Award on Reproductive Medicine with 4th Congress of Reproductive Genetics

Poster Presentations

P-60

Investigation and comparison of the effect of TGF- β 3, kartogenin and Avocado/Soybean unsaponifiables on the in-vitro and in-vivo chondrogenesis of human adipose derived stem cells on fibrin scaffold

Hashemibeni B¹, Izadi MA¹, Valiani A¹, Esfandiari I¹, Bahramian H¹, Dortaj H², Pourentezari M³.

1.Department of Anatomical Sciences and Molecular Biology, Isfahan University of Medical Sciences, Isfahan, Iran.

2.Department of Tissue Engineering and Applied Cell Science, Shiraz University of Applied Medical Science and Technologies, Shiraz, Iran.

3.Department of Biology and Anatomical Sciences, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

Email: m.pourentezari@gmail.com

Background: There are lack of suitable therapeutic approaches to cartilage defect.

Objective: This study was designed to determine the effect of TGF- β 3, avocado/soybean (ASU) and Kartogenin (KGN) on chondrogenic differentiation in human adipose-derived stem cells (hADSCs) on fibrin scaffold.

Materials and Methods: hADSCs seeded in fibrin scaffold and cultured in chondrogenic media. These cells divided in to 4 groups (control, TGF- β 3, ASU and KGN). Cell viability was estimated by 3-(4, 5-

dimethylthiazol-2-yl)-2, 5-diphenyltetrazolium bromide, or MTT assay, differentiated cells evaluated by histological and immunohistochemical techniques. Expression genes [sex determining region Y-box 9 (SOX9), Aggrecan (AGG), type II collagen (Coll II) and type X collagen (Coll X)] assessed by real-time PCR. For study on animal model, differentiated cells in fibrin scaffolds were subcutaneously transplanted in rats. Histological and immunohistochemistry was done in animal model.

Results: The results of the real-time PCR indicated that SOX9, AGG and *Col II* genes expression in TGF- β 3, KGN and ASU groups were significantly higher ($p < 0.01$) compared to the control group, *Col X* gene expression only in TGF- β 3 group was significantly higher ($p < 0.01$) compared to the control group. The glycosaminoglycan (GAG) deposition was higher in TGF- β 3, KGN and ASU groups compared to the control group. The immunohistological analysis showed the distribution of collagen type X in the extracellular matrix in fibrin scaffold TGF- β 3 group was significantly higher in control, KGN and ASU groups, ($p < 0.001$).

Conclusion: ASU, and in particular KGN was suitable for successful chondrogenic differentiation of hADSCs and a suppressor of the consequent hypertrophy.

Key words: TGF β 3, Avocado/Soybean, Kartogenin, Human adipose-derived stem cells.