

## 9<sup>th</sup> Yazd International Congress and Student Award on Reproductive Medicine with 4<sup>th</sup> Congress of Reproductive Genetics

### Poster Presentations

#### P-82

#### Protective effects of curcumin on sperm and stereological parameters in testes of formaldehyde-exposed NMRI mice: An experimental study

Fathi E<sup>1</sup>, Shahedi A<sup>1</sup>, Hosseini Sharifabad M<sup>1</sup>, Vakili M<sup>2</sup>.

1. Department of Biology and Anatomy, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

2. Department of Community Medicine, Faculty of Medicine, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

Email: elahe72.fathi@gmail.com

**Background:** Formaldehyde (FA) exposure is one of the important causes of cellular injury and oxidative damage in testis, and leads to infertility.

**Objective:** This study aimed was to assess the protective effects of curcumin on sperm and stereological parameters in testes from formaldehyde-exposed NMRI mice.

**Materials and Methods:** At 6-8 weeks of age, 24 adult male NMRI mice weighing 30-35 g were separated into four groups (n = 6) based on the treatment they received: Group I (control) no treatment, group II received FA (10 mg/kg), group III received FA (10 mg/kg) and curcumin (100 mg/kg), and group IV (Solvent) received dimethyl sulfoxide (DMSO) (0.2 ml/day). Materials were

administered intraperitoneally for 35 days. After excision, epididymis tissues were placed in 1-ml aliquots of Ham's F10 medium at 37°C for 20 min and were then used in analyses of sperm parameters. Testes were fixed and stained with Hematoxylin and Eosin (H & E) for investigations of stereological indices. We also determined lipid peroxidation levels using malondialdehyde (MDA) assays.

**Results:** Mean sperm parameters (count, motility, viability, and morphology) differed significantly between groups II and III ( $p \leq 0.001$ ). Stereological indices, including Leydig and spermatogonia cell numbers and surface-to-volume ratios of seminiferous tubules were significantly higher in group III than in group II ( $p \leq 0.001$ ). Finally, MDA levels in group III were significantly lower than in group II ( $p = 0.001$ ).

**Conclusion:** Our data showed that the curcumin, as an antioxidant, reduced FA-induced damage in sperm parameters and stereological indices in mice testis.

**Key words:** Formaldehyde, Curcumin, Testis, Mice.

*The original full text of this abstract has been published in International Journal of Medical Laboratory 2021; 8(1): 35-43. <https://doi.org/10.18502/ijml.v8i1.5671>.*

*How to cite to this article: Fathi E, Shahedi A, Hosseini Sharifabad M, Vakili M. Protective effects of curcumin on sperm and stereological parameters in testes of formaldehyde-exposed NMRI mice: An experimental study. International Journal of Medical Laboratory 2021; 8(1): 35-43.*