

## 9<sup>th</sup> Yazd International Congress and Student Award on Reproductive Medicine with 4<sup>th</sup> Congress of Reproductive Genetics

### Poster Presentations

P-37

#### The association between fatty acids and steroids gene expression in abdominal subcutaneous fat depot: A comparison between pregnant PCOS and non-PCOS pregnant women

Emami N<sup>1, 2</sup>, Alizadeh AR<sup>2</sup>, Moini A<sup>3, 4, 5</sup>, Yaghmaei P<sup>1</sup>, Shahhosseini M<sup>6, 7, 8</sup>.

1. Department of Biology, Faculty of Science, Science and Research Branch, Islamic Azad University, Tehran, Iran.

2. Department of Embryology, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran.

3. Department of Endocrinology and Female Infertility, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran.

4. Breast Disease Research Center (BDRC), Tehran University of Medical Sciences, Tehran, Iran.

5. Department of Gynecology and Obstetrics, Arash Women's Hospital, Tehran University of Medical Sciences, Tehran, Iran.

6. Reproductive Epidemiology Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran.

7. Department of Genetics, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran.

8. Department of Cell and Molecular Biology, School of Biology, College of Science, University of Tehran, Tehran, Iran.

Email: neda.emami58@gmail.com

**Background:** It was reported that steroid-related gene expressions in the adipose tissue (AT) of women differ between women affected with polycystic ovary syndrome (PCOS) (case) and non-PCOS (control). Although association between PCOS in mother and offspring's health is a crucial issue, there are few studies focusing on effectiveness of AT profiles on

steroids genes expression in pregnant women suffering from PCOS.

**Objective:** Our objectives were to assess association between fatty acid (FA) and genes related to steroids metabolism expression in abdominal subcutaneous AT of 12 PCOS (case) vs. 32 non-PCOS (control) age- and BMI-matched pregnant women.

**Materials and Methods:** Twelve pregnant women with PCOS (case) and thirty two non-PCOS pregnant women (control) (age- and BMI-matched) undergoing cesarean section were enrolled for the present study. Expressions of fifteen genes related to steroidogenesis in abdominal subcutaneous AT were investigated using quantitative real-time PCR. Fatty acids profiles assessed by gas chromatography. Linear regression was performed to determine the association of FA and gene expression in subcutaneous AT.

**Results:** Age and BMI were similar among two groups at delivery day. Current study showed that omega-3 fatty acids had the highest association with steroids gene expression rate ( $r = 0.500$ ;  $p < 0.05$ ).

**Conclusion:** It seems that fatty acids, both direct and by metabolites, can play a role in many diseases through extensive signaling pathways, specifically in exacerbating PCOS, although pregnancy can double the role of nutrition in exacerbating these effects.

**Key words:** Polycystic ovary syndrome, Subcutaneous adipose tissue, Sex steroid, Fatty acids.

The original full text of this abstract has been published in *Adipocyte* 2020 Jan 1; 9(1): 16-23. <https://doi.org/10.1080/21623945.2019.1710021>.

How to cite to this article: Emami N, Alizadeh A, Moini A, Yaghmaei P, Shahhosseini M. Differences in fatty acid profiles and desaturation indices of abdominal subcutaneous adipose tissue between pregnant women with and without PCOS. *Adipocyte* 2020; 9(1): 16-23.