

9th Yazd International Congress and Student Award on Reproductive Medicine with 4th Congress of Reproductive Genetics

Poster Presentations

P-16

Antioxidant effects of royal jelly on lead induced sperm DNA damage and sperm abnormality in male mice

Kakebaraei S¹, Jalili C².

1. Student Research Committee, Kermanshah University of Medical Sciences, Kermanshah, Iran.

2. Department of Anatomy, School of Medicine, Kermanshah Medical University, Kermanshah, Iran.

Email: seyranbaraie@yahoo.com

Background: Royal jelly, a secretion vintage of the salivary glands of worker bees, is an extremely impressive antioxidant and possesses eminent free radical scavenging trait. It has been shown to have anti-tumor, antibiotic, anti-inflammatory, immunomodulatory and as well as antioxidant properties. Royal jelly has remarkable positive effects on reproductive system and fertility. On the other hand, lead, which is widely used in industry, can be detected in foods, drinking water, ambient air, dust, and various cosmetics products. Its toxicity causes excess levels of reactive oxygen species (ROS) and cellular oxidative stress. More importantly, our knowledge of the effects of lead on sperm DNA integrity is significantly limited.

Objective: We investigated the effect of Royal jelly

against reproductive toxicity caused by lead exposure in mice.

Materials and Methods: Male mice received lead acetate (50 mg/kg) and plus Royal jelly (25, 50 or 100 mg/kg) via oral gavage for 28 days. Adult male mice were divided into five groups (n = 5). Caudal epididymal sperm characteristics, lipid peroxidation and in vitro fertilizing capacity were evaluated after 4 weeks.

Results: Lead acetate-intoxicated mice exhibited testicular tissue injury and decreased serum levels of SOD, TAC, testosterone and increased serum level of MDA and nitric oxide. The count, viability, motility and normal morphology of the sperms were decreased in lead-induced group. Royal jelly prevented testicular injury, increased serum levels of SOD, TAC and improved the semen quality and decreased serum level of MDA and nitric oxide. However, Royal Jelly can reduce the regulation of Bax and Caspase-3 pro-inflammatory factors in lead-treated mice by reducing oxidative stress.

Conclusion: Our findings showed that Royal jelly with its antioxidant effects reduces inflammation and cell death in testis and sperm DNA structure following exposure to lead. However, further studies are needed to illuminate other mechanisms of Royal jelly's effect on testis function.

Key words: Royal jelly, Lead acetate, Sperm, Caspase3, Bax.