

## 9<sup>th</sup> Yazd International Congress and Student Award on Reproductive Medicine with 4<sup>th</sup> Congress of Reproductive Genetics

### Oral Presentations

#### O-13

#### Evaluation the effect of human sperm incubation time in polyvinylpyrrolidone on sperm structure reactive oxygen species, acrosome reaction, and mitochondrial membrane potential

Sabour M<sup>1</sup>, Agha-Rahimi A<sup>2</sup>, Dehghani Ashkezari M<sup>1</sup>, Seifati SM<sup>1</sup>, Kalantar SM<sup>3</sup>, Anbari F<sup>2</sup>, Nabi A<sup>2</sup>.

1. Medical Biotechnology Research Center, Ashkezar Branch, Islamic Azad University, Ashkezar, Yazd, Iran.

2. Research and Clinical Center for Infertility, Yazd Reproductive Sciences Institute, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

3. Abortion Research Center, Yazd Reproductive Sciences Institute, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

Email: 63rahimi@gmail.com

**Background:** Polyvinylpyrrolidone (PVP) is a chemical used in intracytoplasmic sperm injection for sperm immobilization. In human sperm, PVP has been shown to damage sperm membranes, DNA integrity, mitochondrial membrane, and destroy axonal tubules and fibrous sheaths.

**Objective:** The aim of this study was to investigate the

ideal time that sperm can be safely incubated in PVP with less possible damage.

**Materials and Methods:** Twenty-five normospermic samples were used. Sperm samples were prepared by swim-up method. Sperm samples incubated in 10% PVP at different time intervals (0, 15, 30, and 60 min). The effect of PVP was assessed on sperm structure, reactive oxygen species, acrosome reaction, Mitochondrial Membrane potential at different time intervals.

**Results:** Sperm parameters, DNA integrity and chromatin quality in 15, 30 and 60 min after incubation sperm with PVP were significantly changed compared to the 0 min. Moreover, in 30 and 60 min after incubation with PVP, above parameters were significantly changed compared to the 15 min. 60 min after incubation sperm with PVP, these parameters were significantly changed compared to the 30 min.

**Conclusion:** Sperm samples could be incubated with PVP for 15 min with less possible damage. While, prolonged incubation may damage the sperm parameters, DNA integrity and chromatin quality significantly.

**Key words:** Polyvinylpyrrolidone, Sperm mitochondria, Sperm ROS.