

## 9<sup>th</sup> Yazd International Congress and Student Award on Reproductive Medicine with 4<sup>th</sup> Congress of Reproductive Genetics

### Oral Presentations

#### O-52

#### Endometrial scratching affects gene expression of *NLRP3* in patients with unexplained repeated implantation failure: A randomized control trial

Aghajanpour S<sup>1</sup>, Hosseini E<sup>2</sup>, Amirchaghmaghi E<sup>3</sup>, Zandieh Z<sup>4</sup>, Amjadi F<sup>4</sup>, Yahyaei A<sup>1</sup>, Ashrafi M<sup>1</sup>, Aflatoonian R<sup>1</sup>.

1. Department of Endocrinology and Female Infertility, Reproductive Biomedicine Research Center, Royan Institute for Reproductive Biomedicine, ACECR, Tehran, Iran.

2. Department of Obstetrics and Gynecology, IVF Clinic, Mousavi Hospital, School of Medicine, Zanzan University of Medical Sciences, Zanzan, Iran.

3. Department of Regenerative Biomedicine, Cell Science Research Center, Royan Institute for Stem Cell Biology and Technology, ACECR, Tehran, Iran.

4. Department of Anatomical Sciences, School of Medicine, Iran University of Medical Sciences, Tehran, Iran.

Email: r.aflatoonian@gmail.com

**Background:** Alternative strategies have been used to augment success rate of implantation in IVF/ICSI cycles in unexplained repeated implantation failure patients. Endometrial scratching is one of these procedures. It seems scratching can affect *NLRP3* gene expression which has an important role on receptivity of endometrium. *NLRP3* is an intracellular sensor that detects a broad range of harmful sterile or infectious stimuli, resulting in the formation and activation of the *NLRP3* inflammasome

**Objective:** In the present study, we investigated whether gene expression of *NLRP3* (*NOD*-, *LRR*- and *pyrin domain-containing protein 3*) is affected by endometrial injury during proliferative phase of menstrual cycle before embryo transfer.

**Materials and Methods:** Twenty women with unexplained repeated implantation failure who failed to conceive during three or more IVF/ICSI cycles and embryo transfer were selected. The patients randomly classified into two study groups (N = 10 in each group). In the intervention group (not in the control group), endometrial scratching was done on day 9-13 in the proliferative phase of the preceding menstrual cycle. Then, endometrial biopsies of the intervention and control groups were performed in the luteal phase (on 19-21 day). The RNA of all samples was extracted and cDNA synthesis was performed. The expression of *NLRP3* was quantified by quantitative real-time PCR.

**Results:** *NLRP3* gene expression from all samples was investigated. Relative expression of *NLRP3* was lower in the intervention samples compared to the controls.

**Conclusion:** The inflammasome components are suggested as a novel family of endometrial biomarkers. This result is in consistent with other studies that showed dysregulated inflammasome activation has involved in the disruption of maternal-fetal immune-tolerance and in pregnancy complications.

**Key words:** Endometrial scratching, *NLRP3*, Unexplained repeated implantation failure.