

## 9<sup>th</sup> Yazd International Congress and Student Award on Reproductive Medicine with 4<sup>th</sup> Congress of Reproductive Genetics

### Poster Presentations

#### P-128

#### Determining an optimal cut-off value for follicle-stimulating hormone to predict microsurgical testicular sperm extraction outcome in patients with non-obstructive azoospermia

Namavar Jahromi B<sup>1</sup>, Zeyghami Sh<sup>2</sup>, Parsanezhad ME<sup>1</sup>, Ghaemmaghami P<sup>3</sup>, Zarei A<sup>1</sup>, Azizi Kutenae M<sup>4</sup>, Sohail P<sup>3</sup>, Keshavarz P<sup>5</sup>.

1. Infertility Research Center, Department of Obstetrics and Gynecology, Shiraz University of Medical Sciences, Shiraz, Iran.

2. Ghadir Madar Hospital, Infertility Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

3. Department of Biostatistics, Medical School, Shiraz University of Medical Sciences, Shiraz, Iran.

4. Ghadir Madar Hospital, Hormozgan Fertility and Infertility Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

5. Department of Radiology, Medical Imaging Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

Email: namavarb@sums.ac.ir

**Background:** Determining the success of sperm retrieval for infertile men before using assisted reproductive technologies can decrease the costs. Using endocrine markers, as an inexpensive and non-invasive method is considered to be a clinically suitable marker in assessment of infertile men.

**Objective:** To determine the optimal cut-off value for follicle stimulating hormone (FSH) to predict the outcome of microsurgical testicular sperm extraction (micro-TESE) in patients with nonobstructive azoospermia (NOA).

**Materials and Methods:** We included a total number

of 180 patients with NOA. The serum level of FSH was determined and all the subjects underwent micro-TESE. We determined the optimal cut-off value for FSH and assessed whether the test could be effectively used as a successful predictor of sperm retrieval by calculating the receiver operating characteristic area under the curve.

**Results:** Overall, we included a total number of 171 patients with mean age of  $34.3 \pm 8.6$  yr. The micro-TESE was considered to be successful in 79 (43.8%) while it failed in 92 (56.2%) patients. We found that the mean level of serum FSH was significantly higher in group those with failed micro-TEST compared to successful group ( $P < 0.001$ ). The cut-off value for FSH was calculated to be 14.6 mIU/mL to predictive the outcome of micro-TESE with a sensitivity of 83.5% [73.5%-90.9%] and a specificity of 80.3% [69.5%-88.5%]. At this value, the other parameters were calculated to be PPV, 81.5%; NPV, 82.4; LR+, 4.23; and LR-, 0.21.

**Conclusion:** The results of the current study indicate that FSH plasma levels above 14.6 mIU/mL can be considered to be the failure predictor of the micro-TESE in NOA patients.

**Key words:** Follicle-stimulating hormone, Non-obstructive azoospermia, Microsurgical testicular sperm extraction.

*The original full text of this abstract has been published in Archives of Endocrinology and Metabolism 2020; 64(2): 165-170. <http://dx.doi.org/10.20945/2359-3997000000217>.*

*How to cite to this article: Jahromi BN, Zeyghami S, Parsanezhad ME, Ghaemmaghami P, Zarei A, Azizi Kutenae M, Sohail P, Keshavarz P. Determining an optimal cut-off value for follicle-stimulating hormone to predict microsurgical testicular sperm extraction outcome in patients with non-obstructive azoospermia. Archives of Endocrinology and Metabolism 2020; 64(2): 165-170.*