

9th Yazd International Congress and Student Award on Reproductive Medicine with 4th Congress of Reproductive Genetics

Poster Presentations

P-139

The histopathological effects of maternal caffeine consumption on testis, prostate, seminal vesicle and epididymis of male rat offspring (stereological study)

Yadegari Dehnavi M^{1, 2}, Belali Kharaji M³, Zare Mehrjerdi F⁴, Pourntezari Zarch M⁵.

Department of Anatomy and Cell Biology, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

Email: yadegarimy@gmail.com

Background: Caffeine is an alkaloid of the methyl xanthine group. It is conceivable that caffeine consumption would induce gonadal changes.

Objective: The aim of this study was to assess the impact of embryonic caffeine exposure on rat testis, prostate, seminal vesicle and epididymis.

Materials and Methods: Female rats were divided into three major groups (n = 7) including: A control which only received drinking water. B and C groups received low dose of caffeine (26 mg/kg) and high dose (45 mg/kg) respectively via drinking water during pregnancy and lactation. Structural changes in testis, prostate, seminal vesicle and epididymis were studied by using stereological methods at 21, 60 and 120 days of postnatal development.

Results: The result showed the decreases in body and testis weight and testis volume of offspring in group C compared to other groups at all ages (p < 0.05). There were significant differences in number of sperm cells

of offspring of experimental groups compared to control group in different ages (p < 0.05). Number of sertoli, spermatogonia, spermatocyte and spermatid cells in group C showed a significant decrease compared with other groups at all ages (p < 0.01). The prostate of caffeine-fed groups presented various degrees of epithelial and stromal hyperplasia and epithelial cell proliferation in the prostatic lobes. In high dose caffeine treated group, epithelium height of seminal vesicle decreased, but it was not significant. The thickness of the muscle layer of seminal vesicles in high dose caffeine treated group increased compared to the control group (p ≤ 0.001). Significant decreases in epithelium height of tubules of epididymis in high dose caffeine treated group was seen compared with control group (p < 0.05). In high dose caffeine treated group diameter of tubules of epididymis decreased (p ≤ 0.001).

Conclusion: Results showed that maternal caffeine consumption altered the structure of testis and prostate gland, seminal vesicle, epididymis, and spermatogenesis of offspring in adulthood.

Key words: Caffeine, Testis, Prostate, Seminal Vesicle, Epididymis.

The original full text of this abstract has been published in *Journal of Rafsanjan University of Medical Sciences* 2020; 19(4): 325-340. <http://eprints.rums.ac.ir/id/eprint/7102>.

How to cite this article: Yadegari Dehnavi M, Belali Kharaji M, Zare Mehrjerdi F, Pourntezari Zarch M. The effects of caffeine consumption on histomorphometric changes of seminal vesicle and epididymis in rat offspring: An experimental study. *Journal of Rafsanjan University of Medical Sciences* 2020; 19(4): 325-340.