

Correlation between follicular fluid 25-OH vitamin D and assisted reproductive outcomes

Laya Farzadi¹ M.D., Homa Khayatzaeh Bidgoli¹ M.D., Morteza Ghojazadeh² M.D., Ph.D., Zahra Bahrami¹ M.D., Amir Fattahi³ M.Sc., Zeinab Latifi³ M.Sc., Vahideh Shahnazi¹ M.Sc., Mohammad Nouri¹ Ph.D.

1. Women's Reproductive Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.
2. Liver and Gastrointestinal Disease Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.
3. Department of Clinical Biochemistry, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran.

Corresponding Author:

Mohammad Nouri, Women's Reproductive Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.
 Email: nourimd@yahoo.com
 Tel: (+98) 9144054268

Received: 12 July 2014

Revised: 16 November 2014

Accepted: 14 February 2015

Abstract

Background: Vitamin D in complex with its receptor by regulating gene expression, endometrium immune response and stimulation of endometrium decidualization can be involved in implantation. So, it seems that the amount of vitamin D in follicular fluids (FF) may have an association with ART success.

Objective: First, we intended to investigate the possible association between levels of follicular fluids 25-OH vitamin D with assisted reproductive outcomes. Second, we examined relationship between 25-OH vitamin D levels with number and quality of oocytes.

Materials and Methods: In a prospective study, 80 infertile female candidates for IVF/ICSI were enrolled. Blood samples (on the day of human chorionic gonadotropin administration) and follicular fluids were taken, and then levels of serum estradiol and follicular fluids 25-OH vitamin D were measured. Also clinical characteristics of patients (duration of infertility, causes of infertility, menstrual status), number and quality of oocytes, number of fertilized oocytes, estradiol levels, and clinical pregnancy were evaluated.

Results: Concentration of FF 25-OH vitamin D in pregnant women was significantly higher than non-pregnant women ($p=0.007$) but there were no significant differences in age, body mass index (BMI), duration of infertility, menstrual status, number of oocytes, oocytes quality, number of fertilized oocytes, and serum estradiol levels between the two groups. Statistically positive correlation was found between 25-OH vitamin D levels with patient age and implantation rate ($r=0.264$, $p=0.018$ and $r=0.301$, $p=0.007$ respectively).

Conclusion: The obtained results suggest that vitamin D without affecting the number and quality of oocytes can independently improve implantation rate and IVF outcome.

Key words: Embryo implantation, Vitamin D, In vitro fertilization, Outcome.

Introduction

Existence of vitamin D receptors in the female reproductive system including the uterus, endometrium, ovaries and placenta, represents pivotal role of vitamin D in female fertility (1). Further studies have suggested that vitamin D deficiency is associated with some female reproductive abnormalities especially gestational diabetes, endometriosis, PCOS, bacterial vaginosis, premature labor pains and preeclampsia (2-7). Also it has been reported that in mice with knocked out vitamin D receptor (VDR) gene, impaired follicle development, gonadal malfunctions, decreased aromatase expression and activity, pregnancy complications, uterine hypoplasia and infertility can be seen (8, 9). This vitamin

in complex with its receptor by regulating of *HOXA10* gene expression, endometrium immune response and stimulation of endometrium decidualization can also be involved in implantation (10).

Considering the importance of vitamin D in pregnancy and implantation and according to Estes *et al* report about decreased expression level of vitamin D binding protein in follicular fluid (FF) as a potential biomarker for ART outcome, it seems that the amount of vitamin D in FF can has a significant association with assisted reproductive technology (ART) success (11). In this regard, Ozkan *et al* have documented that high level of vitamin D in FF and serum is related with chance of conceiving following IVF (12). Bodnar and colleagues also have reported that vitamin D deficiency is associated with reduced

pregnancy outcome (2). Rudick *et al* in a similar study showed a direct correlation between vitamin D and IVF success in non-Asian patients but not in Asian population (13). While, some studies have mentioned the lack of this association or even inverse correlation (14, 15).

Roles of estradiol on uterine growth and folliculogenesis during IVF are absolutely clear, but its effects on implantation and IVF outcome are not well understood. Some researchers have investigated association between serum estradiol level and pregnancy rates following IVF, but the results are contradictory (16-18). It has been documented that serum estradiol levels on the day of human chorionic gonadotropin (hCG) administration have a direct correlation with the number of oocytes and an inverse relationship with implantation and pregnancy rate, while some have emphasized the absence of such a relation (17-19).

Recently, in a study on 1712 patients who underwent their first IVF cycle, it was found that over 10% reduction in estradiol levels after hCG administration leads to a 50% decreasing in clinical pregnancy rate (20). Regarding the role of vitamin D in steroidogenesis and increasing expression and activity of aromatase, it is likely that this vitamin exerts its effects on the number and quality of oocytes and the outcome of pregnancy through increasing estradiol production (21).

Because of mixed results about association of vitamin D and estradiol levels with IVF outcome, in this study we intend to investigate the possible association between levels of serum estradiol and FF 25-OH vitamin D with the number and quality of oocytes and rate of clinical pregnancy.

Materials and methods

In a prospective study, 80 infertile female (age 31 ± 6 years) who were referred to Al Zahra Hospital of Tabriz between February 2009-February 2010 and were candidate for IVF/ICSI were enrolled. Informed consent was obtained from participants according to the Ethical Committee of Tabriz University of Medical Sciences criteria. All of the 25 to 37 years old patients who were in first ART cycle with one of the following causes of infertility were enrolled in the study; male factor

infertility, ovulatory dysfunction, fallopian tube obstruction, and idiopathic. Subjects with history of inflammatory disease, endocrine disorders (hyper- or hypothyroidism, Cushing's syndrome, hyperprolactinemia, etc.), liver, kidney, or heart disease, immune system diseases, endometriosis, consuming drugs interfering with vitamin D metabolism and previous ART cycle experience were excluded.

In all patients, long protocol with administration of exogenous gonadotropin (Gonal-F, Merck, Germany and Menogon[®], Ferring, Germany) was used to stimulate ovulation and when at least 2-3 follicles reached 18 mm, venous blood samples were taken for assessment of serum estradiol and 10,000 IU human chorionic gonadotropin (Choriomon, Lugano, Switzerland) was administered intramuscularly on the same day. Serum was isolated by centrifugation, and for subsequent assay was kept at -70°C . Transvaginal-guided follicular puncture and aspiration of FF was performed under general anesthesia 34 hr after administration of HCG.

FF was centrifuged at 3000g for 15 min and the supernatant was stored at -70°C for 25-OH vitamin D measuring. Then the number and quality of oocytes and the number of fertilized oocytes and also clinical pregnancy rate in each patient were evaluated. For grading oocyte we used method which explained previously by Xia (22). Briefly oocytes were graded into 4 groups; 1) large perivitelline space with fragmented first polar body, 2) large perivitelline space with intact first polar body, 3) normal perivitelline space with fragmented first polar body and 4) normal perivitelline space with intact first polar body.

The number of fertilized oocytes was counted 84 hr after ICSI and depending on the number of embryos available, a maximum of 3 high-quality embryos (based on cleavage and morphology) were transferred to the uterus. Clinical pregnancy was considered when the transvaginal ultrasound revealed intrauterine gestational sac. Implantation rates were obtained from ratio of visible sacs number to transferred embryos number. In all samples serum estradiol level was measured by commercial ELISA kit (Roche, Germany) and 25-OH vitamin D level in FF was evaluated by enzyme immunoassay method using DIAsource ImmunoAssays[®] kit (Belgium).

Statistical analysis

Data are provided as mean±SD and normal distribution of data was examined by Kolmogorov-Smirnov test. To compare quantitative data between groups, Mann-Whitney U test was used and Chi-square test was employed for qualitative data. Logistic regression was used to determine the regression model. Spearman test was used to evaluate correlation between the study parameters. Data analysis was performed using SPSS V.16 software and statistical significance was assumed at the p≤0.05 level.

Results

Of 80 patients, in 18 (22.5%) subjects clinical pregnancies were observed, but in 62 (77.5%) cases the pregnancy did not occurred. Basal characteristics of pregnant and non-pregnant patients are shown in table I. Statistical analysis showed no significant differences in age, body mass index (BMI), duration of infertility, menstrual status, between the two groups. As shown in table II, we did not find significant differences in the

number of oocytes, oocytes quality, number of fertilized oocytes and serum estradiol levels (day of hCG administration) between pregnant and non-pregnant women. Only a significant difference was found in concentration of FF 25-OH vitamin D between the two groups which the concentration was higher in the group with clinical pregnancy. Evaluating the ratio of appeared sacs number to transferred embryos number in pregnant women demonstrated that the average rate of implantation is 52.3±22.4.

In the assessment of correlation between variables, no significant correlation was observed between FF 25-OH vitamin D levels with BMI, duration of infertility, number of oocytes, number of fertilized oocytes and estradiol levels. Statistically positive correlation was found between 25-OH vitamin D levels with patient age and implantation rate (Table III). Logistic regression analysis revealed that after controlling for the effects of age, for each ng/ml increasing of 25-OH vitamin D level in follicular fluid, the rate of clinical pregnancy following IVF increases 1.2 times.

Table I. Basal characteristics of clinical pregnant and non-pregnant patients

Parameters	Clinical pregnant (n=18)	Non-Clinical pregnant (n=62)	p-value
Age (year) ^a	31.9 ± 4.9	29.9 ± 4.8	0.052
BMI (kg/m ²) ^a	27.2 ± 3.8	27.6 ± 3.8	0.954
Duration of infertility (year) ^a	7.3 ± 4	6.5 ± 3.6	0.513
Causes of infertility ^{a,b}			
Male factor (n=46)	11 (61.1%)	35 (56.5%)	
Ovulatory dysfunction (n=13)	2 (11.1%)	11 (17.7%)	
Fallopian tube obstruction (n=8)	1 (5.6%)	7 (11.3%)	0.70
Idiopathic (n=13)	4 (22.2%)	9 (14.5%)	
Menstrual status ^{a,b}			
Regular	16 (88.9%)	53 (85.5%)	
Irregular	2 (11.1%)	9 (14.5%)	0.529

BMI: Body mass index.
a: Mann-Whitney U

* Data are presented as mean±SD.
b: Chi-square test

** Data are presented as n (%)

Table II. Cycle characteristics of clinical pregnant and non-pregnant patients

Parameters	Clinical pregnant (n=18)	Non-Clinical pregnant (n=62)	p-value
Number of oocytes ^a	12.2 ± 4.6	11.4 ± 7.1	0.352
Oocytes quality ^{a,b}			
G1	0 (0%)	0 (0%)	
G2	78 (35.45%)	233 (33.28%)	
G3	128 (58.18%)	350 (50%)	0.395
G4	14 (6.37%)	117 (16.72%)	
Number of fertilized oocytes ^a	6.2 ± 3.3	5.3 ± 4.7	0.120
Estradiol (pg/ml) ^a	2751.4 ± 1224.6	2546.6 ± 1464	0.595
25-OH vitamin D (ng/ml) ^a	15.8 ± 4.5	11.6 ± 5.5	0.007

* Data are presented as mean±SD.

** Data are presented as n (%)

a: Mann-Whitney U test

b: Chi-square test

G1: large perivitelline space with fragmented first polar body.

G2: large perivitelline space with intact first polar body

G3: normal perivitelline space with fragmented first polar body and

G4: normal perivitelline space with intact first polar body.

Table III. Correlation between 25-OH vitamin D levels with other study variables

Variable	rho	p-value
Age	0.264	0.018
BMI	0.054	0.633
Duration of infertility	0.002	0.988
Number of oocytes	0.198	0.078
Number of fertilized oocytes	0.163	0.149
Estradiol levels	0.067	0.560
Implantation rates	0.301	0.007
Oocytes quality		
G1 ^a	-	-
G2	0.045	0.605
G3	0.068	0.520
G4	0.024	0.833

^a Oocyte with G1 quality did not observed

BMI: Body mass index

Spearman test was used to evaluate correlation between 25-OH vitamin D levels with the factors and correlation coefficient (rho) with p-value is presented for each variable.

Discussion

Several studies have investigated association between IVF outcome and different factors in follicular fluid such as anti-mullerian hormone, pregnancy-associated plasma protein-A, cholesteryl ester transfer protein and vitamin D (12, 23-25). Studies have offered that vitamin D could be involved in important functions of female reproductive system such as steroidogenesis, folliculogenesis, implantation and developing fetus (8, 21, 26, 27). It was recently reported that this vitamin could possibly be a prognosis factor for pregnancy success following IVF, although the results are contradictory and the mechanisms of impact on IVF results has not been exactly determined (12).

In this study, we have examined the association between 25-OH vitamin D of FF with the number and quality of oocytes, number of fertilized oocytes and also implantation and clinical pregnancy rates. No significant differences between pregnant and non-pregnant groups were found in age, BMI, duration of infertility, menstrual status, number of oocytes, oocytes quality, number of fertilized oocytes and serum estradiol levels (day of hCG administration), whereas FF levels of 25-OH vitamin D in pregnant group was significantly higher than non-pregnant subjects.

Our results also revealed a positive correlation between 25-OH vitamin D levels with implantation rates and age, while such a relationship were not found with BMI, duration of infertility, number of oocytes, number of fertilized oocytes and serum estradiol levels. The obtained results suggest that vitamin D without affecting the number and quality of oocytes can independently improve

implantation. In similar studies, Anifandis, Ozkan and Rudick have observed a direct link between vitamin D levels and IVF outcome (12, 13, 15). Studies have shown that vitamin D deficiency or defect in its receptor can reduce chances of pregnancy by disrupting follicles and oocytes development and gonads function especially in estradiol synthesis, so it has been proposed that maybe high levels of vitamin D through affecting these items can improve pregnancy outcome (6, 8, 9, 12).

However, our study demonstrated that vitamin D without having effect on the number and quality of oocytes, basal estradiol levels and number of fertilized oocytes can increase chance of pregnancy by 1.2-fold. In some of previous studies association between vitamin D level with number and quality of oocytes and embryos has not been found (14, 28). This indicates that potential impact of vitamin D on implantation rate is probably by binding to its receptors in the endometrium. Ozkan *et al* have also suggested that vitamin D may increase endometrial receptivity and chance of pregnancy via binding to its receptor in the endometrium (12). One of the possible mechanisms of vitamin D-receptor complex can be increasing the expression of *HOXA10* gene which has an essential role in implantation (10). Moreover, it is likely that vitamin D may play a role in uterine receptivity by modulating endometrial immune response, as it has been seen that this vitamin can reduce the decidual T-cell activity and synthesis of cytokines, interleukin 1 and 6 and TNF α which are essential in embryo implantation and uterine receptivity (27, 29).

On the other hand some researchers have documented that there is an inverse correlation between vitamin D levels with the number and quality of follicles and oocytes

and also IVF success, although our results did not show such a correlation (15). Anifandis *et al* have reported that vitamin D decreases chance of pregnancy through affecting insulin function and decreasing FF glucose which is essential for oocyte and granulosa development (15). Some studies also have emphasized the lack of association between vitamin D levels with the quality and number of oocytes and pregnancy rate (30).

These conflicting results are probably due to the low sample size, different causes of infertility in women, lack of identical protocol in ovulation induction and eclipse of pregnancy achieving by various factors. The advantages of our study are using similar protocol of treatment for patients, lack of differences in age, BMI, menstrual status, causes of infertility, number and quality of oocytes, number of fertilized oocyte and basal estradiol level between pregnant and non-pregnant patients which cause revealing the effects of vitamin D on implantation and pregnancy success. However the sample size in this study was relatively low, and for obtaining a clear conclusion and understanding underlying mechanisms more investigations with high sample size are needed.

Conclusion

In conclusion we have found that vitamin D without affecting the number and quality of oocytes can independently improve implantation rate and IVF outcome.

Acknowledgments

We are indebted to Al Zahra Hospital of Tabriz for providing the samples. This study was financially supported by Tabriz University of Medical Sciences.

Conflict of interest

The authors have no conflicts of interest to disclose.

References

1. Shahbazi M, Jeddi-Tehrani M, Zareie M, Salek-Moghaddam A, Akhondi M, Bahmanpoor M, et al. Expression profiling of vitamin D receptor in placenta, decidua and ovary of pregnant mice. *Placenta* 2011; 32: 657-664.

2. Kramer CK, Swaminathan B, Hanley AJ, Connelly PW, Sermer M, Zinman B, et al. Vitamin D and Parathyroid Hormone Status in Pregnancy: Impact on Insulin Sensitivity, Beta-cell Function, and Gestational Diabetes Mellitus. *J Clin Endocrinol Metab* 2014; 99: 4506-4513.
3. Zhang C, Qiu C, Hu FB, David RM, van Dam RM, Bralley A, et al. Maternal plasma 25-hydroxyvitamin D concentrations and the risk for gestational diabetes mellitus. *PLoS One* 2008; 3: e3753.
4. Abdi J, Redinger D, Broughton KS. Plasma vitamin D level among women with and without polycystic ovarian syndrome. *FASEB J* 2013; 27: 635.
5. Mahmoudi T, Gourabi H, Ashrafi M, Yazdi RS, Ezabadi Z. Calcitropic hormones, insulin resistance, and the polycystic ovary syndrome. *Fertil Steril* 2010; 93: 1208-1214.
6. Bodnar LM, Krohn MA, Simhan HN. Maternal vitamin D deficiency is associated with bacterial vaginosis in the first trimester of pregnancy. *J Nutr* 2009; 139: 1157-1161.
7. Abedi P, Mohaghegh Z, Afshary P, Latifi M. The relationship of serum vitamin D with pre-eclampsia in the Iranian women. *Matern Child Nutr* 2014; 10: 206-212.
8. Yoshizawa T, Handa Y, Uematsu Y, Takeda S, Sekine K, Yoshihara Y, et al. Mice lacking the vitamin D receptor exhibit impaired bone formation, uterine hypoplasia and growth retardation after weaning. *Nat Genet* 1997; 16: 391-396.
9. Kinuta K, Tanaka H, Moriwake T, Aya K, Kato S, Seino Y. Vitamin D Is an Important Factor in Estrogen Biosynthesis of Both Female and Male Gonads 1. *Endocrinology* 2000; 141: 1317-1324.
10. Lerchbaum E, Obermayer-Pietsch B. Mechanisms in endocrinology: Vitamin D and fertility: a systematic review. *Eur J Endocrinol* 2012; 166: 765-778.
11. Estes SJ, Ye B, Qiu W, Cramer D, Hornstein MD, Missmer SA. A proteomic analysis of IVF follicular fluid in women ≤ 32 years old. *Fertil Steril* 2009; 92: 1569-1578.
12. Ozkan S, Jindal S, Greenseid K, Shu J, Zeitlian G, Hickmon C, et al. Replete vitamin D stores predict reproductive success following in vitro fertilization. *Fertil Steril* 2010; 94: 1314-1319.
13. Rudick B, Ingles S, Stanczyk F, Chung K, Paulson R, Bendikson K. Characterizing the role of vitamin D levels on IVF outcomes: stimulation, embryo, or endometrium? *Fertil Steril* 2010; 94: S72.
14. Aleyasin A, Hosseini MA, Mahdavi A, Safdarian L, Fallahi P, Mohajeri MR, et al. Predictive value of the level of vitamin D in follicular fluid on the outcome of assisted reproductive technology. *Eur J Obstet Gynecol Reprod Biol* 2011; 159: 132-137.
15. Anifandis GM, Dafopoulos K, Messini CI, Chalvatzas N, Liakos N, Pournaras S, et al. Prognostic value of follicular fluid 25-OH vitamin D and glucose levels in the IVF outcome. *Reprod Biol Endocrinol* 2010; 8: 91.
16. Joo BS, Park SH, An BM, Kim KS, Moon SE, Moon HS. Serum estradiol levels during controlled ovarian hyperstimulation influence the pregnancy outcome of in vitro fertilization in a concentration-dependent manner. *Fertil Steril* 2010; 93: 442-446.
17. Imudia AN, Goldman RH, Awonuga AO, Wright DL, Styer AK, Toth TL. The impact of supraphysiologic serum estradiol levels on peri-implantation embryo

- development and early pregnancy outcome following in vitro fertilization cycles. *J Assist Reprod Genet* 2014; 31: 65-71.
18. Peña JE, Chang PL, Chan L-K, Zeitoun K, Thornton MH, Sauer MV. Supraphysiological estradiol levels do not affect oocyte and embryo quality in oocyte donation cycles. *Hum Reprod* 2002; 17: 83-87.
 19. Ozdegirmenci O, Dilbaz S, Cinar O, Aydin S, Beydilli G, Cakir L, et al. Can serum oestradiol be a predictor of quality of oocytes and embryos, maturation of oocytes and pregnancy rate in ICSI cycles? *Gynecol Endocrinol* 2011; 27: 279-285.
 20. Kondapalli L, Molinaro T, Sammel M, Dokras A. A decrease in serum estradiol levels after human chorionic gonadotrophin administration predicts significantly lower clinical pregnancy and live birth rates in in vitro fertilization cycles. *Hum Reprod* 2012; 27: 2690-2697.
 21. Parikh G, Varadinova M, Suwandhi P, Araki T, Rosenwaks Z, Poretsky L, et al. Vitamin D regulates steroidogenesis and insulin-like growth factor binding protein-1 (IGFBP-1) production in human ovarian cells. *Horm Metab Res* 2010; 42: 754.
 22. Xia P. Intracytoplasmic sperm injection: correlation of oocyte grade based on polar body, perivitelline space and cytoplasmic inclusions with fertilization rate and embryo quality. *Hum Reprod* 1997; 12: 1750-1755.
 23. Aflatoonian A, Mashayekhy M, Mohamadian F, Moghaddam FM. The correlation between follicular fluid anti-mullerian hormone levels and fertilization and embryo quality in ART cycles. *Iran J Reprod Med* 2010; 8: 157-160.
 24. Dehghani Firouzabadi R, Mohammadian F, Mashayekhy M, Davar R, Eftekhar M. The correlation between follicular fluid pregnancy-associated plasma protein A levels, fertilization, and embryo quality in GnRH agonist and GnRH antagonist protocols in ART cycles. *Iran J Reprod Med* 2012; 10: 477-482.
 25. Mehdizadeh A, Rahimipour A, Farzadi L, Darabi M, Shahnazi V, Shaaker M, et al. Correlation between the level of cholesteryl ester transfer protein in follicular fluid with fertilization rates in IVF/ICSI cycles. *Iran J Reprod Med* 2011; 9: 193-198.
 26. Liu N, Hewison M. Vitamin D, the placenta and pregnancy. *Arch Biochem Biophys* 2012; 523: 37-47.
 27. Pasco JA, Wark JD, Carlin JB, Ponsonby A-L, Vuillermin PJ, Morley R. Maternal vitamin D in pregnancy may influence not only offspring bone mass but other aspects of musculoskeletal health and adiposity. *Med Hypothes* 2008; 71: 266-269.
 28. Rudick B, Ingles S, Stanczyk F, Chung K, Paulson R, Bendikson K. The role of vitamin D levels on IVF outcomes in donor-recipient cycles. *Fertil Steril* 2011; 95: S8.
 29. Vignani P, Lattuada D, Mangioni S, Ermellino L, Vignali M, Caporizzo E, et al. Cycling and early pregnant endometrium as a site of regulated expression of the vitamin D system. *J Mol Endocrinol* 2006; 36: 415-424.
 30. Aflatoonian A, Modarresi S, Sekhavat L, MohammadTaheri S. Therapeutic effects of calcium & vitamin D supplementation in women with PCOS. *Comp Ther Clin Pract* 2012; 18: 85-88.