

Effect of quercetin on the number of blastomeres, zona pellucida thickness, and hatching rate of mouse embryos exposed to actinomycin D: An experimental study

Hamid Reza Sameni Ph.D., Sara Sadat Javadinia M.Sc., Manouchehr Safari Ph.D., Mohammad Hasan Tabrizi Amjad M.Sc., Nasrin Khanmohammadi M.Sc., Houman Parsaie M.Sc., Sam Zarbakhsh Ph.D.

Research Center of Nervous System Stem Cells, Department of Anatomy, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran.

Corresponding Author:
Sam Zarbakhsh, Research Center of Nervous System Stem Cells, Department of Anatomy, Faculty of Medicine, Semnan University of Medical Sciences, Semnan, Iran. P.O.Box: 35198-99951.
Email: smzarbakhsh@gmail.com
Tel: (+98) 2333654218

Received: 9 January 2017
Revised: 5 July 2017
Accepted: 15 November 2017

Abstract

Background: Quercetin is a flavonoid with the ability to improve the growth of embryos in vitro, and actinomycin D is an inducer of apoptosis in embryonic cells.

Objective: The aim was to evaluate the effect of quercetin on the number of viable and apoptotic cells, the zona pellucida (ZP) thickness and the hatching rate of preimplantation embryos exposed to actinomycin D in mice.

Materials and Methods: Two-cell embryos were randomly divided into four groups (Control, Quercetin, actinomycin D, and Quercetin + actinomycin D group). Blastocysts percentage, hatched blastocysts, and ZP thickness of blastocysts was measured. The number of blastomeres was counted by Hoechst and propidium iodide staining and the apoptotic cells number was counted by TUNEL assay.

Results: The results showed that the use of quercetin significantly improved the growth of embryos compared to the control group ($p=0.037$). Moreover, quercetin reduced the destructive effects of actinomycin D on the growth of embryos significantly ($p=0.026$).

Conclusion: quercetin may protect the embryos against actinomycin D so that increases the number of viable cells and decreases the number of apoptotic cells, which can help the expansion of the blastocysts, thinning of the ZP thickness and increasing the hatching rate in mouse embryos.

Key words: *Quercetin, Embryonic development, Zona pellucida, Apoptosis, Blastocyst inner cell mass.*

This article extracted from M.Sc. thesis. (Sara Sadat Javadinia)

Introduction

In vitro production of the embryo is an important method for improving reproductive technology. The development potential of embryos produced by in vitro production is related to the composition of the culture medium (1). It is known that embryo development in culture media can be damaged by several stressors such as high oxygen concentration and high level of reactive oxygen species (ROS) (2). It is important to keep embryos in culture media from oxidative stress and for this purpose, antioxidants are valuable candidates (3).

Flavonoids are phyto phenolic compounds with the strong antioxidant effect that work as free radical scavengers. Quercetin is an important member of flavonoid family in the

human diet which is abundant in fruits and vegetables such as, apple, onion, strawberry, tea, green tea and broccoli (4, 5). It has been reported that quercetin has anti-oxidative, anti-inflammatory and anti-mutagenic activities due to free radical scavenging so that quercetin reduces anomalies in rat embryos, decreases neural tube defects in embryos of diabetic mice and reduces hair cell damage in zebrafish embryos (5-8).

Actinomycin D is a DNA ligand drug that clinically used as an antibiotic to treat cancer. Actinomycin D inhibits DNA transcription by connecting to guanine-cytosine base pairs (9). In some reports, actinomycin D has been used to induce apoptosis in preimplantation embryos (10, 11). By studying the embryonic morphology, predicting the fate of the embryos is largely possible. The most

morphological indicators to select the best embryos for transfer are zona pellucida (ZP) thickness and the number of embryonic cells (12). ZP thickness is a reliable indicator of in vitro fertilization success rate that can be applied as a criterion for embryo selection. Actually, ZP thickness is inversely correlated with embryonic viability and hatching rate (13, 14).

Moreover, cleavage rate and development to blastocyst are applied as a quality parameter of mammal embryos (12, 15). In the previous study, we showed l-carnitine as an antioxidant could increase embryo development and blastocyst quality but it is still unclear which antioxidant is more effective in embryo culture and so different antioxidants should be investigated (16). Recent studies have reported quercetin has the protective effect on the development of mouse embryos against H₂O₂ and hydroxyurea (17, 18).

The aim of this study was to evaluate the antioxidant and anti-apoptotic effects of quercetin on some morphological indicators of preimplantation embryos in vitro. The morphological indicators include the ZP thickness, the number of viable and apoptotic blastomeres and the hatching rate. To evaluate the antioxidant and anti-apoptotic properties of quercetin, actinomycin D was used as an inducer of apoptosis in blastomeres.

Materials and methods

Animals

In this experimental study, female C57BL/6 mice (6-8 wk) were kept under controlled temperature (25±2°C) and controlled light (12 hr light/dark), with free access to food and water.

Superovulation and embryo collection

The mice were superovulated by an intraperitoneal injection of 10 IU pregnant mare's serum gonadotropin (Gonaser 5.000 UI, Spain) followed 48 hr later by an intraperitoneal injection of 10 IU human chorionic gonadotropin (hCG) (Sigma, China) (19). Then they were mated overnight with males and the mating was emphasized by the presence of vaginal plug on the morning after hCG injection. Two-cell embryos were flushed

from the oviduct at about 48 hr after hCG injection and washed in human tubal fluid (HTF) medium containing HEPES (Sigma, USA). From 20 female mice, a total of 240 two-cell embryos were obtained and used in this study.

Embryo culture

The 240 two-cell embryos were transferred into the HTF medium (supplemented with 10% human serum albumin) (Sigma, USA). Two-cell embryos were randomly divided into four groups (60 embryos in each group): I. Control group, without any treatment, II. Quercetin (Sigma, China) group, 5 µM of quercetin (17) was added into the medium, III. Actinomycin D (Sigma, USA) group, 0.005 µg/ml of actinomycin D (10) was added into the medium, IV. Quercetin + actinomycin D group, 5 µM of quercetin and 0.005 µg/ml of actinomycin D were added together into the medium.

In all groups, 10 embryos were situated in a drop of 20 µl of HTF medium under mineral oil (Sigma, USA) in a 35 mm Petri dish (Jet Biofil, Canada) and were incubated at 37°C with 95% humidity and 5% CO₂. To evaluate the anti-apoptotic effect of quercetin, when the two-cell embryos reached the eight-cell stage, they were incubated with 0.005 µg/ml of actinomycin D in the medium for 4 hr. Actinomycin D concentration was derived from the other report (10). In 120 hr after the onset of incubation, the percentage of embryos that reached to blastocyst and hatched blastocyst stages was assessed.

Measurement of ZP thickness

The ZP thickness measurement was performed in two stages of the early blastocyst and full blastocyst. First, blastocysts were randomly selected. Then the thickness of each ZP was measured at three steady points and the average was calculated (12, 16). The measurement was taken from images using an inverted microscope (Nikon, Eclipse Ti-U, Japan) and motic images plus 2.0 software.

Differential staining and detection of apoptotic nuclei

When the blastocysts were expanded, randomly selected for counting the number of

total blastomeres by differential staining and TUNEL assay. Differential staining to separate the inner cell mass and trophoctoderm cells and TUNEL assay to detect the apoptotic nuclei were done according to the method described by Fouladi-Nashta and colleagues (20, 21). Briefly, the blastocysts were treated with 30 µg/ml propidium iodide (PI) (Sigma, China) and 1% Triton X-100 (Sigma, China) at 37°C for 5 min. Immediately after, the blastocysts were washed twice and fixed in 4% paraformaldehyde containing 10 µg/ml bisbenzimidazole (Hoechst 33342) (Sigma, USA) for 20 min at room temperature that allows fixation of blastocysts and staining total cell nuclei.

Next, the blastocysts were washed and incubated in droplets of in situ cell death detection (TUNEL) kit solution (Roche, Germany) for 45 min according to the manufacturer's instructions. Then the blastocysts were mounted on glass slides in glycerol droplets and were observed under an inverted fluorescent microscope (Motic, AE31, Spain). Inner cell mass (ICM) nuclei labeled with Hoechst appeared blue, trophoctoderm (TE) nuclei labeled with both Hoechst and PI appeared red and apoptotic cells labeled with TUNEL appeared green. The number of ICM, TE and apoptotic cells was counted.

Ethical consideration

All animal protocols were approved by the Research Council of Semnan University of Medical Sciences (IR.SEMUMS.REC.1394.83).

Statistical analysis

Comparison of the percentage of embryos from two-cell to hatched blastocyst was analyzed by χ^2 test. The ZP thickness, the blastocyst cell count and the TUNEL positive cell count were analyzed by independent t-test as mean \pm standard deviation. A difference with $p < 0.05$ was considered statistically significant.

Results

Developmental rate of embryos

There was no statistically significant difference in the percentage of two-cell embryos to morula stage between quercetin and control groups ($p=0.559$); and between quercetin + actinomycin D and actinomycin D groups ($p=0.690$) (Table I). The percentage of embryos that reached to blastocyst and hatched blastocyst stages in quercetin group was significantly higher than the control group ($p=0.037$), and in quercetin+actinomycin D group was significantly higher than actinomycin D group ($p=0.026$) (Figure 1A, B).

The ZP thickness of blastocysts

The results of ZP thickness of blastocysts showed the ZP thickness in quercetin group was significantly thinner than the control group ($p=0.043$), and in quercetin + actinomycin D group was significantly thinner than actinomycin D group ($p=0.027$) (Figure 1A, C).

Blastocyst cell count

Expanded blastocysts were stained with Hoechst and PI and the number of ICM and TE was counted. The results showed the number of ICM and TE in quercetin group was significantly higher than the control group ($p=0.034$), and in quercetin+ actinomycin D group was significantly higher than actinomycin D group ($p=0.037$) (Figure 2A, B).

Anti-apoptotic effect of quercetin

The TUNEL assay was used to determine the anti-apoptotic effect of quercetin on blastocyst development in the presence or absence of actinomycin D. The results showed the number of TUNEL-positive nuclei in quercetin group was significantly lower than control group ($p=0.011$), and in quercetin+actinomycin D group was significantly lower than actinomycin D group ($p=0.019$) (Figure 2A, C).

Table I. two-cell embryos up to morula stage in different groups

Groups	2-Cell	4-Cell	8-Cell	Morula
Control	60 (100)	60 (100)	59 (98.33)	58 (96.66)
Quercetin	60 (100)	60 (100)	60 (100)	59 (98.33)
Actinomycin D	60 (100)	59 (98.33)	58 (96.66)	41 (68.33)
Quercetin+Actinomycin D	60 (100)	60 (100)	58 (96.66)	43 (71.66)

Data presented as n (%)





Figure 1. A. The blastocysts of I) Control, II) Quercetin, III) Actinomycin D, and IV) Quercetin + actinomycin D (×200). B. The results of the percentage of blastocysts and hatched blastocysts. C. The results of zona pellucida thickness of early and expanded blastocysts. *p<0.05 vs. the control group. #p<0.05 vs. the actinomycin D group.






Figure 2. A. Differential staining and TUNEL labeling of blastocysts. The triple staining with propidium iodide (PI) for trophoblast cells (red), Hoechst for total cells (blue), and TUNEL for apoptotic cells (green) ($\times 200$). B. The results of blastocyst cell numbers with PI and Hoechst staining. C. The results of apoptotic cells of blastocysts with TUNEL assay.

* $p < 0.05$ vs. the control group.

$p < 0.05$ vs. the actinomycin D group.

Discussion

In this study, we evaluated the effect of quercetin on some morphological indicators of mouse embryo development and blastocyst quality including the ZP thickness, the hatching rate, the number of blastomeres and the rate of apoptosis induced by actinomycin D. Overall, the results showed 5 μM of quercetin was able to promote the development of embryos exposed to actinomycin D and reduced apoptosis in the blastomeres, suggesting quercetin as an antioxidant had a protective effect on mouse embryo development and blastocyst quality against actinomycin D.

Quercetin is one of the most important flavonoids. The antioxidant capacity of quercetin has been shown in different cell types (22, 23). In relation to the effect of quercetin on mouse embryo development, Yu *et al* have reported that quercetin has the protective effect on mouse embryos against H_2O_2 (17). In the other study, Perez-Pasten and co-worker have reported that using more than 100 μM of quercetin causes some defects and anomalies, and 3 μM of quercetin has the protective effect on mouse embryos against hydroxyurea (18). Moreover, Shahzad *et al* have reported that administration of 50 mg/kg/day to pregnant rats decreased implantation rate (24). It seems that high doses of antioxidants may act as enzyme inhibitors and mutagens due to inhibition of

topoisomerases, proteasome synthesis or fatty-acid synthesis (25, 26). To induce apoptosis, we used actinomycin D which is known as an inducer of apoptosis on different cell types by inhibiting DNA transcription (11, 27). Some articles have shown that actinomycin D decreases development and cleavage rate of embryos, and increases apoptotic cells in blastocyst (10, 11, 28).

The results showed the quercetin improved development of two-cell embryos to hatched blastocyst stage and reduced apoptotic cells of the embryos exposed to actinomycin D. These results are in accordance with the other reports that have shown quercetin has protective effects on mouse embryos against H_2O_2 and hydroxyurea (17, 18). Due to the different mechanism of actinomycin D with H_2O_2 and hydroxyurea, our results confirmed quercetin could protect mouse embryos in vitro from various harmful factors. These effects of quercetin are probably related to its antioxidant action in reducing ROS level because it is well known that ROS induces apoptotic cell death in different cell types (29, 30). Moreover, quercetin protects the mitochondrial function (31) and regulates enzyme antioxidant defense systems (32). In the other study, Sovernigo *et al* have reported that quercetin can improve bovine blastocyst development (33). The results of our study are in agreement with Fan and Sovernigo reports.

By studying morphological indicators, embryo quality is checked and the best

embryos are selected for transfer (34-36). From the most important indicators of embryonic morphology are the number of viable and apoptotic cells of the blastocyst, the ZP thickness of blastocyst and the ability of hatching (37, 38). The number of blastocyst cells is important for appropriate implantation and various studies have reported that reduction in the number of blastocyst cells leads to reduced embryonic viability (39, 40). Blastocoel fluid has H₂O₂ that is cytotoxic and induces apoptosis in blastocyst cells (41). Antioxidant effect of quercetin probably increases intracellular glutathione level and because glutathione is involved in the removal of H₂O₂, blastocyst cells would not become apoptosis (6). Our results showed quercetin increased the number of viable cells of the blastocyst and decreased apoptotic cells that these results were in agreement with the other report (17).

ZP thickness is a marker to select the best frozen-thawed embryos for transfer (42) because the thinner ZP increases the probability of the hatching rate and implantation. ZP thickness depends on inherent features of embryos to generate the lytic factors needed for ZP thinning (13, 43). Khanmohammadi and colleagues showed L-carnitine could cause thinning of the ZP thickness of blastocysts (16). Prior to the present study, the effect of quercetin on ZP thickness has not been investigated. We showed that quercetin could cause thinning of the ZP thickness of blastocysts. It seems that blastocyst expansion depends on the number of viable cells of the blastocyst and an increase in these cells leads to thinning of the ZP thickness and hatching of the blastocyst (16, 44). More research is required to clarify the molecular mechanisms underlying quercetin function on development and quality of embryos in different conditions.

Conclusion

The results suggest that quercetin may protect the embryos against actinomycin D so that increases the number of viable cells and decreases the number of apoptotic cells, which can help the expansion of the blastocysts, thinning of the ZP thickness and

increasing the hatching rate in mouse embryos.

Acknowledgments

This study was supported by a thesis grant number 880 from Semnan University of Medical Sciences. We would like to thank the Research Center of Nervous System Stem Cells of Semnan University of Medical Sciences for cooperation and providing facilities to this work.

Conflict of interest

There is no conflict of interest in this article.

References

1. Paternot G, Debrock S, D'Hooghe TM, Spiessens C. Early embryo development in a sequential versus single medium: a randomized study. *Reprod Biol Endocrinol* 2010; 8: 83.
2. Li XX, Lee KB, Lee JH, Kim KJ, Kim EY, Han KW, et al. Glutathione and cysteine enhance porcine preimplantation embryo development in vitro after intracytoplasmic sperm injection. *Theriogenology* 2014; 81: 309-314.
3. Truong TT, Soh YM, Gardner DK. Antioxidants improve mouse preimplantation embryo development and viability. *Hum Reprod* 2016; 31: 1445-1454.
4. Perez A, Gonzalez-Manzano S, Jimenez R, Perez-Abud R, Haro JM, Osuna A, et al. The flavonoid quercetin induces acute vasodilator effects in healthy volunteers: correlation with beta-glucuronidase activity. *Pharmacol Res* 2014; 89: 11-18.
5. Lee SK, Oh KH, Chung AY, Park HC, Lee SH, Kwon SY, et al. Protective role of quercetin against cisplatin-induced hair cell damage in zebrafish embryos. *Hum Exp Toxicol* 2015; 34: 1043-1052.
6. Kang JT, Moon JH, Choi JY, Park SJ, Kim SJ, Saadeldin IM, et al. Effect of antioxidant flavonoids (quercetin and taxifolin) on in vitro maturation of porcine oocytes. *Asian-Australas J Anim Sci* 2016; 29: 352-358.
7. Sistani Karampour N, Arzi A, Najafzadeh Varzi H, Mohammadian B, Rezaei M. Quercetin preventive effects on theophylline-induced anomalies in rat embryo. *Jundishapur J Nat Pharm Prod* 2014; 9: e17834.
8. Cao L, Tan C, Meng F, Liu P, Reece EA, Zhao Z. Amelioration of intracellular stress and reduction of neural tube defects in embryos of diabetic mice by phytochemical quercetin. *Sci Rep* 2016; 6: 21491.
9. Cesconetto EC, Junior FS, Crisafuli FA, Mesquita ON, Ramos EB, Rocha MS. DNA interaction with Actinomycin D: mechanical measurements reveal the details of the binding data. *Phys Chem Chem Phys* 2013; 15: 11070-11077.
10. Abdelrazik H, Sharma R, Mahfouz R, Agarwal A. L-carnitine decreases DNA damage and improves the

- in vitro blastocyst development rate in mouse embryos. *Fertil Steril* 2009; 91: 589-596.
11. Niknafs B, Mehdi-pour A, Mohammadi Roushandeh A. Melatonin improves development of early mouse embryos impaired by actinomycin-D and TNF- α . *Iran J Reprod Med* 2014; 12: 799-804.
 12. Marco-Jiménez F, Naturil-Alfonso C, Jiménez-Trigos E, Lavara R, Vicente JS. Influence of zona pellucida thickness on fertilization, embryo implantation and birth. *Anim Reprod Sci* 2012; 132: 96-100.
 13. Filho ES, Noble JA, Wells D. A review on automatic analysis of human embryo microscope images. *Open Biomed Eng J* 2010; 4: 170-177.
 14. Ghannadi A, Kazerooni M, Jamalzadeh F, Amiri S, Rostami P, Absalan F. The effects of laser assisted hatching on pregnancy rates. *Iran J Reprod Med* 2011; 9: 95-98.
 15. Lundin K, Bergh C, Hardarson T. Early embryo cleavage is a strong indicator of embryo quality in human IVF. *Hum Reprod* 2001; 16: 2652-2657.
 16. Khanmohammadi N, Movahedin M, Safari M, Sameni HR, Yousefi B, Jafari B, et al. Effect of L-carnitine on in vitro developmental rate, the zona pellucida and hatching of blastocysts and their cell numbers in mouse embryos. *Int J Reprod Biomed* 2016; 14: 649-656.
 17. Yu S, Long H, Lyu QF, Zhang QH, Yan ZG, Liang HX, et al. Protective effect of quercetin on the development of preimplantation mouse embryos against hydrogen peroxide-induced oxidative injury. *PLoS One* 2014; 9: e89520.
 18. Pérez-Pastén R, Martínez-Galero E, Chamorro-Cevallos G. Quercetin and naringenin reduce abnormal development of mouse embryos produced by hydroxyurea. *J Pharm Pharmacol* 2010; 62: 1003-1009.
 19. Dai SJ, Xu CL, Wang J, Sun YP, Chian RC. Effect of culture medium volume and embryo density on early mouse embryonic development: tracking the development of the individual embryo. *J Assist Reprod Genet* 2012; 29: 617-623.
 20. Fouladi-Nashta AA, Alberio R, Kafi M, Nicholas B, Campbell KH, Webb R. Differential staining combined with TUNEL labelling to detect apoptosis in preimplantation bovine embryos. *Reprod Biomed Online* 2005; 10: 497-502.
 21. Fouladi-Nashta AA, Wonnacott KE, Gutierrez CG, Gong JG, Sinclair KD, Garnsworthy PC, et al. Oocyte quality in lactating dairy cows fed on high levels of n-3 and n-6 fatty acids. *Reproduction* 2009; 138: 771-781.
 22. Barreca D, Bellocco E, D'Onofrio G, Nabavi SF, Daglia M, Rastrelli L, et al. Neuroprotective Effects of Quercetin: From Chemistry to Medicine. *CNS Neurol Disord Drug Targets* 2016; 15: 964-975.
 23. Sohn US, Lee SE, Lee SH, Nam Y, Hwang WK, Sohn UD. The protective mechanism of quercetin-3-O- β -D-glucuronopyranoside (QGC) in H₂O₂-induced injury of feline esophageal epithelial cells. *Arch Pharm Res* 2016; 39: 1324-1334.
 24. Shahzad H, Giribabu N, Karim K, Kassim N, Muniandy S, Kumar KE, et al. Quercetin interferes with the fluid volume and receptivity development of the uterus in rats during the peri-implantation period. *Reprod Toxicol* 2017; 71: 42-54.
 25. Cortázar TM, Coombs GH, Walker J. Leishmania panamensis: comparative inhibition of nuclear DNA topoisomerase II enzymes from promastigotes and human macrophages reveals anti-parasite selectivity of fluoroquinolones, flavonoids and pentamidine. *Exp Parasitol* 2007; 116: 475-482.
 26. Crespo I, García-Mediavilla MV, Almar M, González P, Tuñón MJ, Sánchez-Campos S, et al. Differential effects of dietary flavonoids on reactive oxygen and nitrogen species generation and changes in antioxidant enzyme expression induced by proinflammatory cytokines in Chang Liver cells. *Food Chem Toxicol* 2008; 46: 1555-1569.
 27. Liu XF, Xiang L, Zhou Q, Carralot JP, Prunotto M, Niederfellner G, et al. Actinomycin D enhances killing of cancer cells by immunotoxin RG7787 through activation of the extrinsic pathway of apoptosis. *Proc Natl Acad Sci U S A* 2016; 113: 10666-10671.
 28. Moura MT, de Sousa RV, de Oliveira Leme L, Rumpf R. Analysis of actinomycin D treated cattle oocytes and their use for somatic cell nuclear transfer. *Anim Reprod Sci* 2008; 109: 40-49.
 29. Bayir H, Kagan VE. Bench-to bedside review: Mitochondrial injury, oxidative stress and apoptosis--there is nothing more practical than a good theory. *Crit Care* 2008; 12: 206.
 30. Jazvinščak Jembrek M, Vuković L, Puhović J, Erhardt J, Oršolić N. Neuroprotective effect of quercetin against hydrogen peroxide-induced oxidative injury in P19 neurons. *J Mol Neurosci* 2012; 47: 286-299.
 31. Carrasco-Pozo C, Mizgier ML, Speisky H, Gotteland M. Differential protective effects of quercetin, resveratrol, rutin and epigallocatechin gallate against mitochondrial dysfunction induced by indomethacin in Caco-2 cells. *Chem Biol Interact* 2012; 195: 199-205.
 32. Nabavi SF, Nabavi SM, Latifi AM, Mirzaei M, Habtemariam S, Moghaddam AH. Mitigating role of quercetin against sodium fluoride-induced oxidative stress in the rat brain. *Pharm Biol* 2012; 50: 1380-1383.
 33. Sovrnigo TC, Adona PR, Monzani PS, Guemra S, Barros F, Lopes FG, et al. Effects of supplementation of medium with different antioxidants during in vitro maturation of bovine oocytes on subsequent embryo production. *Reprod Domest Anim* 2017; 52: 561-569.
 34. Adler A, Lee HL, McCulloh DH, Ampeloquio E, Clarke-Williams M, Wertz BH, et al. Blastocyst culture selects for euploid embryos: comparison of blastomere and trophectoderm biopsies. *Reprod Biomed Online* 2014; 28: 485-491.
 35. Basirat Z, Adib Rad H, Esmailzadeh S, Jorsaraei SG, Hajian-Tilaki K, Pasha H, et al. Comparison of pregnancy rate between fresh embryo transfers and frozen-thawed embryo transfers following ICSI treatment. *Int J Reprod Biomed* 2016; 14: 39-46.
 36. Aflatoonian A, Karimzadeh Maybodi MA, Aflatoonian N, Tabibnejad N, Amir-Arjmand MH, Soleimani M, et al. Perinatal outcome in fresh versus frozen embryo transfer in ART cycles. *Int J Reprod Biomed* 2016; 14: 167-172.
 37. Matos FD, Rocha JC, Nogueira MF. A method using artificial neural networks to morphologically assess mouse blastocyst quality. *J Anim Sci Technol* 2014; 56: 15.
 38. Rocha JC, Passalia F, Matos FD, Maserati MPJr, Alves MF, Almeida TG, et al. Methods for assessing the quality of mammalian embryos: How far we are

- from the gold standard? *JBRA Assist Reprod* 2016; 20: 150-158.
39. Zullo G, Albero G, Neglia G, De Canditiis C, Bifulco G, Campanile G, et al. L-ergothioneine supplementation during culture improves quality of bovine in vitro-produced embryos. *Theriogenology* 2016; 85: 688-697.
40. Du QY, Wang EY, Huang Y, Guo XY, Xiong YJ, Yu YP, et al. Blastocoele expansion degree predicts live birth after single blastocyst transfer for fresh and vitrified/warmed single blastocyst transfer cycles. *Fertil Steril* 2016; 105: 910-919.
41. Bain NT, Madan P, Betts DH. The early embryo response to intracellular reactive oxygen species is developmentally regulated. *Reprod Fertil Dev* 2011; 23: 561-575.
42. Karlsson JO, Szurek EA, Higgins AZ, Lee SR, Eroglu A. Optimization of cryoprotectant loading into murine and human oocytes. *Cryobiology* 2014; 68: 18-28.
43. Balakier H, Sojecki A, Motamedi G, Bashar S, Mandel R, Librach C. Is the zona pellucida thickness of human embryos influenced by women's age and hormonal levels? *Fertil Steril* 2012; 98: 77-83.
44. Montag M, Koll B, Holmes P, van der Ven. Significance of the number of embryonic cells and the state of the zona pellucida for hatching of mouse blastocysts in vitro versus in vivo. *Biol Reprod* 2000; 62: 1738-1744.