

## Effect of *Nelumbo nucifera* seeds on the reproductive organs of female rats

Anju Mutreja M.Sc., Meera Agarwal M.Sc., Ph.D., Sonalika Kushwaha M.Sc., Alka Chauhan M.Sc.

Reproductive Physiology and Biochemistry Section, Center for Advanced Studies, Department of Zoology, University of Rajasthan, Jaipur, India.

Received: 4 October 2007; accepted: 10 March 2008

### Abstract

**Background:** *Nelumbo nucifera* has been used as antifertility agent in females by the local tribals of Rajasthan especially of Udaipur district India.

**Objective:** The present study was conducted to investigate the effect of *Nelumbo nucifera* on the fertility and general physiology in female rats.

**Materials and Methods:** 20 female albino rats were taken and divided into two groups. Group 1 served as control and group 2 received ethanolic extract at dose of 800mg/kg b.wt. for investigating the nature of the drug and antifertility effect. Vaginal smear was monitored everyday in the morning to study the estrous cycle. On day 41<sup>st</sup> all the animals were sacrificed and various haematological and biochemical parameters were estimated.

**Results:** Data revealed that oral administration of *Nelumbo nucifera* extract brought about a significant decline in the weight of Ovary; Control (43±4.75mg), *Nelumbo* extract treated (25±3.86mg), Uterus; Control (236±0.004mg), *Nelumbo* extract treated (214±0.007mg) and Vagina; Control (221±0.002mg), *Nelumbo* extract treated (178±0.003mg) as well as protein and glycogen level, however cholesterol level increased significantly. In addition, the diestrous phase of the estrous cycle was found to be prolonged; Control (1.81±0.21) days, *Nelumbo* extract treated (3.62±0.42) days.

**Conclusion:** These results suggest that *Nelumbo nucifera* has the anti-estrogenic nature without altering the general physiology of the female rats.

**Key words:** *Anti-estrogenic, Female rats, Nelumbo nucifera, Estrous cycle.*

### Introduction

The development of new fertility regulating drug from medicinal plants is an attractive proposition. A wide variety of synthetic contraceptive agents are available but these are not without side effects. Plants like *Prangos ferulacia* has been reported to have abortifacient effect on the pregnant rats (1).

Antifertility effect has been studied in the plant *Woodfordia fruticosa* (2). Antisteroidogenic activity of two Indian medicinal plants, *Croton*

*roxburghii* and *Zizyphus jujuba* has been observed in mice (3). *Nelumbo nucifera* commonly known as Lotus and Kamala in Hindi belongs to family Nymphaeacea. Presences of various alkaloids have been reported from the entire plant including nuciferine, neferine, lotusine and isoliensinine. The seeds of *Nelumbo nucifera* contain 2-3% oil comprised of myristic, palmitic, oleic and linoleic acid. Mukherjee et al (4) showed that the rhizome extract of *Nelumbo nucifera* has antipyretic activity. Gupta et al (5) showed that the seed extract of *Nelumbo nucifera* has antisteroidogenic effect in the testis and ovary of the rat and Mazumdar et al (6) reported antifertility effect in mice. The present study was therefore carried out to study its effect on the reproductive organs and blood profile including total erythrocyte count, total leucocyte count, haematocrit, haemoglobin

#### Correspondence Author:

Dr. Meera Agarwal, Reproductive Physiology and Biochemistry Section, Center for Advanced Studies, Department of Zoology, University of Rajasthan, Jaipur- 302 004, India.

E-mail: drmeera\_agarwal@yahoo.co.in

and blood sugar for studying toxic side effects in female albino rats.

## Materials and methods

### Preparation of extract

The seeds of *Nelumbo nucifera* were purchased from the local market and authenticated by the Department of Botany, University of Rajasthan. The extract was prepared according to WHO protocol CG-04 (7) for the preparation of an alcoholic extract. The seeds were dried, powdered and then subjected to Soxhlet apparatus for extraction with 50% ethanol. The extract obtained was filtered and then evaporated to dryness under reduced pressure which yielded about 8.5% of solid residue.

### Animal study

Antifertility experiments were performed on inbred Wistar strain cyclic female albino rats weighing 200-210g. Twenty animals were maintained under standard husbandry conditions with food and water ad libitum.

Vaginal smear for each rat was monitored daily in the morning. Only rats with normal estrous cycle were selected for the experiment. To study the effect of ethanolic extract of *Nelumbo nucifera* on the estrous cycle, the selected animals were divided into two groups; group 1 served as control and group 2 received ethanolic extract at dose of 800 mg/kg b.wt.

The treatment was continued for 40 days. During this period the vaginal smear of the rats were examined daily in the morning hours. On day 41<sup>st</sup> the rats were sacrificed under light ether anesthesia. Ovary, uterus and vagina were dissected out, weighed and kept at -20°C for biochemical estimations. Blood was collected by cardiac puncture for haematological studies.

All the experiments were performed according to the guidelines of University Animal Ethics Committee.

### Acute toxicity Test

To determine the accurate dose level, acute toxicity test was performed and for this 3 separate animal groups were taken.

The study was conducted in female rats weighing (200±10g). The animals were divided into 3 groups.

The extract was administered orally starting from 500 mg/kg b.wt for ten days (0% mortality), 1gm/kg b.wt. for ten days (50% mortality) and 2gm/kg b.wt for ten days (100% mortality) doses

in 1 ml of vehicle. Records of mortality and manifestation of toxicity were made during 24 hr. Based on the mortality rate, probit values; the oral LD<sub>50</sub> was determined by McLeod method (8).

### Estrous cycle

Vaginal smear of each animal were evaluated as described by Vogel (9). The duration of the estrous cycle together with that of various phases was determined as described by Makonnen et al (10).

### Biochemical estimation

Protein and glycogen level were estimated according to the method of Lowry et al (11) and Montgomery (12) respectively. The cholesterol content was estimated according to Libermann and Burchard method (13).

### Total protein

Total protein was assayed by the method of Lowry et al (11). Briefly, the assay mixture contained; 1 N NaOH, reagent D (2%) Sodium tartarate, Folin's reagent, and 10% T.C.A. Protein reacts with Folin's reagent to give colour complexes. The intensity of this colour was measured at green filter against blank on colorimeter.

### Total cholesterol

Total cholesterol was determined by the method of Oser (13). Briefly it contains FeCl<sub>3</sub> solution, conc.

H<sub>2</sub>SO<sub>4</sub> and glacial acetic acid. The phenathrene ring of cholesterol reacts with FeCl<sub>3</sub> .7H<sub>2</sub>O and gives pinkish to brown colour depending upon the concentration of cholesterol.

### Glycogen

Glycogen was assayed by the method of Montgomery (12). Briefly the assay mixture contained 30% KOH, H<sub>2</sub>SO<sub>4</sub>, 80% phenol and absolute alcohol.

Polysaccharides are treated with conc. H<sub>2</sub>SO<sub>4</sub> and phenol due to which they undergo degradation and form a complex which is pink in colour. The intensity of this colour indicates intensity of glycogen in the tissue

### Blood and serum analysis

Total erythrocyte count and Haematocrit were measured by Microhaematocrit method (14). Total leucocyte count (15) was estimated. The haemoglobin level was estimated by cynomethanoglobin method (16) and blood sugar by Astoor and King method (17).

### Statistical analysis

Results are expressed as mean±SEM. Statistical analysis was done by student's t-test (18).

## Results

### Effects on body and organs weight

There was a non significant change in the body weight of the experimental animals but there was a significant decrease ( $p<0.01$ ) in the weight of the reproductive organs as compared to the control as follows: Ovary; Control ( $43\pm4.75$ ), Nelumbo extract treated ( $25\pm3.86$ ), Uterus; Control ( $236\pm0.004$ ), Nelumbo extract treated ( $214\pm0.007$ ) and Vagina; Control ( $221\pm0.002$ ), Nelumbo extract treated ( $178\pm0.003$ ) mg (Table I).

### Effect on tissue biochemistry


The glycogen decreased significantly ( $p<0.001$ ) in the: Ovary; Control ( $8.03\pm0.32$ ), Nelumbo treated ( $4.78\pm0.62$ ), Uterus; Control ( $18.31\pm0.50$ ), Nelumbo treated ( $11.66\pm0.81$ ), Vagina; Control ( $8.71\pm0.48$ ), Nelumbo treated ( $4.18\pm0.57$ ), and protein: Ovary; Control ( $184\pm4.49$ ), Nelumbo treated ( $134\pm3.94$ ), Uterus; Control ( $181.13\pm6.8$ ), Nelumbo treated ( $140\pm5.49$ ), Vagina; Control ( $169.37\pm7.84$ ), Nelumbo treated ( $119\pm4.66$ ). Where as there was a significant increase in the level of cholesterol of the experimental animals as compared to the control: Ovary; Control ( $8.84\pm0.55$ ), Nelumbo treated ( $12.08\pm0.44$ ), Uterus; Control ( $4.15\pm0.18$ ), Nelumbo treated ( $8.23\pm0.79$ ), Vagina; Control ( $5.17\pm0.14$ ), Nelumbo treated ( $8.41\pm0.65$ ) (Figures 1-3).

### Effect on haematological studies


There was a non significant change in haematological parameters. Total erythrocyte count, total leucocyte counts, haemoglobin, blood sugar and haematocrit values were found to be within normal range when compared to control (Table III).

### Effect on estrous cycle of rats


Ethanollic extracts of seeds of *Nelumbo nucifera* prolonged the estrous cycle significantly ( $p<0.05$ ). There had been a significant increase ( $p<0.01$ ) in the length of the diestrus phase (Control;  $1.81\pm0.21$  days, Nelumbo treated;  $3.62\pm0.42$  days) (Table II).



**Figure 1.** Effect of *Nelumbo nucifera* 50% ethanolic extract treatment on the protein content of ovary, uterus and vagina of female albino rats.



**Figure 2.** Effect of *Nelumbo nucifera* 50% ethanolic extract treatment on the glycogen content of ovary, uterus and vagina of female albino rats.



**Figure 3.** Effect of *Nelumbo nucifera* 50% ethanolic extract treatment on the cholesterol content of ovary, uterus and vagina of female albino rats.

**Table I.** Body and organ weight changes in female albino rats after the administration of 50 % ethanolic extract of *Nelumbo nucifera*.

Groups	Body weight (g)		Reproductive organs weight (mg)		
	Initial	Final	Ovary	Uterus	Vagina
Group 1: Control	200.20±2.85	209.80±2.57 <sup>ns</sup>	43±4.75	236±0.004	221±0.002
Group 2: <i>Nelumbo nucifera</i> treated	205.00±1.41	214.70±2.71 <sup>ns</sup>	25±3.86*	214±0.007*	178±0.003*

(Mean ±SEM of 10 animals)

Group 2 compared with group 1: \* = significant (p≤0.01), ns = non-significant

**Table II.** Effects on estrous cycle of female albino rats after the administration of 50 % ethanolic extract of *Nelumbo nucifera*.

Phases	Estrous cycle	Proestrous phase	Estrous phase	Metaestrous phase	Di estrous phase
Group 1: Control	4.28±0.12	0.82±0.09	0.83±0.18	0.82±0.16	1.81±0.21
Group 2: <i>Nelumbo nucifera</i> treated	5.46±0.51*	0.45±0.12*	0.89±0.19*	0.48 ±0.21*	3.62±0.42**

(Mean ±SEM of 10 animals)

Group 2 compared with group 1: \*\*=highly significant (p≤0.001), \* = significant (p≤0.01) ns = non-significant

**Table III.** Haematological changes in female albino rats after the administration of 50 % ethanolic extract of *Nelumbo nucifera*.

Groups	RBC Count million/mm <sup>3</sup>	WBC Count per. cu. mm.	Haemoglobin gm%	Haematocrit %	Blood Sugar mg/100 ml
Group 1: Control	3.52±0.45	6.4±0.44	13.49±0.56	49.38±1.39	81.32±0.66
Group 2: <i>Nelumbo nucifera</i> treated	3.80±0.06 <sup>ns</sup>	6.7±0.30 <sup>ns</sup>	13.17±0.42 <sup>ns</sup>	43.19±0.99 <sup>ns</sup>	81.41±0.81 <sup>ns</sup>

(Mean ±SEM of 10 animals)

Group 2 compared with group 1: ns = non-significant

## Discussion

Plants have the property to inhibit the estrogen surge for implantation (19). In mice and humans, estrogens play a pivotal role in the implantation since they participate in the estrogen, progesterone balance and thereby in the uterine receptivity to the embryo (20).

Administration of *Nelumbo nucifera* to female rats caused estrogen inhibition due to its antiestrogenic nature. The decrease in the weight of ovary and uterus shows antiestrogenic nature of *Nelumbo nucifera* since antiestrogenic substance decreases the wet weight of the uterus (21).

The prolonged estrous cycle and diestrous phase observed with the extract suggests the antifertility effect of *Nelumbo nucifera* seeds. The prolongation of diestrous phase may explain the remote chances of the rats to get pregnant. The results obtained in the present study on the

prolonged estrous cycle and on its diestrous phase methanolic root extract of *Rumex steudelii* (22).

Cholesterol derived from the different sources is the precursor for the steroidogenesis of ovarian endocrine tissue (23).

The significant increase in the cholesterol level of the group receiving extract indicates that cholesterol was not used for steroidogenesis hence accumulated in the ovary (24).

The decrease in the glycogen content of *Nelumbo nucifera* treated uterus confirms the antiestrogenic nature of the drug (25). Reduction in protein content of the female genital tract of *Nelumbo nucifera* treated rats suggests an inhibition of estrogen production (26).

Blood parameters have been found to be within normal range indicating non toxic action of *Nelumbo* extract on general body metabolism. Further work on the mechanism of antifertility action and histological changes of the genital organs are in progress.

## Conclusion

Over all, it may be concluded that the ethanolic extract of *Nelumbo nucifera* seeds has anti-estrogenic effect in female rats.

## Acknowledgement

The authors are thankful to the Center for Advanced Studies, Department of Zoology, University of Rajasthan, Jaipur, for providing necessary facilities.

## References

1. Talieh K, Kazem M, Alireza A, Mohsen S, Alireza S. Abortifacient effect of *Prangos ferulacia* on pregnant rats. *Contraception* 2006; 73: 554-556.
2. Khushalani H, Taltke P, Singh K. Antifertility activity of dried flowers of *Woodfordia fruticosa* kurz. *Indian Journal of Pharmaceutical Sciences* 2006; 68: 528-529.
3. Gupta M, Mazumdar UK, Vamsi MLM, Sivakumar T, Kandar CC. Antisteroidogenic activity of the two Indian medicinal plants in mice. *Journal of Ethnopharmacology* 2004; 90: 21-25.
4. Mukherjee PK, Das J. Antipyretic activity of *Nelumbo nucifera* rhizome extract. *Indian Journal of Experimental Biology* 1996; 34: 275-276
5. Gupta M, Mazumdar UK. Antisteroidogenic effect of the seed extract of *Nelumbo nucifera* in the testis and ovary of rat. *Indian Journal of Pharmaceutical Sciences* 1996; 58: 326-242.
6. Mazumdar UK, Gupta M. Antifertility activity of seeds of *Nelumbo nucifera* in mice. *Indian Journal of Experimental Biology* 1992; 30: 533-534.
7. WHO: protocol CG-04. Preparation of alcoholic extract for bioassay and phytochemical studies (APJF/IP, 1001 A). Geneva, World Health Organization 1983 a.
8. McLeod LJ. Pharmacological experiments on intact preparation. Treatment of all or none principle, First ed. Churchill Livingstone Ltd., Edinburgh 1976: 105
9. Vogel HG. Ovarian hormones. In: Pharmacological Assay, 2<sup>nd</sup> ed. Springer-verlag, Berlin, Heidelberg, 1997: 637-645
10. Makonnen E, Rostum Amr AH, Assefa G, Zerihun L. Antifertility effect of *Jatropha curcus* L. seeds in guinea pigs. *Ethiopian Journal of Health Development* 1997; 11: 145-148.
11. Lowry OH, Rosen burgh NJ, Farr AL, Ran dell RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951; 193: 265-275
12. Montgomery R. Determination of Glycogen Arch. *Biochem Biophys* 1957; 67: 378.
13. Oser BL. In: Hawk's Physiological Chemistry 14<sup>th</sup> Ed. Mc Graw Hill, New York, 1965: 246
14. Schalm OM, Jain NC, Carrolt EJ. Veterinary haematology 3<sup>rd</sup> ed, Lea and Febiger Philadelphia, 1975: 324-335.
15. Lynch JM, Raphael SS, Melvir LD, Spare PD, Inwood MJM. In: medical laboratory and clinical pathology pub, Saunders Company Sohm Ltd., Tokyo, 1969: 626, 647-662.
16. Makrem A. Haemoglobin, myoglobin and hepatoglobin. In: Henry Cannon Winkelmann. Ed. Clinical Chemistry. Principles and techniques. 1969; 1111-1214
17. Astoor A, King EJ. Simplified calorimetric blood sugar method. *J Biol Chem* 1954; 50 XIV.
18. Ipstein J, Poly F. Brancroft's Introduction to biostatics, 2<sup>nd</sup> ed. Harper International, New York : 44
19. Gandhi M, Lal R, Sankaranarayanan, Sharma PL. Post-coital antifertility activity of *Ruta graveolens* in female rats and hamsters. *J Ethnopharm* 1991: 49-59.
20. Denker HW. Cell biology of endometrial receptivity and of trophoblast- endometrial interaction .In: Glasser SP, Mulholland J, Psychoyos A. Endocrinology of embryo-endometrium interactions, Plenum Press, New York, 1994: 17-32.
21. Mukherjee P. Quality control Herbal drugs: an approach to evaluation of botanicals, First ed. Business Horizons, New Delhi 2002.
22. Gebrie E, Makonnen E, Debella A, Zerihun L. Phytochemical screening and pharmacological evaluations for the antifertility effect of the methanolic root extract of *Rumex steudelii*. *J Ethnopharm* 2005; 96: 139-143.
23. Strauss JF III, Schuler LA, Tanaka T. Cholesterol metabolism by ovarian tissue. *Advances in Lipid Research* 1981; 18: 99.
24. Shivalingappa H, Satyanarayan ND, Purohit MG, Sharanabasappa A, Patil SBJ. Effect of ethanol extract of *Rivea hypocrateriformis* on the estrous cycle of the rat. *Ethnopharm* 2002; 82: 11-17.
25. Agarwal M, Dixit VP, Sandhu JS. Possible mechanism of antifertility activity of 3-chloro-1, 2-Propanediol (U-5897) on the female genital tract of *Rattus rattus Rufescens*- A Biochemical and Histophysiological study. *Proc Ind Natl Sci Acad* 1980; 46: 293-301.
26. Mohla S, Prasad MRN. Oestrogen antiestrogen interaction: Effect of U-11100 A, MRL-41 (Clomiphene) and U-11555 A on oestrogen induced uterine glycogen and protein synthesis in the rat during delayed implantation. *Acta Endo* 1969; 162: 482-487.

