

## Change in sex hormone profile due to exposure to nuclides in nuclear detonation crisis: a topic to be discussed in reproductive medicine

Dear Editor,

The big public health concern in this year is the nuclear crisis in Japan that causes contamination of nuclides around the world. The effect of the exposure to leaked nuclides from the present nuclear crisis on reproductive health is a topic to be concerned. In my previous publication (1), the evidence on the relationship between nuclear exposure and infertility can be confirmed. However, there are also other adverse effects on the reproductive system of the exposed subjects.

An interesting topic is the disturbance of the normal sex hormone system. There are some reports on the effect of exposure on sex hormone profile in exposed subjects. Based on the data from the referencing nuclear crisis, the Chernobly crisis, the change in the level of testosterone in exposed subjects were controversial. Some reports mentioned for low testosterone level (2) while the others noted for rising level (3-5) among exposed subjects.

However, more reports points to the increasing level of testosterone. Nevertheless, the observation on the increased testosterone level in adolescent offspring of exposed subjects was also reported (5). Therefore, the problem of testosterone hormone alteration is a problem in reproductive health of the exposed subjects and this might lead to infertility (1). Hence, this is an interesting topic to be studied in the present Fukushima crisis. Unlike the report on testosterone, there are few reports on estrogen. The interesting report is on the estrogen related cancers in females exposed to the nuclides in the Chernobyl crisis. The study on breast cancers revealed that "BRCA1 mutations were strongly associated with earlier age at diagnosis, with estrogen receptor (ER) negative tumors (6)."

However, there is a report on the change of estrogen and progesterone in animals in the contaminated areas (7). For FSH and LH, although FSH and LH are not the actual sex hormones from gonad, the change of FSH and LH is another interesting issue. According to the study of Goncharov *et al* (4), there was a significant lowering of LH level but there was no significant change in FSH. However, in other studies, the increased FSH indicating the infertility was observed (2, 8).

In conclusion, there are evidences on sex hormone change in the subjects exposed to nuclides from nuclear accident crisis. This is an important issue in reproductive medicine that should be the focus in following up of the exposed subjects in the present nuclear crisis.

**Viroj Wiwanitkit M.D.**

Wiwanitkit House, Bangkhuae, Bangkok Thailand.

**Corresponding Author:**

Viroj Wiwanitkit, Wiwanitkit House, Bangkhuae, Bangkok, 10160, Thailand.

Email: wviroj@yahoo.com

Tel/Fax: 6624132436

Received: 7 July 2011

Accepted: 30 October 2011

## References

1. Wiwanitkit V. Nuclear denotation and infertility: evidence. *J Hum Reprod Sci* 2011; 4: 103-104.
2. Birioukov A, Meurer M, Peter RU, Braun-Falco O, Plewig G. Male reproductive system in patients exposed to ionizing irradiation in the Chernobyl accident. *Arch Androl* 1993; 30: 99-104.
3. Mishchenko VP, Gritsaï NN, Litvin AA, Sokolenko VN, Tserbrzhinskiï OI, Kutsenko LA, et al. Human physiological blood protection systems in the late periods after ionizing radiation exposure related to the accident at the Chernobyl Atomic Electric Power Station. *Gematol Transfuziol* 1993; 38: 30-33.
4. Goncharov NP, Katsiya GV, Kolesnikova GS, Dobracheva GA, Todua TN, Vax VV, et al. Endocrine and reproductive health status of men who had experienced short-term radiation exposure at Chernobyl. *Int J Androl* 1998; 21: 271-276.
5. Huizink AC, Bartels M, Rose RJ, Pulkkinen L, Eriksson CJ, Kaprio J. Chernobyl exposure as stressor during pregnancy and hormone levels in adolescent offspring. *J Epidemiol Community Health* 2008; 62: e5.
6. Bogdanova NV, Antonenkova NN, Rogov YI, Karstens JH, Hillemanns P, Dörk T. High frequency and allele-specific differences of BRCA1 founder mutations in breast cancer and ovarian cancer patients from Belarus. *Clin Genet* 2010; 78: 364-372.
7. Belov AD, Lysenko NP, Fomichëva NA. Evaluation of biological sequelae for cattle in the zone of the Chernobyl accident. *Radiats Biol Radioecol* 1997; 37: 629-639.

8. Gottlöber P, Steinert M, Weiss M, Bebeshko V, Belyi D, Nadejina N, et al. The outcome of local radiation injuries: 14 years of follow-up after the Chernobyl accident. *Radiat Res* 2001; 155: 409-416.