

Effects of selenium, calcium and calcium ionophore on human oocytes in vitro maturation in a chemically defined medium

Mehri Makki¹ M.Sc., Elham Saboorti² M.Sc., Mohammad Ali Sabbagh^{2,3} M.Sc., Raheleh Aram² M.Sc., Mohammad Hasan Fallahian⁴ M.Sc., Fatemeh Peyghambari⁵ Ph.D., Hessam Roustaei² M.D., Abbas Ahmadi⁶ Ph.D.

¹ Department of Midwifery, Islamic Azad University, Meybod Branch, Meybod, Iran.

² Department of Assisted Reproduction Technology, Laboratory of Embryology, Novin Fertility and Infertility Center, Mashhad, Iran.

³ Department of Biology, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad, Iran.

⁴ Department of Chemistry, Islamic Azad University, Meybod Branch, Meybod, Iran.

⁵ Department of Anatomy, Islamic Azad University, Yazd Branch, Yazd, Iran.

⁶ Department of Basic Sciences, Laboratory of Embryology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

Corresponding Author:

Abbas Ahmadi, Department of Basic Sciences, Laboratory of Embryology, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran.

Email: ivf_center@yahoo.com
Tel/Fax: (+98) 9141498524

Received: 2 May 2011

Revised: 17 August 2011

Accepted: 27 August 2011

Abstract

Background: In vitro maturation (IVM) of human oocytes is an emerging procedure quickly incorporated into the world of assisted reproductive technologies. As an effective method of in vitro maturation, several studies have reported the critical role of differentiations on activating the complex process involved in both gamete maturation and fertilization.

Objective: In this study, we supplemented a chemically defined medium with different combinations of selenium, calcium and calcium ionophore concentrations to obtain the best rate of human oocytes maturation, survival, and fertilization.

Materials and Methods: As an experimental study, Three combinations of [selenium (5 μ g/ml), calcium (5 μ g/ml) and calcium ionophore (1 μ g/ml)], [selenium (10 μ g/ml), calcium (7 μ g/ml) and calcium ionophore (2 μ g/ml)] and [selenium (15 μ g/ml), calcium (10 μ g/ml) and calcium ionophore (5 μ g/ml)] added to the chemically defined medium and the morphology of oocytes assessed after 22-24 hours in vitro maturation of the oocytes.

Results: The highest percentage of MII (meiosis II) oocytes (68%), developing beyond the morula (20.1%) and the blastocyst formation (11.1%) observed in oocytes treated with 15 μ g/ml selenium, 10 μ g/ml calcium and 5 μ g/ml calcium ionophore. Moreover, we showed the significant rate of survival in each three combinations after 36, 72 and 96 hours.

Conclusion: Maturation and activation of oocytes may be triggered by changes in intracellular ion concentrations as second messengers in signal transduction pathways. Here, we received the highest percentage of in vitro maturation and fertilization among three combinations of selenium, calcium and calcium ionophore treatments. Using this combination of ions beside other factors might be useful for the enrichment of the human oocytes IVM medium.

Key words: Human oocytes, In vitro maturation (IVM), Selenium, Calcium, Calcium ionophore.

Introduction

In vitro maturation of oocytes encompasses a variety of cellular processes must be completed in order to conception and development into normal embryos and offspring (1, 2). Minimizing hormone's analogues usage, elimination of ovarian hyperstimulation syndrome (OHS), also simplicity of the protocol are some of the clinical potential advantages of this new technology (3). Furthermore, the conservation of genetic resources, cloning by nuclear transfer, either for reproductive purposes or the generation of embryonic stem cells in

regenerative medicine, will belong to the large number of high-quality oocytes (4).

Oocytes growth occurs primarily in preantral follicles where it arrested in prophase I of meiosis. Then, follicles undergo an increase in size and specialization of granulosa cells into cumulus and mural granulosa cells. At the final stage, follicle maturation involves the period between luteinizing hormone (LH) surge and ovulation. During this time, the oocytes resume meiosis and complete their cytoplasmic maturation (5, 6). The corresponding process takes place during in vitro maturation to achieve a high number of oocytes. Several supplementary factors have used for in vitro maturation of

oocytes both in human and other species. Fetal bovine serum (7), patient's own follicular fluid (FF) (8) and/or patient's own serum (9) were the earlier supplementation that used for human oocytes maturation.

It has shown that changes in enormous ion concentrations can serve as second messengers of signal transduction in the different systems (10). In recent years, several studies have investigated the roles of different ions in order to oocytes maturation. The effects of ammonium (11), sodium pyruvate (12, 13), calcium (14, 15), nitrogen oxide (NO) (16) and selenium (17, 18) have examined on oocytes maturation and subsequent embryonic development.

In this study, we used three combinations of selenium, calcium and calcium ionophore concentrations in a "chemically defined medium" to compare the developmental competency of immature human oocytes and achieve highest rate of oocytes maturation. Using different ion concentrations beside other factors might be useful in developing well defined human in vitro maturation medium for various basic and clinical purposes.

Materials and methods

Seventy six 25-40 years old women attending in Novin Fertility and Infertility Center of Mashhad for in vitro fertilization (IVF) and intra cytoplasmic sperm injection (ICSI) invited to participate in this study. An information sheet gave to all patients and informed written consent obtained from those agreeing to take part and all of the sheets documented in this center. All the procedure of surgery, in vitro and in vivo maturation was performed according to the human ethical protocols of Ministry of Health and Medical Education of Iran.

Experimental design

In this study, selenium (Sigma, USA), calcium (Merck) and calcium ionophore A23187 (Sigma, USA) used in a chemically defined medium (SAGE, USA). Human GV (Germinal vesicles) oocytes evaluated in three experiments (six replicates in each experiment). Experiment 1, examined the effects of selenium (5 μ g/ml), calcium (5 μ g/ml) and calcium ionophore (1 μ g/ml) on in vitro maturation of the oocytes.

Experiment 2, examined the effects of selenium (10 μ g/ml), calcium (7 μ g/ml) and calcium ionophore (2 μ g/ml) supplementation on GV oocytes. Ultimately, Experiment 3 assessed the effects of selenium (10 μ g/ml), calcium (7 μ g/ml) and calcium ionophore (5 μ g/ml) in a chemically defined medium. As control group, GV oocytes evaluated in SAGE as a chemically defined medium without the treatments.

Semen preparation

Semen samples from a single ejaculation rapidly thawed (30 seconds) in a water bath at 37°C. Total motility after thawing was 70%, with 50-60% progressive motility. Sperms prepared by the swim-up procedure for ICSI. The optical microscopy based method performed to determine the concentration of sperms, Furthermore, one drop of sperm suspension placed on a microscopic slide, and the movement of 200 sperm cells examined using a light microscope in $\times 40$ magnifications. Besides that, at least 200 sperms from each sample examined for sperm's quality assessment. The semen quality evaluated according to the laboratory manual for the examination and processing of human semen by WHO as described in 2010 (19).

Oocytes preparation and evaluation

According to the center's protocol, stimulation achieved by the administration of two injections of gonadotropin each day. When the ovaries were ready, the patients received an HCG injection. The egg retrieval process took place 34-36 hours after the HCG injection. Oocytes collected from the ovary and transfer to medium supplemented by different combinations of selenium, calcium and calcium as in vitro maturation medium.

After 22-24 hours, oocytes denuded of cumulus and corona cells by incubation with 80 IU hyaluronidase/ml (Sigma, USA) and in/out aspiration with finely drawn glass pipettes. Oocyte that showing an extruded 1st polar body submitted to microinjection after denuding (20).

In vitro culture of oocytes

Injected oocytes placed in drops of medium under sterile mineral oil at 37°C in a humidified atmosphere of 5% (v/v) CO₂ after

microinjection. Once daily, embryos assessed and their developmental stage recorded.

Statistical analysis

All data converted to proportions using 2-proportion analysis. Within each experiment, data reported as percentages. P-values of less than 0.05 considered statistically significant. The data analyzed by one way ANOVA (SPSS version 11.5).

Results

The quality of in vitro matured oocytes

In order to evaluate the effects of selenium, calcium and calcium ionophore on in vitro maturation, we assessed the quality of oocytes in various developmental stages after different combinations of treatments. The number of MI, MII and abnormal oocytes counted in different time lines (8, 16 and 24 hours). We observed the highest percentage of MII oocytes (68%) in oocytes treated with 15 µg/ml selenium, 10 µg/ml calcium and 5 µg/ml calcium ionophore after 24 hours.

The significant reduction in MI oocytes also increases in MII oocytes percentage observed after 24 hours in comparison to 8 and 16 hours in comparison to the control group ($p<0.05$). Interestingly, there is no significant difference between the percentages of abnormal oocytes in three combinations of treatments in comparison to control group. In addition, the lowest percentages of abnormality (22.1%) observed in oocytes that treated with 15 µg/ml selenium, 10 µg/ml

calcium and 5 µg/ml calcium ionophore among each three time lines.

Fertilization and survival rate evaluation

Following ICSI, the normal fertilization rate of in vitro maturation evaluated. We showed that the proportion of normal oocytes was not significant after fertilization. Nevertheless, the highest percentage of fertilization (77.4%) observed in 15 µg/ml selenium, 10 µg/ml calcium and 5 µg/ml calcium ionophore after 72 hours (Table II). In addition, we observed the significant survival rate in all combinations after 36, 72 and 96 hours in comparison to the control group ($p<0.05$). As the same as the fertilization rate, the highest survival rate (84.4%) observed in oocytes treated with 15 µg/ml selenium, 10 µg/ml calcium and 5 µg/ml calcium ionophore (Table II).

Embryo development assessments of in vitro matured oocytes

In final step, the proportion of oocytes which is cleaved, developing beyond the morula and blastocyst assessed. Our results showed the significant percentage of cleavage in medium supplemented with all combinations. The highest rate of developing beyond the morula (20.1%) and blastocyst formation (11.1%) observed in oocytes treated with 15 µg/ml selenium, 10 µg/ml calcium and 5 µg/ml calcium ionophore.

Nevertheless, we observed the significant increase in percentage of oocytes proceed to blastocyst in all combinations in comparison to control group ($p<0.05$ and $p<0.01$) (Table III).

Table I. Mean assessment of oocytes quality at various stages of *in vitro* maturation (six replicated experiments).

Time (hours)	Treatments	No. of oocytes	MI (%)	MII (%)	Abnormal (%)
8	Se (5µg/ml), Ca (5µg/ml), CaI (1 µg/ml)	32	68.2	1.4	30.4
	Se(10µg/ml), Ca (7µg/ml), Cal (2 µg/ml)	34	61.6	1.7	36.7
	Se (15µg/ml), Ca (10µg/ml), Cal(5µg/ml)	34	68.1	2.1	29.8
	Control	30	69.2	1.2	29.6
16	Se (5µg/ml), Ca (5µg/ml), CaI (1 µg/ml)	31	42.0	20.2	37.8
	Se(10µg/ml), Ca (7µg/ml), Cal (2 µg/ml)	32	38.4	22.6	39.0
	Se (15µg/ml), Ca (10µg/ml), Cal(5µg/ml)	31	38.2	22.9	38.9
	Control	29	40.0	20.2	39.8
24	Se (5µg/ml), Ca (5µg/ml), CaI (1 µg/ml)	30	13.5	60.4	26.1
	Se(10µg/ml), Ca (7µg/ml), Cal (2 µg/ml)	31	12.2	61.2	26.6
	Se (15µg/ml), Ca (10µg/ml), Cal(5µg/ml)	30	9.9*	68.0*	22.1
	Control	29	13.8	61.4	24.8

* $p<0.05$.

Table II. The effect of different concentration of selenium, calcium and calcium ionophore on GV oocytes *in vitro* (six replicated experiments).

Interval (hours)	Treatments	No. Injected	Survival rate (%)	Fertilization rate (%)
36	Se (5µg/ml), Ca (5 µg/ml), CaI (1µg/ml)	30	76.1*	71.6
	Se (10µg/ml), Ca (7 µg/ml), CaI(2µg/ml)	34	78.4*	70.9
	Se (15µg/ml), Ca (10µg/ml), CaI(5µg/ml)	31	81.6*	72.0
	Control	33	61.2	70.2
72	Se (5µg/ml), Ca (5 µg/ml), CaI (1 µg/ml)	31	78.4*	75.1
	Se(10µg/ml), Ca (7 µg/ml), CaI (2µg/ml)	35	79.8*	72.7
	Se (15µg/ml), Ca (10 µg/ml), CaI(5µg/ml)	34	84.4*	77.4
	Control	32	62.1	74.9
96	Se (5µg/ml), Ca (5 µg/ml), CaI (1 µg/ml)	34	80.7*	73.2
	Se(10µg/ml), Ca (7 µg/ml), CaI (2µg/ml)	30	83.1*	74.7
	Se (15µg/ml), Ca (10 µg/ml), CaI(5µg/ml)	30	82.2*	75.4
	Control	31	59.4	72.1

Se: Selenium , Ca: Calcium, CaI: Calcium Ionophore.

*p<0.05.

Table III. Embryo development of *in vitro* matured oocytes according to the interval between *in vitro* maturation and blastocyst formation.

Treatments	*No of oocytes activated	% **oocytes cleaved	%**oocytes developing beyond the morula stage	% **oocytes developing to the blastocyst
Se (5µg/ml), Ca (5 µg/ml), CaI (1 µg/ml)	32	62.3*	15.1*	7.1*
Se(10µg/ml), Ca (7 µg/ml), CaI (2µg/ml)	34	66.1*	16.6*	9.5*
Se (15µg/ml), Ca (10 µg/ml), CaI(5µg/ml)	34	76.2**	20.1**	11.1**
Control	30	50.1	10.3	5.4

Se: Selenium, Ca: Calcium, CaI: Calcium Ionophore.

*Oocytes were cultured for 24 h in chemically defined medium supplemented by Selenium, Calcium and Calcium Ionophore.

**Percentage based on the number of oocytes stimulated.

*p<0.05 and **p<0.01.

Discussion

To our knowledge, this is the first study to investigate the effects of selenium, calcium and calcium ionophore supplementation in a chemically define medium to receive the highest rate of IVM. Selenium is an essential trace element which may have antioxidant activity in biological systems (21-23). Here, we used selenium in combination of calcium and calcium ionophore. Calcium elevation is a crucial event in triggering the complex machinery involved gamete maturation and fertilization (24, 25).

Besides that, calcium ionophore, facilitates calcium current into the oocytes and helps the elevation of extracellular calcium concentration (26-28). The combination of these ions may be activating immature oocytes to receive the highest rate of maturation and fertilization. After counting the number of MI, MII and abnormal oocytes in

different time lines, our data showed the highest percentage of MII oocytes (68%) after 24 hours treatment with 15µg/ml selenium, 10 µg/ml calcium and 5µg/ml calcium ionophore.

Jeong et al (17) recommended the supplementation of the porcine IVM medium with 10 µg/ml insulin, 5.5 µg/ml transferrin and 5 µg/ml selenium. In addition, ITS plus L-ascorbic acid supplementation during the first 12 hours of in vitro maturation improves cytoplasm maturation (83.3%) in calf oocytes (18).

Choi et al (29) showed the highest rate of oocytes activation (82%) in TCM-199, treated with 50 µmol calcium ionophore and cultured with cycloheximide for 24 hours. Furthermore, 61% of oocytes that had cultured with Calcium-EDTA for 48 hours formed a pronucleus (30). The percentage of MII oocytes was not significant in other time lines (8 and 16 hours), except the treatment with 15 µg/ml selenium, 10 µg/ml calcium and 5 µg/ml calcium ionophore after 16 hours. There was

no difference between the percentages of abnormal oocytes in different combinations. It shows that these combinations do not cause to the abnormality in oocytes treated for in vitro maturation. The lowest percentage of abnormality was 22.1% among different treatments (Table I).

In the next step, we assessed the normal fertilization and survival percentage following in vitro matured oocytes ICSI. There was no significant difference in percentage of normal fertilization. The highest percentage of fertilization was 77.4% in medium supplemented with 15 µg/ml selenium, 10 µg/ml calcium and 5 µg/ml calcium ionophore after 72 hours. As the same as fertilization rate, we observed the highest rate of survival (84.4%) in oocytes treated with 15 µg/ml selenium, 10 µg/ml calcium and 5 µg/ml calcium ionophore. The survival rate is significant in all combinations (Table II). Treatment with ITS plus porcine follicular fluid significantly improved the fertilization parameter of oocytes (17).

At the final step, embryo development of in vitro matured oocytes evaluated. There was the significant difference between the percentages of oocytes that cleaved, developed beyond to the morula and blastocyst in all combinations, in compare to the chemically defined medium without treatment. The highest percentage of oocytes cleavage (76.2%) and developing beyond morula (20.1%) observed in medium supplemented with 15 µg/ml selenium, 10 µg/ml calcium and 5 µg/ml calcium ionophore. Furthermore, the highest percentage of blastocyst formation (11.1%) observed as the same group as later (Table III). When bovine oocytes cultured further in medium without Calcium-EDTA after 48 hours Calcium-EDTA supplementation, only 26% of the oocytes developed to the cleaved stage (30).

In conclusion, these results will allow future approaches on gamete in vitro maturation and the role of different ion concentrations on it. We need future studies to investigate the signaling pathways which involved in this process.

Acknowledgements

This work was supported by grants from Islamic Azad University (IAU), Meybod Branch and Novin center of fertility and infertility of

Mashhad. Authors would like to thank M. Fadavi and M. Daneshvar for their invaluable and excellent technical assistance in IVM. The authors wish to thank Dr. Mansoori for experimental design helps and her kind managements.

References

1. Glazier AM, Mate KE, Rodger JC. In vitro and in vivo maturation of oocytes from gonadotrophin-treated brushtail possums. *Mol Reprod Dev* 2002; 62: 504-512.
2. Heikinheimo O, Gibbons WE. The molecular mechanisms of oocytes maturation and early embryonic development are unveiling new insights into reproductive medicine. *Mol Hum Reprod* 1998; 4: 745-756.
3. Jurema MW, Nogueira D. In vitro maturation of human oocytes for assisted reproduction. *Fertil Steril* 2006; 86: 1277-1291.
4. Martín-Coello J, González R, Crespo C, Gomendio M, Roldan ER. Superovulation and in vitro oocytes maturation in three species of mice (Mus musculus, Mus spretus and Mus spicilegus). *Theriogenology* 2008; 70: 1004-1013.
5. Moor R, Dai Y. Maturation of pig oocytes in vivo and in vitro. *Reprod Suppl* 2001; 58: 91-104.
6. Moor RM, Dai Y, Lee C, Fulka J Jr. Oocytes maturation and embryonic failure. *Hum Reprod Update* 1998; 4: 223-236.
7. Younis AI, Brackett BG, Fayer-Hosken RA. Influence of serum and hormones on bovine oocytes maturation and fertilization in vitro. *Gamete Res* 1989; 23: 189-201.
8. Cha KY, Koo JJ, Ko JJ, Choi DH, Han SY, Yoon TK. Pregnancy after in vitro fertilization of human follicular oocytes collected from nonstimulated cycles, their culture in vitro and their transfer in a donor oocytes program. *Fertil Steril* 1991; 55: 109-113.
9. Chian RC, Buckett WM, Too LL, Tan SL. Pregnancies resulting from in vitro matured oocytes retrieved from patients with polycystic ovary syndrome after priming with human chorionic gonadotropin. *Fertil Steril* 1999; 72: 639-642.
10. Ben-Yosef D, Shalgi R. Early ionic events in activation of the mammalian egg. *Rev Reprod* 1998; 3: 96-103.
11. Yuan Y, Krisher RL. Effect of ammonium during in vitro maturation on oocytes nuclear maturation and subsequent embryonic development in pigs. *Anim Reprod Sci* 2010; 117: 302-307.
12. Geshi M, Takenouchi N, Yamauchi N, Nagai T. Effects of sodium pyruvate in nonserum maturation medium on maturation, fertilization, and subsequent development of bovine oocytes with or without cumulus cells. *Biol Reprod* 2000; 63: 1730-1734.
13. Gonzales-Figueroa H, Gonzales-Molino HM. Maturation of pig oocytes in vitro in a medium with pyruvate. *Braz J Med Biol Res* 2005; 38: 869-872.
14. Miyara F, Pesty A, Migne C, Djediat C, Huang XB, Dumont-Hassan M, et al. Spontaneous calcium oscillations and nuclear PLC-beta1 in human GV oocytes. *Mol Reprod Dev* 2008; 75: 392-402.

15. Asano A, Niwa K. Activation and penetration in vitro of pig oocytes treated with calcium ionophore. *J Reprod Dev* 2004; 50: 77-85.
16. Voznesens'ka T, Blashkiv TV. Effect of NO on meiotic maturation of the oocytes in mice in vitro. *Fiziol Zh* 2003; 49: 100-103.
17. Jeong YW, Shamim Hossein M, Bhandari DP, Kim YW, Kim JH, Park SW, et al. Effects of insulin-transferrin-selenium in defined and porcine follicular fluid supplemented IVM media on porcine IVF and SCNT embryo production. *Anim Reprod Sci* 2008; 106: 13-24.
18. Córdoba B, Morató R, Izquierdo D, Paramio T, Mogas T. Effect of the addition of insulin-transferrin-selenium and/or L-ascorbic acid to the in vitro maturation of prepubertal bovine oocytes on cytoplasmic maturation and embryo development. *Theriogenology* 2010; 74: 1341-1348.
19. Cooper TG, Noonan E, von Eckardstein S, Auger J, Baker HW, Behre HM, et al. World Health Organization reference values for human semen characteristics. *Hum Reprod Update* 2010; 16: 231-245.
20. Rienzi L, Vajta G, Ubaldi F. Predictive value of oocytes morphology in human IVF: a systematic review of the literature. *Hum Reprod Update* 2011; 17: 34-45.
21. Wu SH, Oldfield JE, Whanger PD, Weswig PH, et al. Effect of selenium, vitamin E, and antioxidants on testicular function in rats. *Biol Reprod* 1973; 8: 625-629.
22. Gutteridge JM. Antioxidant properties of the proteins caeruloplasmin, albumin and transferrin. A study of their activity in serum and synovial fluid from patients with rheumatoid arthritis. *Biochim Biophys Acta* 1986; 869: 119-127.
23. Kara H, Cevik A, Konar V, Dayangac A, Yilmaz M. Protective effects of antioxidants against cadmium-induced oxidative damage in rat testes. *Biol Trace Elem Res* 2007; 120: 205-211.
24. Boni R, Gualtieri R, Talevi R, Tosti E. Calcium and other ion dynamics during gamete maturation and fertilization. *Theriogenology* 2007; 68: 156-164.
25. Ahmad K, Bracho GE, Wolf DP, Tash JS. Regulation of human sperm motility and hyperactivation components by calcium, calmodulin, and protein phosphatases. *Arch Androl* 1995; 35: 187-208.
26. Isachenko E, Isachenko V, Todorov P, Ostashko V, Kreienberg R, Kaufmann M, et al. Pregnancy after the calcium ionophore correction of pronuclei position in oocytes after intracytoplasmic sperm injection. *Fertil Steril* 2010; 94: 2770. e3-5.
27. Terada Y, Hasegawa H, Takahashi A, Ugajin T, Yaegashi N, Okamura K. Successful pregnancy after oocytes activation by a calcium ionophore for a patient with recurrent intracytoplasmic sperm injection failure, with an assessment of oocytes activation and sperm centrosomal function using bovine eggs. *Fertil Steril* 2009; 91: 935. e11-14.
28. Swain JE, Pool TB. ART failure: oocytes contributions to unsuccessful fertilization. *Hum Reprod Update* 2008; 14: 431-446.
29. Choi YH, Love CC, Varner DD, Thompson JA, Hinrichs K. Activation of cumulus-free equine oocytes: effect of maturation medium, calcium ionophore concentration and duration of cycloheximide exposure. *Reproduction* 2001; 122: 177-183.
30. Azuma T, Kondo T, Ikeda S, Imai H, Yamada M. Effects of EDTA saturated with Ca²⁺ (Ca-EDTA) on pig, bovine and mouse oocytes at the germinal vesicle stage during maturation culture and the involvement of chelation of Zn²⁺ in pronuclear formation induction by Ca-EDTA. *Reproduction* 2002; 124: 235-240.