Treatment of recurrent pregnancy loss by Levothyroxine in women with high Anti-TPO antibody

Mohammad Hosein Mosaddegh¹ Pharm D, Ph.D., Nasrin Ghasemi² M.D., Ph.D., Tahere Jahaninejad³ M.Sc., Fatemeh Mohsenifar⁴ M.D., Abbas Aflatoonian⁵ M.D.

- 1. Department of Pharmacology, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- 2. Department of Medical Genetics, Research and Clinical Centre for Infertility, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- 3. Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- 4. Yazd Azad University, Yazd, Iran.
- 5. Department of Obstetrics and Gynecology, Research and Clinical Centre for Infertility, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

Corresponding Author:

Nasrin Ghasemi, Department of Medical Genetics, Research and Clinical Centre for Infertility, Shahid Sadoughi University of Medical Sciences, Yazd, Iran. Email: n479g@yahoo.co.uk Tel/Fax: (+98) 3518247085-6

Received: 7 January 2012 Revised: 24 May 2012 Accepted: 10 June 2012

Abstract

Background: Recurrent pregnancy loss (RPL) is defined as two or more consecutive pregnancy losses before twenty weeks of gestation. It is caused by a variety of genetics and non-genetics factors. Thyroid autoimmunity could associate with pregnancy loss.

Objective: To investigate the effectiveness of Levothyroxine in treatment of RPL in women with high auto-thyroid antibodies.

Materials and Methods: In this observational cross sectional study, 900 women who had a history of recurrent pregnancy loss were studied. All women with high anti-TPO antibody without any other problems entered in this study. Levothyroxine was given to them two months before pregnancy till the end of pregnancy. The doses of levothyroxine were depended on the anti-TPO levels, which were decided by endocrinologist. Women followed for the results of pregnancies.

Results: The success rate of pregnancy in women with abnormal anti-TPO with Levothyroxine therapy was 82.85%. Mean of anti-TPO in women with treatment before taking medication was 488.35 and after that it was 123.35 UI/ml. This difference was significant (p<0.05). The mean of the antibodies was not significantly different in women without treatment.

Conclusion: This study showed that Levothyroxine reduces the incidence of spontaneous abortions in women with high Anti-TPO antibody. It decreased anti-TPO antibody levels after 2-3 months treatment.

Key words: Recurrent pregnancy loss, Levothyroxine, Anti-TPO antibody.

Introduction

ecurrent pregnancy loss (RPL) is defined as two or more consecutive pregnancy losses before twenty week of gestation, which affects 1-3% of couples. It is caused by variety of genetics and non-genetics factors. Genetic disorders, reproductive tract anatomical pathologies, infectious diseases, endocrine dysfunctions, thrombophilia and autoimmune diseases are known to be the most important risk factors for RPL (1-6).

thyroid There evidence is that autoimmunity is associated with pregnancy loss, because maternal thyroid hormones (TH) play a critical role in the development of both fetus and placenta (7). Pregnancy loss in women with positive thyroid autoantibodies occurs within the first trimester of gestation, when the fetus is dependent on maternal thvroid hormones (8. 9). Following implantation, the preservation the pregnancy is reliant on a mass of immunological events that will assistance in the successful growth and development of the fetus (7).

So, Thyroid autoantibodies generally enhanced autoimmune reactivity against the feto-placental unit as a consequence of hypothyroidism (10). The presence of thyroid autoantibodies can cause infertility and delay pregnancy. Women with high concentration thyroid autoantibodies in their blood circulation usually become pregnant in older age and encounter with a higher risk of pregnancy loss (11-13).

In the other hand, the presence of thyroid autoantibodies in women during pregnancy could be associated with a lack in thyroid hormone concentrations or a poorer ability of the thyroid gland to sufficiently conform to the requests of pregnancy. Treatment of thyroid insufficiency, caused by auto antibodies, during pregnancy is important in avoiding hostile maternal and fetal outcome (14).

This study assessed the effect of Levothyroxine therapy on the live-birth rate in women with a history of at least two recurrent unexplained pregnancy loss, which were anti-TPO antibody positive. It also compared antithyroid peroxidase antibodies levels before and after treatment.

Materials and methods

This project evaluated 900 women with recurrent pregnancy loss. They had referred to the Recurrent Abortion Clinic of Yazd Reproductive Sciences Institute. They were evaluated for uterine and cervical anatomical disorders (using pelvic ultrasonography or hysteroscopy), ovarian dysfunction, chromosome abnormality (using conventional karyotype), and then evaluated for thyroid disorders, and autoimmune antibodies. This study approved by ethic committee of yazd Reproductive Science Institute.

Women with positive thyroid peroxidase antibodies (anti-TPO) and normal TSH were entered to this study. Thyroid peroxidase (TPO) was tested with a chemiluminescence immunoassay, and women with anti-TPO more than 40 UI/ml were treated with levothyroxine after signing inform consent.

After two months anti-TPO was tested again. Levothyroxine doses were depended on the levels of anti-TPO, which were decided by endocrinologist. It was 25-100 µg every day. Treatment continued with levothyroxine and aspirin till pregnancy happened and these

continued during pregnancy until delivery. All women were followed during pregnancy to the end.

Statistical analysis

Mean of anti-TPO antibodies in each group were tested before and after 2 months with paired t-test using SPSS (version 16) and p-value<0.05 is significant.

Results

Forty five unexplained RPL women had anti-TPO more than 40 UI/ml with normal TSH. Mean of anti-TPO in women without treatment and with treatment 462.45 488.35 pregnancy were and respectively, which was not significantly different (p>0.05). Mean of anti-TPO in women after treatment was 123.35. The difference between mean before and after treatment was significant (p<0.05) (Table I).

Thirty nine women received Levothyroxine and six women never used Levothyroxine by their own decision. Four of 39 cases never get pregnant (in follow up time). From the remaining cases 35 treated with Levothyroxine, 14 had normal delivery, 9 passed 20 weeks of gestation, 6 had preterm labors with normal child, and 6 had miscarriages. Three of 6 women never used levothyroxine, get pregnant, but all had abortion again. Three of six cases never get pregnant again (Table II).

Table I. Mean of anti-TPO in women with recurrent abortion.

Patients		Anti-TPO (UI/ml)	p-value			
	Mean± SD					
With treatment						
	Before medication	488.35±113.58	< 0.05			
	After 2 months	123.35±38.7				
Without treatment						
	Before medication	462.45±149.33	>0.05			
	After 2 months	453.12±139.2				

Anti-TPO significantly decreased after treatment with levothyroxine (p-value<0.05).

Table II. Success rate of pregnancy in women with abnormal anti-TPO with and without treatment.

Levothyroxine	Normal delivery (n)	Preterm labor with normal child (n)	Pregnancy more than 20 weeks (n)	Abortion (n)	Sum	Successes rate
Take	14	6	9	6	35	82.85%
Not take	-	-	-	3	3	0%

Pregnancy rate after treatment with levothyroxine in women with high anti-TPO was 82.85%.

Discussion

Recurrent pregnancy loss is an important clinical problem and classically defined as two or more pregnancy losses before the fetus has reached viability. The leading etiologies associated with it include a variety of causes such as autoimmune diseases (15).

Nilsson et al in 1975 found for first time that levels of circulating anticoagulant were higher in women with recurrent pregnancy loss than in control subjects (16). This anticoagulant was later found to be the lupus anticoagulant antibody. Since that time several studies have established the relationship between autoimmune diseases such as antiphospholipid syndrome, rheumatoid arthritis and systemic lupus erythematous with abortion (17, 18).

A number of studies have revealed relation between thyroid autoantibodies and recurrent abortions. It is suggested that the presence of thyroid autoantibodies could cause a generalized activation of the immune system, which unregulated activity of the immune system at the fetal-maternal interface (19). Lejune *et al* found that frequency of circulating anti-thyroid antibodies were higher in women with recurrent abortion than in control subjects (20).

In 1990, Stagnaro-Green *et al* tested 552 women for thyroid auto-antibodies in the first trimester of pregnancy. Pregnancy loss rate in women with high antibody was twice than women with normal antibody (10). Regarding prevention of abortion, there are a few studies showing that thyroxin treatment may be efficient in decreasing the number of abortion when given during the early stages of pregnancy. Negro *et al* treated euthyroid women with high thyroid autoantibodies who underwent IVF.

Half of them received levothyroxine and other half received placebo. The pregnancy loss rate in the placebo group was 52% compared to 33% in the group in which levothyroxine was given. However, because of the small number of investigated patients in their study the difference was not statistically significant (21). In Roussev study, 187 patients were randomized for evaluating the

effect of levothyroxine treatment on pregnancy outcomes (22).

Also in Poppe study levothyroxine was prescribed for 187 women with unexplained recurrent pregnancy loss, and compare live birth rates among two groups (23). In both studies women with normal thyroid function with thyroid autoantibodies were selected. Both studies showed a decrease in pregnancy loss rates (36% and 75% relative reductions). Wang *et al* screened a total of 756 women in the first trimester of pregnancy for thyroid functions after Levothyroxine treatment and then follow up them until delivery. They found that therapy decreased the incidence of spontaneous abortions (24).

Conclusion

In conclusion, high anti-TPO antibody in women with RPL could increase risk of pregnancy loss, and treatment with levothyroxine helps them to have normal pregnancy.

Acknowledgments

Authors like to thank Yazd Reproductive Sciences Institute for financial support of this project, and thank Prof. Bashardoost for statistical analysis.

Conflict of interest

None of the authors have any conflict of interest in this project.

References

- Toth B, Jeschke U, Rogenhofer N, Scholz C, Würfel W, Thaler CJ, et al. Recurrent miscarriage: current concepts in diagnosis and treatment. J Reprod Immunol 2010; 85: 25-32.
- Carrington B, Sacks G, Regan L. Recurrent miscarriage: pathophysiology and outcome. Curr Opin Obstet Gynecol 2005; 17: 591-597.
- Monien S, Kadecki O, Baumgarten S, Salama A, Dörner T, Kiesewetter H. Use of heparin in women with early and late miscarriages with and without thrombophilia. Clin Appl Thromb Hemost 2009; 15: 636-644.
- Rai R, Regan L. Recurrent miscarriage. Lancet 2006; 368: 601-611.
- 5. Redline RW. Thrombophilia and placental pathology. *Clin Obstet Gynecol* 2006; 49: 885-894.
- Firouzabadi RD, Ghasemi N, Rozbahani MA, Tabibnejad N. Association of p53 polymorphism with

- ICSI/IVF failure and recurrent pregnancy loss. *Aust N Z J Obstet Gynaecol* 2009; 49: 216-219.
- Galofre JC, Davies TF. Autoimmune thyroid disease in pregnancy: a review. J Women's Health (Larchmt) 2009; 18: 1847-1856.
- 8. Poppe K, Glinoer D, Tournaye H, Devroey P, van Steirteghem A, Kaufman L, et al. Assisted reproduction and thyroid autoimmunity: an unfortunate combination? *J Clin Endocrinol Metab* 2003; 88: 4149-4152.
- Calvo RM, Jauniaux E, Gulbis B, Asunción M, Gervy C, Contempré B, et al. Fetal tissues are exposed to biologically relevant free thyroxine concentrations during early phases of development. *J Clin* Endocrinol Metab 2002; 87: 1768-1777.
- Stagnaro-Green A, Roman SH, Cobin RH, el-Hazary E, Alvarez-Marfany M, Davies TF. Detection of at-risk pregnancy by means of highly sensitive assays for thyroid autoantibodies. *JAMA* 1990; 264: 1422-1425.
- Lejeune B, Grun JP, de Nayer P, Servais G, Glinoer D. Antithyroid antibodies underlying thyroid abnormalities and pregnancy lossor pregnancy induced hypertension. *Br J Obstet Gynaecol* 1993; 100: 669-672.
- 12. Menken J, Trussell J, Larsen U. Age and infertility. *Science* 1986; 233: 1389-1394.
- Osmanagaoglu K, Tournaye H, Kolibianakis E, Camus M, van Steirteghem A, Devroey P. Cumulative delivery rates after ICSI in women aged >37 years. Hum Reprod 2002; 17: 940-944.
- Rashid M, Rashid MH. Obstetric management of thyroid disease. Obstet Gynecol Surv 2007; 62: 680-688.
- 15. Poppe K, Glinoer D. Thyroid autoimmunity and hypothyroidism before and during pregnancy. *Hum Reprod Update* 2003; 9: 149-161.

- Nilsson IM, Astedt B, Hender U, Bezerin D. Intrauterine death and circulating anticoagulant "antithromboplastin". Acta Med Scand 1975; 197: 153-159.
- Cowchock S, Smith JB, Gocial B. Antibodies to phospholipids and nuclear antigens in patients with repeated abortions. *Am J Obstet Gynecol* 1986; 155: 1002-1010.
- 18. Kaplan D. Fetal wastage in patients with rheumatoid arthritis. *J Rheumatol* 1986; 13: 875-877.
- 19. Challis JR, Lockwood CJ, Myatt L, Norman JE, Strauss JF 3rd, Petraglia F. Inflammation and pregnancy. *Reprod Sci* 2009; 16: 206-215.
- Lejeune B, Grun JP, de Nayer P, Servais G, Glinoer D. Anti-thyroid antibodies underlying thyroid abnormalities and miscarriage and pregnancy induced hypertension. *Br J Obstet Gynecol* 1993; 100: 669-672.
- 21. Negro R, Mangieri T, Coppola L, Presicce G, Casavola EC, Gismondi R, et al. Levothyroxine treatment in thyroid peroxidase antibody-positive women undergoing assisted reproduction technologies: a prospective study. *Hum Reprod* 2005; 20: 1529-1533.
- Roussev RG, Kaider BD, Pride DE, Coulam CB. Laboratory evaluation of women experiencing reproductive failure. Am J Reprod Immunol 1996; 35: 415-420.
- Poppe K, Glinoer D, Tournaye H, Devroey P, Schiettecatte J, Haentjens P, et al. Thyroid autoimmunity and female infertility. Verh K Acad Geneeskd Belg 2006; 68: 357-377.
- 24. Wang S, Teng WP, Li JX, Wang WW, Shan ZY. Effects of maternal subclinicalhypothyroidism on obstetrical outcomes during early pregnancy. *J Endocrinol Invest* 2012; 35: 322-325.