# Downloaded from ijrm.ir on 2025-11-18]

# A comparison of general health and coping strategies in fertile and infertile women in Yazd

Ali Reza Bakhshayesh Ph.D., Mahsa Kazeraninejad B.Sc., Mahsa Dehghan Mongabadi B.Sc., Malihe Raghebian B.Sc.

Department of Psychology and Education, Faculty of Humanities, Yazd University, Yazd, Iran.

### **Corresponding Author:**

Ali Reza Bakhshayesh, Department of Psychology and Education, Faculty of Humanities, Yazd University, Yazd, Iran. Email: abakhshayesh@yazduni.ac.ir Tel/Fax: (+98) 913 450 2254

Received: 4 January 2011 Revised: 18 April 2011 Accepted: 29 March 2012

### **Abstract**

**Background:** Infertility affects various aspects of personality and psychology, familial and career performances, and relationships. Studies show that stress, anxiety, life dissatisfaction, and other psychological problems follow infertility. Infertility issue, its tests and remedy are stressful and may lead to anxiety and depression and have destructive effects on couple relationships.

**Objective:** The present study was done in order to comparison general health and coping strategies in fertile and infertile women.

**Materials and Methods:** This is an analytic cross-sectional study and was done through random sampling on 70 fertile women and 70 infertile women who visited Yazd's clinics. The age range of participants was between 20-40 years. General Health Questionnaire (GHQ) and Ways of coping questionnaire (WOCQ) were filled by women who were agreed to participate in the study, following some explanations about aims and ways of doing the study. In the next step, data were analyzed through statistic methods and independent t-test. We considered a significant level p<0.05 in all tests.

**Results:** The results indicated a significant difference (p<0.05) with respect to general health in two groups, but no significant difference was found in problem-centered and emotion centered coping strategies and depression anxiety.

**Conclusion:** This study shows that general health in both groups is below average which means women are not sensitive about their general health. So planning on improving women's general health by providing consultation and training courses is suggested.

Key words: General health, Coping strategies, Fertility and infertility.

## Introduction

n today's modern world, the problem of infertility is increasing and becoming a social concern. Infertility was described as a global sanitation problem with physical, psychological, and emotional aspects in Bangkok International Conference in 1988 (1). Infertility and its remedies are major crises in one's life that can prolong several years. "This incident provokes tension in life and couples are subjected to an emotional attack" (2). As Stage mentioned (2007), the uncertainty associated with infertility is the key challenge for infertile women (3).

With respect to these processes, it can be implied that any good or bad changes in life requires a type of further compatibility. Strategies to encounter these changes in life and the resultant tension vary in different

individuals regarding different situations (4). Author's views in coping definition are arranged on a continuum with two opposite poles in two sides; in one hand it is believed that coping means how people overcome their problems (5) and in two other poles there is Folkman and Lazarus theory which defines coping stressful, regarding environmental conditions and environment management quality (6). "Coping strategies are collection of one's cognitive and behavioral efforts which are used to interpret, analyze, and reform a stressful condition, resulting in the reduction of its discomfort" (4).

Two main coping strategies are: emotional coping strategies, including efforts to set emotional consequences of stressful incident and keep the emotional and sentimental balance by controlling resultant emotions from stressful conditions; and problem centered

coping strategies including one's effective acts with respect to stressful conditions, and is trying to remove or change the source of stress (4). Dynamism is known as the common feature of problem-centered coping strategies. Dynamism provides requisite equipment's for active coping with stressful situations. This condition needs person's all potential abilities for opposite coping and solving the problem, and increases his success probability (7).

Another feature of people who use problem-centered coping strategies is the low level of tension. On the other side, denial and passivity are two characteristics of those who ineffective emotion-centered coping strategies. Denial of stressful situations and inability to use potential abilities and initiative will cause the occurred problem left unsolved, which leads to dissatisfaction of the person (7).

Moller *et al* have found that psychic pressures and worries resulting from infertility have a direct effect on physiological functions of body which ultimately leads to a negative effect on fertility (8). Omoaregba *et al* (2011), indicated that "the prevalence of probable psychological distress was significantly higher among the infertile group compared with their fertile counterparts (p<0.001)" (9).

Daniel et al (2010), have the same idea, and mention that, compared with fertile women, infertile women have higher odds of self-reported depression (10). Substantial evidences point that problems resulting from infertility and inappropriate coping strategies might be a factor which helps exacerbate infertility (11). However, Bringer et al (2010) provide evidence against the hypothesis, casual relationship between mental distress and infertility (12).

On the other side, those people who have peace of mind and general health, experience less psychic pressure, consequently their chance of fertility increase (11). Depression is one of the main psychological problems in infertile individuals, which has a noticeable impression on all aspects of their life (12). As Peterson *et al* (2010) have reported in their study, infertility stress which is linked with

depression and psychological distress, can lead to premature dropout from medical treatments and unresolved feelings of loss and grief (13).

Hariiyan, Mohammadpur, and Aghajanlu have found that 58% of 100 infertile women, to some extent, are suffering from depression (12). Litt *et al* reported that "25% of infertile women will get depression following IVF unsuccessful treatment "(14, 15). Infertile couples experience the resultant tension of depression individually and cooperatively. Such tension may lead to mental problems (16, 17).

Ramezanzadeh (2004) has shown that, in comparison to fertile women, infertile women experience more anxiety and depression (18). Aghanwa, Dare, and Oguniy in 1999, reported in their research that "7.29% of infertile women suffer from depression and anxiety disorder "(19). In fact, psychological factors not only can cause infertility, but also can lead to different psychological issues (18).

With respect to the fact that, not only Yazd province is one of the poles of infertility in Iran, and in Iranian culture, ability to become fertile has a great significant, but also, so far, no research was done with the subject of both coping strategies, and general health of fertile and infertile women.

Therefore this research was done to examine general health and coping strategies in fertile and infertile women who visited Yazd clinics, and tried to answer the following questions:

- 1. Is there a difference in emotional-centered coping strategies scale in fertile and infertile women?
- 2. Is there a difference in problem-centered coping strategies scale in fertile and infertile women?
- 3. Is there a difference in general health in fertile and infertile women?

# Materials and methods

The strategy for this study was analytic cross-sectional and required information was collected from fertile and infertile women by Ways of Coping Questionnaire (WOCQ) and

General Health Questionnaire (GHQ). Statistical society in the present research was all fertile and infertile women conferring to women's clinics of Yazd.

In order to choose sample, fertile and infertile women (in the age range of 20 to 40 years) who conferred to women's clinics were chosen by simple random sampling. There were 140 participants including 70 fertile and 70 infertile women, and 133 questionnaires were returned. Results were analyzed by SPSS statistics software and independent t-test.

### **Measurement tools**

General Health Questionnaire (GHQ): This test is designed by Goldberg and consists of 28 questions that evaluate 4 factors: physical, anxiety. social function disorder depression. So far, more than 70 researches have been done on validity of GHQ all over the world. Goldberg and Marry in 1987 (20) have reported that average sensibility of the questionnaire is 82% and the result of validity has shown that sensibility and feature of this questionnaire in the best cut-off score are 86.5% and 82% respectively, and the final coefficient with retest and Cronbach's alpha is 88%, which is in 99% trust level (21).

Ways of Coping Questionnaire (WOCQ): this is a 66 article test which is built with respect to Lazarus and Folkman's inventory, by themselves, and evaluates a wide range of thoughts and deeds that are done when the person is confronted to external and internal stressful situations. At first, participants were asked to describe recently experienced stressful situation, oral or written, and then, by reading the questionnaire, specify how he/she has used each of the following strategies; this test contains 7 subscales: 1) direct coping, 2) going away, 3) self-controlling, 4) social sensitivity demand, 5) responsibility, escape-avoidance, 7) solving and the problem.

# A. Problem centered strategies

Social support demand: describes efforts seeking informational and emotional support (questions: 8, 18, 22, 31, 42, 45).

Responsibility: accepting self-role in problem which is always accompanied by an effort to place everything in the right place (questions: 9, 25, 29, 51).

Thoughtfully solving the problem describes thoughtful problem-centered efforts for changing the situation which is followed by solving the problem (questions: 1, 26, 39, 48, 49, 52).

Double positive evaluation: efforts which adds positive meaning by focusing on personal growth. This scale has religious concept. (questions: 20, 23, 30, 36, 38, 56, 60)

## B. Emotion-centered strategies

Encounter coping: describes aggressive efforts done to change the situation and has some levels of hostility and risk-taking (questions: 6, 7, 17, 28, 34, 46).

Avoidance: cognitive efforts to make yourself alone and minimizing the importance of situation (questions: 44, 41, 21, 15, 13, 12).

Continence: describes efforts which set one's emotions and actions. (questions: 10, 14, 35, 43, 54, 62, 63).

Escape: describes dream thoughts and cognitive efforts in order to avoid the problem. This scale is different from avoidance scale (questions: 11, 16, 33, 40, 47, 50, 58, 59). Lazarus and Folkman (1988) reported internal stability =0.66-0.79 through Cronbach's Alpha for all coping strategies.

# Statistical analysis

In order to analyze information in this research, descriptive statistic (mean, standard deviation) and independent t-test were used.

# Results

This research evaluates emotion-centered strategies, problem-centered strategies and psychological health in fertile and infertile groups. The values of all variables are listed below (Table I, Figure 1).

Our findings showed that the two groups have no significant differences in using coping strategies. The mean of the resultant data in emotion-centered coping strategies in fertile women was 9.7±3.5, and in infertile women was 9.5±1.14. The mean of the resultant data in problem-centered coping strategies in fertile women was 9.4±2.19 and in infertile women was 9.7±1.65. The results of independent t-

test showed no significant differences between two groups in coping strategies (p=0.6). However, we can notice such difference in the case of their general health (p=0.004).

**Table I.** Mean, standard deviation, maximum and minimum score in coping strategies and psychological health and its derivative subtests in fertile and infertile women.

Variable	Emotion-centered strategies	Problem-centered strategies	Psychological health
Number of tests	133	133	133
Mean	9.6	9.6	29.9
Standard deviation	2.6	1.9	11.5
Maximum score	26	20.1	57
Minimum score	6.5	5.7	4




Figure 1: A comparison between mean in fertile and infertile groups.

As it can be seen in the above diagram, average scores with regard to emotion-centered and problem-centered coping strategies are in one line, however, general health in these two groups is in two different directions, and this shows the difference between these two groups with regard to general health.

### **Discussion**

Is there a difference in emotional-centered coping strategies scale in fertile and infertile women? This was the first question of this study. Results show that there is no difference in emotion-centered coping strategies in fertile and infertile women, which means that the level of using emotion-centered coping strategies in fertile and infertile women is almost the same. In this aspect, this study is

the same as Pahlavani *et al* research. In their study, 40 fertile (20 males, 20 females) and 40 infertile (20 males, 20 females) were compared.

They used SCL-90-R and Carver-Schier coping skills check list. Pahlavani *et al* (2002) got to this point that there is no difference in using emotion-centered coping strategies in fertile and infertile women (22). But the result is not the same in Shakeri's research (2006). In their study 150 infertile women conferring to

infertility clinics were studied and data were collected through GHQ-28 general health questionnaire and CRI coping responses. It shows that most of the infertile women use emotion-centered strategy (11).

Panagopoulou *et al* (2009) "do not support the hypotheses that emotional disclosure will reduce infertility-related psychological distress". In their study, "148 participants were randomized to an emotional-writing condition, a fact-writing condition and a control condition" (23).

The second question was: Is there a difference in problem-centered coping strategies scale in fertile and infertile women? Results show that there is no difference in using problem-centered coping strategies in fertile and infertile women. Despite the fact that we tried to find a study in this field, no study was found to compare the results.

Next question was: Is there a difference in general health in fertile and infertile women? Results show that there is a significant difference in fertile and infertile women with respect to general health.

General health in fertile women was in higher level than infertile women. In this aspect, our results were the same as Besharat et al in 2006 (24). In their research 45 fertile and 45 infertile women conferring to infertility clinics were studied and data were collected through the Golombok Rust Inventory of Marital State and Veit-Ware Mental health Inventory. They have got to the point that general health in fertile women stands in a higher level than infertile women.

Similarly, the same results were obtained by Bahrami *et al* (2007). In their study 150 fertile and 150 infertile couples were studied and Beck depression inventory and Larson sexual satisfaction inventory were used.

### Conclusion

This study shows that tension and worry effect general health in infertile women and reduce it. But there are no significant difference in anxiety and depression and using emotion centered and problem-centered coping strategies in fertile and infertile women.

## References

- Mohammadi Hosseini F, Farzadi L, Seyed Fatemi L. [The study of stressor factors and coping strategies of infertile women referring to Tabriz Infertility Educational and Therapeutic center in 1998]. Fertil Infertil J 1998; 2: 27-34. (in Persian)
- Hirxsch AM, Hirxsch SM. The longer psychological effects of infertility. J Obstet Gynecol Neonatal Nurs 1995; 24: 517-521.
- Stege RL. Women's experience of infertility. University of Alberta (Canada); 2007: 1-96.
- Ghazanfari F, kadampoor E. [Relationship between mental health and coping strategies in citizenship of Khoramabad city]. J Fundamentals Mental Health 2008; 37: 47-54. (in Persian)
- Sarason IG, Sara Son BR. Abnormal psychology. New Jersey. Prentice Hall; 1989: 1-108.
- Folkman S. personal control and stress and coping processes: a theoretical analysis. J Personal Social Psychology 1984; 46: 839-852.
- 7. Pearlinli Scholar C. The structure of coping. *J Health and Social Behavior* 1987; 2: 2-21.
- Omoaregba J, James B, Lawani A, Morakinyo O, Olotu O. Psychological characteristics of female infertility in a tertiary institution in Nigeria. *Annals of African Medicine* 2011; 10: 19-25.
- Herbert D, Luke JC, Dobson AJ. Depression: an emotional obstacle to seeking medical advice for infertility. J Fertil Steril 2010; 94: 1871-1821.
- Shaker J, Hosseini M, Golshani S, Sadeghi Kh, Feyzollahi V. [general health, coping strategies and marital satisfaction in fertile and infertile women]. Fertil Infertil Quarter 2006; 14: 75-269. (in Persian)
- 11. Haririan HR, Mohammadpur Y, Aghajanlu A. [Depression abundance in infertile women referring to reproduction clinics in Orumiye]. *Iran J Obstet Gynecol Infertil* 2009; 13: 45-49. (in Persian)
- Brennan D, Peterson, Georg H, Eifert. Using acceptance and commitment therapy to treat infertility stress. Chapman University. Cognitive and Behavioral Practice 2010; 18: 577-587.
- Beutel M, Kupfer J, Kirchneyer P, Kehde, Kolin FM, Schroder-Printzen I, et al. Treatment-related stresses and depression in couples undergoing assisted reproductive treatment by IVF or ICSI. *Andrologia* 1999; 31: 27-35.
- Rajvir D, Ceinwen E, Cumming DC. Psychological well-being and coping patterns in fertile men. Fertil Stril 2000; 74: 702-706.
- 15. Mcnaughton C, Bostwick M. Development of brief stress management support groups for couples undergoing in vitro fertilization treatment. *Fertil Steril* 2000: 5: 87-93.
- Strauss B, Hepp U. psychobiological characteristics of infertile couples: Can they predict pregnancy and treatment persistence? *J Comm Appl Soc Psychol* 1998; 2: 289-301.
- 17. Ramezanzadeh F, Aghssa MM, Abedinia N, Zayeri F, Khanafshar N, Shariat M, et al. A survey of relationship between anxiety depression and duration of infertility. BMC Women's Health 2004; 4 6-9
- Aghanwa HS, Dare FO, Ogunniyi SO. Socio demographic factors in mental disorders associated with infertility in Nigeria. J Psychosom Res 1999; 3: 117-123.

- 19. Goldberg DP. Manual of the general health questionnaire, Windsor: NFER Nelson. 1988.
- Omidiyan M. [the study of mental health and happiness among Yazd university students]. Studies in Education and Psychology 2009; 1: 102-116. (in Persian)
- 21. Pahlavani H, Shahrokh E, Malakuti K. [Studying factors causing stress, coping strategies and psychological health in infertile people]. *Iran J Psychiatry and Clinical Psychology* 2002; 7: 79-87. (in Persian)
- 22. Panagopoulou E, Montgomery A, Tarlatzis B.

- experimental, emotional disclosure in women undergoing infertility treatment: are drop outs better off? Social Science and Medicine 2009; 21: 678-681.
- Besharat MA, Hoseinzade R. [Comparison of general health and sexual problems of fertile and infertile women]. *Iran J Psychiatry and Clinical Psychology* 2006; 12: 146-153. (in Persian)
- 24. Bahrami N, Sattarzadeh N, Ranjbar Kuchaksorayi F, Ghojazadeh M. [The relationship between sexual satisfaction and depression among fertile and infertile couples]. Fertil Infertil Quarter 2007; 30: 52-59. (in Persian)