

The Chromosomal Abnormality of Failed Fertilized Human Oocytes in an In Vitro Fertilization Program

Seyed M. Kalnatar Ph.D.¹ Ahmad Ebrahimi B.S., Mehrdad Solimani, Hossein Fazli

Research & Clinical Centre for Infertility, Medical Sciences University, Yazd Iran.

Background: The high fertilization failure after IVF treatment cycles could be related to chromosomal abnormalities. This study was carried out to assess the frequency of chromosomal abnormality on human oocytes lacking signs of fertilization 18-20 h after insemination.

Materials and Methods: On day one, 18-20 h after insemination (IVF), fertilization was confirmed when two pronuclei (normal IVF) or more pronuclei (poly pronucleus FR) were present. Chromosomal analysis of unfertilized oocytes was carried out within 20-24 h of collection. All oocyte did not sign of pronuclei were collected from total fertilization failure, TFF (FR=0) or partial fertilization failure, PFF (FR=10-90%). Chromosomal preparation was carried out as described by Tarkowski's techniques. The average of finding between two groups was compared by X^2 test.

Results: Chromosome spreading permitted adequate analyzing in 348 unfertilized oocytes. In 33.6% chromosomal aneuploidy was observed with the following frequencies; hypo-hyploidy, 22/348 (6.4%), hyper-hyploidy, 42/348 (12.2%) and diploidy, 52/348 (15%). The frequency of aneuploidy was significantly higher in TFF group 33/80 (41%) than PFF group 83/268 (31%), $p < 0.01$, X^2 . The most frequent numerical aberration was observed in chromosome group, G of the human karyotyped.

Conclusion: Since cytogenetic analysis of failed fertilized oocytes and sperm function tests are very helpful for direct information on low success rate of fertilization, further studies analyzing on both gametes function in TFF cycles will be needed.

Key words: chromosome abnormality, cytogenetic, unfertilized oocytes, Total Fertilization Failure

Introduction

Assisted reproductive techniques (ART) have become an established treatment for certain types of human infertility, with high fertilization rate (Edirisinghe et al., 1997). With ARTs used for fertilization of human oocytes *in vitro* are extremely effective, the overall fertilization rates are now consistently about 60-80% in programs per inseminated egg (Bedford and Kim 1993). The fertilization failure may be occurred in some and/or occasionally all oocytes. In other words, in some women, it is common to find that some and occasionally all pre-ovulatory oocytes in a clutch are non-fertilized. TFF occurs in about 10-15% of patients (Barlow et al 1990). Whereas, partial failure in a subset of visually TFF occurs in about 10-15% of patients (Barlow et al. 1990). Whereas, partial failure in a subset of visually normal and mature oocytes from a clutch despite the presence of competent spermatozoa, is more common. There are several reasons for a proportion of oocytes to remain unfertilized after insemination, such as abnormality of sperm DNA, sperm inability to undergo

the acrosome reaction, immature oocyte either GV or MI stages and chromosomal abnormality as well (Kalantar et al., 1995; Plachot, 1995; Abruzzo & Hassold, 1995; Wall et al., 1996). Several researches have been attempted to analyze the chromosomal components reporting the incidence of chromosomal aberrations in the human oocyte to range from 11% to 49% (Bongso et al., 1988; Ma et al., 1989; Macas et al., 1990; Angell et al. 1991; Tarin et al., 1991; Kamiguchi et al., 1993; Almmedia & Bolton 1994; Sengoku, 1997).

This study was carried out on 115 patients underwent IVF treatment cycles who referred to our center for their infertility problem. Cytogenetic analysis was performed on 513 oocytes, which did not present signs of development 18-20 hr after insemination or re-insemination In Vitro.

Materials and Methods

Patients who were underwent ART at Yazd Research & Clinical Center for Infertility gave written consent to donate their excess oocytes and unfertilized oocytes for Cytogenetic analysis. The induction procedure was done as previously described (Aflatoonian et al., 2000). Oocytes were aspirated transvaginally with ultrasound guidance 35 h after the hCG administration. Oocytes

Corresponding Author:

Seyed M. Kalantar. Research & Clinical Centre for Infertility, Medical Sciences University, Yazd Iran.

E-mail: kalantarsm@ystp.ac.ir

Table I: The overall results of failed fertilize oocytes from TFF and Pff cycles

Variables	TFF	PFF	Total
All oocytes	85(16.6)	427(83.4)	512
No. of cycles	26(11.8)	198(88.2)	220
Analysed oocytes	80(23.0)	268(77.0)	348
Haploid (n=23)	47(20.3)	185(80.0)	232
Aneuploidy			
Hypo-haploid	05(22.7)	17(77.3)	22
Hyper-haploid	12(28.6)	30(71.4)	42
Diploid	16(3.8)	36(69.2)	52

* The numbers in parentheses are percentage

Table II: Chromosomal Complement of Unfertilized oocytes

Hypo-haploid	No.	Hyper-haploid	No.	Diploidy	No.
20,X-A,-C,-G	1	26,X,+G,+C,+D	1	46,XX	37
21,X,-D,-D	1	25,X,+c,+A	1	46,XY	15
21,X,-C,-E	1	24,X,+G	11		
21,-C,-X?	1	24,X,+X	1		
22,X,-C	1	24,X,+D	7		
22,X,-E	2	24,X,+C	8		
22,X,-G*	5	24,X,+mar	12		
22,-	6				
22,X,-A	2				
22,-?	2				
Tptal	22		42		52

were cultured in Ham's F10 medium supplemented with 10% FBS. Insemination was carried out 4-6 hr following oocyte recovery using 50,000-100,000 motile spermatozoa/ml which were separated from seminal plasma by Percoll preparation. On day one, 18-20 h after insemination (IVF), fertilization was confirmed when two pronuclei (normal IVF) or more pronuclei (poly pronucleus FR) were present. All embryos were graded on basis of 1-4 scales and transferred the best quality were selected and replacement to the uterus. Chromosomal analysis of unfertilized oocytes was carried out within 20-24 hr of collection.

Cytogenetic methods

Chromosomal preparation was carried out as described by Tarkowski's techniques (Tarkowski, 1966). Briefly, all oocyte with no sign of pronuclei were collected from TFF PFF. Prior to fixation for chromosomal analysis the cumulus cells were removed from the pre-ovulatory oocyte using 0.1% hyaluronidase (Sigma, Chemical Co., St Louis). The zona pelucida removal was done by mouth pipetting in and out. All oocytes were then exposed to a 1% sodium citrate

ashypotonic solution for 10 min. and fixed by a few drops of freshly prepared Carnoy's solution (3 parts ethanol: 1 part acid acetic glacial, Merck). The slides were dried on a warming plate, and then stained in 10% Gimsa solution (Merck), the metaphase cell examined under an oil immersion objectives lens (Nikon, Co.). The results were analyzed by X²-test, using SPSS for windows.

Results

In total, 512 unfertilized oocytes from 220 cycles with mean age 30.6±5.1, ranged 23-41 were prepared for further Cytogenetic investigation. Chromosomal analysis was possible on 348 unfertilized oocytes (68%). The number of oocyte per cycle ranged from 1 to 12. All oocytes remained unfertilized from 26 TFF cycles, and in 194 cycles ranged from 1 to 5 as PFF. The Cytogenetic results of unfertilized oocytes are shown in Table I.A total of 232 unfertilized oocytes (66.3%) had a normal haploid karyotype. 6.4% of oocytes had hypo-haploid karyotype, forty-two oocytes (12.2%) were




Figure 1. Metaphase plate of unfertilized oocytes after insemination in IVF program, prepared by Tarkowski method.

- a) typical metaphase plate with a diploid oocytes
- b) A typical hyperhaploidy (n>23 chromosome) oocyte

hyperhaploid, +1, to +3 (Figure 1.) and 52 were diploid (2n) (Figure 2). Of those only 5 cycles had more than one oocyte with abnormal chromosome, and remaining had only one abnormal oocyte per cycle. The most frequent numerical aberration was observed in chromosome groups, G, D, C, A of the human karyotyped. In cycles with TFF, 33 out of 80 oocytes showed aneuploidy which was significantly higher than cycles with PFF, where aneuploidy was 83 out of 268 oocytes ($p < 0.01$, X^2).

Discussion

The occurrence of unfertilized oocytes considered common in ART treatment cycles. The failed fertilization is caused by abnormal gametes either oocyte or sperm (Bedford et al., 1993). Regarding to MII oocytes and associated with spontaneous miscarriage, may all contribute to the final chromosomal abnormalities in the human abnormal embryos (Plachot, 1995; Abruzzo & Hassold, 1995; Wall et al., 1996; Kamiguchi et al., 1993; Almedia & Bolton 1994; Sengoku et al., 1997). The present data on cytogenetic analysis of unfertilized oocytes is compared with the average finding of the previous studies presented in Table 3 (32.75, ranged 18–73.8). The 33.3% total abnormality rate of oocytes failed to cleave 44–66 h after insemination, compared favorably with results from several studies (Bongso., 1988; Macas et al., 1990; Pellestor., 1991; Kamiguchi et al., 1993; Lim et al., 1995; Edirisinghe et al., 1997), but not with data from others (Ma et al., 1989; Delhanty & Penketh, 1990; Tarin et al., 1991; Almeida & Bolton, 1994; Nakaoka et al., 1998; Martini., et al., 1997). The difference of the chromosomal aberration between different studies may be explained by several factors such as, cause of infertility, age of women, method of culturing of oocytes, stimulation protocol, also method of oocytes preparation for chromosomal analysis as well (Edirisinghe et al., 1992; Lim et al., 1995; Sengoku et al., 1997; Edirisinghe et al., 1997; Nakaoka et al., 1998). It is reported that with the method described by Tarkowski an artificial loss of chromosome might occur while using the gradual fixation/air-drying method leads to poor spreading of chromosomes from oocytes that cause difficulty for analyzing in detail (Dyban et al., 1993; Nakaoka et al., 1998). In our study, the success rate of chromosome analysis was 68% which is in more than the gradual fixation/air-drying method used by others (Edirisinghe et al., 1997; Nakaoka et al., 1998). The incidence of chromosome aneuploidy in our study was not significantly different from previously reported as 29.65%, ranged 25%–37.3% (Pellestor et al., 1991; Tarin et al., 1991; Lim et al., 1995; Edirisinghe et al., 1997; Smith et al., 1998) which obtained using both methods. As shown in Table 2, aneuploidy in unfertilized oocytes was observed in all groups except B and F. the frequency of aneuploidy in group G ($n=20$, 32.2%) was higher than that in the other groups, which the results consistent with the data reported by Zenzens et al., 1992; Kamiguchi et al., 1993; Nakaoka et al., 1998, while we found only one

oocytes with chromosomal anomaly in E group, and non in B and F, groups.

We found hypohaploidy in 6.3% of the all analysed oocytes, which is similar to the value of 5.9% reported by Kamiguchi et al., 1993 while hyperhaploidy was 12% which was much higher than their value of 4.3% but similar to others report by (Ma et al., 1989; Macas et al., 1990; Angell et al., 1991; Almeida & Bolton., 1994). In the recent study, the overall incidence of diploidy was high (14.9%) which was similar of finding by Almeida & Bolton., 1994. Our results showed that the incidence of chromosome abnormality in unfertilized oocytes in TFF cycles was significantly higher than oocytes failed to fertilize in PFF. In the case of TFF cycles the semen parameters according to the WHO, 1998 were normal and did not impair the sperm quality. It is interesting to note that in nine cycles in a repeated IVF cycles, arrest of the oocytes was already observed and in one of those cycles sperm with abnormal DNA was found and no chromosomal abnormality in those cycles detected (Kalantar et al., 1998). To our knowledge, the present study represents the analysis of unfertilized oocytes obtained in TFF vs PFF. In conclusion, in a cytogenetic study carried out in patients groups with similar stimulation regimens, culture condition, treatment cycle, oocytes preparation, we were able to show that total fertilization failure may be affected by chromosomal aneuploidy. Like all previous studies, there are limitations to the approach used in this study. For instance using FISH technique to analysis the chromosomal abnormality, which would helpful to identify anomalies in detail. Also, we did not performed sperm function tests to find any sperm abnormality especially in TFF cycles to look the possible reasons of sperm and egg dysfunction should be caused of fertilization failure. Since cytogenetic analysis of failed fertilized oocytes and sperm function tests are very helpful for direct information on low success rate of fertilization, further studies analyzing on both gametes function in TFF cycles will be needed.

Acknowledgment

The authors wish to thank IVF team of the centre for providing unfertilized oocytes.

References

- Almeida PA., and Bolton VN (1994) The relationship between chromosomal abnormalities in the human oocytes and fertilization in vivo. *Hum Reprod* 9, 343–346.
- Abruzzo, M.A. and Hassold, T.A. (1995) Etiology of nondisjunction in humans. *Environ. Mol. Mutagen.*, 25 (Suppl. 26), 38–47.
- Aflatoonian., Karimzadeh., Dehghani., Kalantar SM., et al. 2000. The role of aspiration of half ovarian follicles prior to administration of hCG or GnRH-a for prevention of severe OHSS in ART programs. *Midd. East. Ferti. Soci.J.* 5.1; 73-75

- Angell, R.R., Ledger, W., Yong, E.L. et al. (1991) Cytogenetic analysis of unfertilized human oocytes. *Hum Reprod* 6, 568–573.
- Barlow, P., Englert, Y., Puissant, F. et al., (1990). Fertilization failure in IVF: why and what next? *Hum Reprod* 5:451-456.
- Bedford, J.M. and Kim, H.H. (1993) Sperm/egg binding patterns and oocyte cytology in retrospective analysis of fertilization failure in vitro. *Hum Reprod* 8, 453–463
- Bongso, A., Ng, S.C., Ratnam, S. et al. (1988) Chromosome analysis in human oocytes failing to fertilize after insemination in vitro. *Hum Reprod* 3, 645–649.
- Delhanty, D.A. and Penketh, R.J.A. (1990) Cytogenetic analysis of unfertilized oocytes retrieved after treatment with the LHRH analogue, buserelin. *Hum Reprod* 5, 699–702.
- Dyban, A., De Sutter, P. and Verlinsky, Y. (1993) Preimplantation cytogenetic analysis. In Verlinsky, Y. and Kuliev, A.M. (eds), *Preimplantation Diagnosis of Genetic Disease*. Wiley-Liss, New York, pp. 98.
- Edirisinghe, W.R., Murch, A.R. and Yovich, J.L. (1992) Cytogenetic analysis of human oocytes and embryos in an in-vitro fertilization programme. *Hum Reprod* 7, 230–236.
- Kalantar SM., Lenton EA., Barratt CLR (1995) Sperm function tests and fertilization failure following IVF. In *Human sperm acrosome reaction*, (Eds) Fenichel P & Parinaud J, Colloque INSERM/John Libbey Eurotext Ltd, 236, pp. 439-440.
- Kalantar SM., Lenton EA (2000) Evaluating the ability of biological substances for induction of acrosome reaction in normospermic samples. *Midd East Ferti. Soci J* 5.2;71-78.
- Kalantar SM., Lenton EA., Brewis I., et al (1998) Fertilization failure of IVF predictive power of SFT. *Andrologia.*, 30:4-5, 180-181.
- Kalantar SM., et al., Evaluation of a cut-off level for DNA normality of sperm to predict fertilization failure following IVF-ET. 5th Annual Meeting of MEFS. Amman Jordan, 1998.
- Kamiguchi, Y., Rosenbusch, B., Sterzik, K. and Mikamo, K. (1993) Chromosome analysis of unfertilized human oocytes prepared by a gradual fixation-air human oocytes. *Hum Reprod* 11, 2230–2238.
- World Health Organization (1999) *WHO Laboratory Manual for the Examination of Human Semen and Sperm-Cervical Mucus Interaction*. Cambridge drying method. *Hum Genet* 90, 533–541.
- Lim, A.S.T., Ho, A.T.N. and Tsakok, M.F.H. (1995) Chromosomes of oocytes failing in-vitro fertilization. *Hum Reprod* 10, 2570–2575.
- Ma, S., Kalousek DK., Zouves C., et al., (1989) Abnormal chromosome arrangements in human oocytes. *Hum Reprod* 5, 703-707.
- Macas E., Floersheim Y., Hotz E. et al., (1990) Chromosome analyzing of human oocytes failing to fertilize in vitro. *Fertil Steril* 51, 992-997.
- Mozdarani H & Aghdai F (2001) Cytogenetics analysis of failed-fertilized oocytes from Iranian infertile women after in vitro fertilization (IVF) and intracytoplasmic sperm injection (ICSI) procedures. *MEFSJ* 6.3:216-225.
- Nakaoka Y, Okamoto E, Miharu N, Ohama K. (1998), Chromosome analysis in human oocytes remaining unfertilized after in-vitro insemination: effect of maternal age and fertilization rate. *Hum Reprod* 13(2):419-24.
- Pellestor F (1991) Frequency and distribution of aneuploidy in human female gametes. *Hum Genet* 86, 283-288.
- Plachot, M. (1995) Oocyte — genetic aspects. In Grudzinskas, J.G. and Yovich, J.L. (eds), *Gametes — The Oocyte*. Cambridge University Press, Cambridge, pp. 95–107.
- Sengoku K., Tamate K., Takuma N, Yoshida T., Goishi K and Ishikawa M (1997) The chromosomal normality of unfertilized oocytes from patients with polycystic ovarian syndrome *Human Reproduction* 12, 3 pp.474–477,
- Smith. R., Walker L., Cobo AC., et al., (1998) Contribution of chromosomal abnormalities to in vitro fertilization failures. *Rev Med Chil* 126(5):511-9.
- Tarkowski, A.K. (1966) An air-drying method for chromosome preparation from mouse eggs. *Cytogenetics*, 5, 394–400.
- Tarin, J.J., Gomez, E., Sampaio, M. et al. (1991) Cytogenetic analysis of human oocytes from fertile women. *Hum Reprod* 8, 1100–1103.
- Wall, M. B., Marks, K., Smith, T. A. et al. (1996) *Cytogenetic and fluorescent in-situ hybridization chromosomal studies on in-vitro fertilized and intracytoplasmic sperm injected 'failed-fertilized'* University Press, Cambridge, UK.
- Zenzes, M.T. and Casper, R.F. (1992) Cytogenetics of human oocytes, zygotes, and embryos after in vitro fertilization. *Hum Genet* 88, 367–37.

