Association of vitamin D receptor gene variants with polycystic ovary syndrome: A case control study

Touraj Mahmoudi^{1, 2} M.Sc., Keivan Majidzadeh-A² M.D., M.PH., Ph.D., Hamid Farahani³ Ph.D., Mojgan Mirakhorli¹ Ph.D., Reza Dabiri⁴ M.D., Hossein Nobakhi⁴ M.D., Asadollah Asadi⁵ Ph.D.

- 1. Gastroenterology and Liver Diseases Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.
- 2. Cancer Genetics Department, Breast Cancer Research Center (BCRC), ACECR, Tehran, Iran.
- 3. Department of Physiology, School of Medicine, Qom University of Medical Sciences, Qom, Iran.
- 4. Internal Medicine Department, Semnan University of Medical Sciences, Semnan, Iran.
- 5. Department of Biology, Faculty of Science, University of Mohaghegh Ardabili, Ardabil, Iran.

Corresponding Author:

Keivan Majidzadeh and Touraj Mahmoudi are co-corresponding authors.

Keivan Majidzadeh, Cancer Genetics Department, Breast Cancer Research Center (BCRC), ACECR, No. 146, South Gandi St, Vanak Square, Postal code: 1517964311, Tehran, Iran. Email: kmajidzadeh@razi.tums.ac.ir Tel: (+98) 9121050564

Touraj Mahmoudi, Department of Cancer, Gastroenterology and Liver Diseases Research Center, Shahid Beheshti University of Medical Sciences, Velenjak, Shahid Chamran Highway, Postal code: 1985711151. Tehran, Iran. **Email:** mahmouditouraj@gmail.com **Tel:** (+98) 9124148851

Received: 11 January 2015 Revised: 3 August 2015 Accepted: 23 August 2015

Introduction

olycystic ovary syndrome (PCOS), a highly complex and heterogeneous disorder with unclear etiology, is endocrine probably common the most disorder, which has а strong genetic component (1). The underlying causes of PCOS are not completely known. However, in addition to the menstrual disturbance and hyperandrogenism, PCOS patients demonstrate an increased prevalence of type 2 diabetes mellitus, impaired glucose tolerance, hyperinsulinemia, insulin resistance, and obesity (2-4). On the other hand, our previous study and other studies have demonstrated that insulin resistance and

Abstract

Background: Vitamin D and insulin play an important role in susceptibility to polycystic ovary syndrome (PCOS), and therefore vitamin D receptor (*VDR*), parathyroid hormone (*PTH*), and insulin receptor (*INSR*) gene variants might be involved in the pathogenesis of PCOS.

Objective: The present study was designed to investigate the possible associations between polymorphisms in *VDR*, *PTH*, and *INSR* genes and the risk of PCOS.

Materials and Methods: *VDR*, *PTH*, and *INSR* gene variants were genotyped in 35 women with PCOS and 35 controls using Polymerase chain reaction – Restriction fragment length polymorphism method. Furthermore, serum levels of glucose and insulin were measured in all participants.

Results: No significant differences were observed for the *VDR FokI*, *VDR Tru9I*, *VDR TaqI*, *PTH DraII*, *INSR NsiI*, and *INSR PmlI* gene polymorphisms between the women with PCOS and controls. However, after adjustment for confounding factors, the *VDR BsmI* "Bb" genotype and the *VDR ApaI* "Aa" genotype were significantly under transmitted to the patients (p=0.016; OR= 0.250; 95% CI= 0.081-0.769, and p=0.017; OR= 0.260; 95% CI= 0.086-0.788, respectively). Furthermore, in the women with PCOS, insulin levels were lower in the participants with the *INSR NsiI* "NN" genotype compared with those with the "Nn + nn" genotypes (P= 0.045).

Conclusion: The results showed an association between the *VDR* gene *BsmI* and *ApaI* polymorphisms and PCOS risk. These data also indicated that the *INSR* "NN" genotype was a marker of decreased insulin in women with PCOS. Our findings, however, do not lend support to the hypothesis that *PTH* gene *DraII* variant plays a role in susceptibility to PCOS.

Key words: Insulin, Insulin receptor, Polycystic ovary syndrome, Vitamin D receptor.

obesity have a negative effect on serum levels of 25-hydroxyvitamin D [25 (OH) D] and a positive effect on parathyroid hormone (PTH) concentrations in the women with PCOS (5-7). Furthermore, these studies have also shown higher serum levels of 25 (OH) D, phosphorous, and PTH in women with PCOS than do controls (5, 7).

The pleiotropic biological actions of 1, 25dihydroxyvitamin D [1, 25 (OH) 2D] are mediated by the vitamin D receptor (VDR) through its ability to modulate the expression of target genes. The *VDR* gene polymorphisms have been shown to be associated with obesity, insulin resistance, serum levels of testosterone, Luteinizing hormone, PTH, and 25 (OH) D (8-12).

Furthermore, in our previous study а significant association was found between VDR gene Apal polymorphism and risk of PCOS (13). In addition, the previous studies have demonstrated significant associations between PTH gene polymorphisms and serum levels of PTH and calcium (14). Finally, in recent years, the associations between various INSR genetic variants and PCOS have been examined in several epidemiologic studies, and the results were contradictory. Some investigations have shown significant associations between INSR gene SNPs and the risk of PCOS (15). In contrast, other studies have found no association (16, 17).

Based on these considerations, the aim of the present study was to investigate whether the *Fokl, Bsml, Apal, Taql,* and *Tru9I* polymorphisms of the *VDR* gene, the *DralI* polymorphism of the *PTH* gene, and the *PmII* and *NsiI* polymorphisms of the *INSR* gene were associated with PCOS risk. Furthermore, the possible relationships of the gene variants with insulin resistance and serum levels of insulin were also evaluated.

Materials and methods

Participants

This case control study was conducted in the Gastroenterology and Liver Diseases Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran between Julv 2011 and September 2012. All participants were provided informed consent to the study and the Ethics Committee of Gastroenterology and Liver Diseases Research Center reviewed and approved this study.

35 women with PCOS as the case group and 35 healthy women as controls were selected. The diagnosis of PCOS was made according to (a) the presence of menstrual dysfunction i.e. oligomenorrhea (fewer than six menstrual periods in the preceding year) or amenorrhea (absence of periods for more than 6 months), and (b) clinical hyperandrogenism (i.e. hirsutism: Ferriman-(18) score>6) and/or Gallwev hyperandrogenemia, and (c) the exclusion of other related disorders, including nonclassic congenital adrenal hyperplasia, androgen secreting tumors, Cushing's syndrome and hyperprolactinaemia. These diagnostic criteria were in accordance with the National Institute of Child Health and Human Development criteria (19). The controls had normal menstrual cycles and none of them had clinical evidence of hyperandrogenism. All 70 women were Iranian and genetically unrelated.

Biochemical measurements

Blood sampling was performed between days 2 and 6 of a menstrual cycle in the control subjects or during a spontaneous bleeding episode or progestin-induced menstrual cycle in the PCOS patients. Blood samples were collected after an overnight fast and were immediately centrifuged, and sera were separated and frozen at -20°C until assayed. In all of the 70 participants serum concentrations of glucose and insulin were measured.

The homeostasis model assessment of resistance index (HoMA-IR) was insulin calculated according to the formula: [fasting insulin (µIU/mI) × fasting glucose (mg/dl)] /405 (20). The serum concentration of insulin was determined by radioimmunoassay (RIA) method (DRG instruments GmbH, Marburg, Germany). Serum glucose measurements were performed using standard methods (CinnaGen Inc., Tehran, Iran). The intra- and inter assay coefficients of variation were 2.2 and 6.5% for insulin, respectively.

Genotype analysis

Blood samples for molecular genetic studies were collected in tubes containing EDTA as an anticoagulant and store at 4C. Genomic DNA was purified from whole blood using a commercial isolation kit (BioNEER, Daejeon, Korea) according to manufacturer's instructions and was amplified by PCR. Genotyping was carried out by RFLP. Details of the studied SNPs, PCR Primers, PCR condition, RFLP condition, the length of PCR products and RFLP products are shown in Table I.

In this study, the SNPs studied were chosen based on their common use in previous genetic epidemiology studies and degree of heterozygosity.

The PCR products were digested overnight by corresponding restriction enzymes (Fermentas, Leon-Rot, Germany) and the RFLP products were run on 2 to 3% agarose gels, and stained with ethidium bromide for visualization under UV light. Genotypes were expressed in RFLP nomenclature: upper case letters denote the absence of a restriction site, while lower case letters indicated the presence of a restriction site. In other words, *VDR, PTH*, and *INSR* genotypes of each participants were identified according to the digestion pattern and alleles according to the presence ("f" or "b" or "a" or "r" or "t" or "d" or "n" or "p") or absence ("F" or "B" or "A" or "R" or "T" or "D" or "N" or "P") of the *Fokl, Bsml, Apal, Tru9l, Taql, Drall, Nsil*, and *Pmll* sites, respectively. The "F", "B", "A", "R", "T", "D", "N", and "P" alleles correspond to "C", "A", "A", "G", "T", "A", "G", and "T" nucleotides, respectively. Approximately 10% of the samples were randomly selected and genotyped in duplicate for confirmation and the results were 100% concordant.

Table I. Information for the studied markers in the VDR, PTH, and INSR genes

Gene (SNP)	SNP reference number	Location (Base change)	Forward primer Reverse primer	PCR condition	PCR fragment size (bp)	Restriction enzyme, Incubation temperature	RFLP fragments size (bp)
VDR (FokI)	rs10735810	Exon 2 (C/T)	5'-AGCTGGCCCTGGCACTGACTCTGCTCT-3' 5'-ATGGAAACACCTTGCTTCTTCTCCCTC-3'	35 cycles 93C 45s, 66C 30s, 72C 45s	265	FoKI, 55C	169, 96
VDR (BsmI)	rs1544410	Intron 8 (A/G)	5'-GGCAACCTGAAGGGAGACGTA-3' 5'-CTCTTTGGACCTCATCACCGAC-3'	35 cycles 93C 45s, 66C 30s, 72C 45s	461	BsmI, 37C	258, 203
VDR (ApaI)	rs7975232	Intron 8 (A/C)	5'-CAGAGCATGGACAGGGAGCAAG-3' 5'-GCAACTCCTCATGGCTGAGGTCTCA-3'	35 cycles 93C 45s, 66C 30s, 72C 45s	740	Apal, 37C	530, 210
VDR (Tru9I)	rs757343	Intron 8 (A/G)	5'-AATACTCAGGCTCTGCTCTT-3' 5'-CATCTCCATTCCTTGAGCCT-3'	35 cycles 93C 45s, 56C 30s, 72C 45s	331	Tru9I, 65C	178, 153
VDR (TaqI)	rs731236	Exon 9 (C/T)	5'-CAGAGCATGGACAGGGAGCAAG-3' 5'- GCAACTCCTCATGGCTGAGGTCTCA-3'	35 cycles 93C 45s, 66C 30s, 72C 45s	740	TaqI, 65C	495,290, 245, 205
PTH (DraII)	rs6256	Exon 3 (A/C)	5'-CATTCTGTGTACTATAGTTTG-3' 5'-GAGCTTTGAATTAGCAGCATG-3'	35 cycles 93C 45s, 56C 30s, 72C 45s	600	DraII, 37C	420, 180
INSR (NsiI)	rs2059806	Exon 8 (A/G)	5'-CGGTCTTGTAAGGGTAACTG-3' 5'-GAATTCACATTCCCAAGACA-3'	35 cycles 93C 45s, 56C 30s, 72C 45s	324	NsiI, 37C	239, 85
INSR (PmlI)	rs1799817	Exon 17 (C/T)	5'-CCAAGGATGCTGTGTAGATAAG-3' 5'-TCAGGAAAGCCAGCCCATGTC-3'	35 cycles 93C 45s, 56C 30s, 72C 45s	317	PmlI, 37C	274, 43

VDR: Vitamin D receptor; PTH: Parathyroid hormone; INSR: Insulin receptor.

Statistical analysis

To examine the normality of distribution of continuous variables between the studied women in two groups, the Kolmogorov-Smirnov goodness-of-fit test was used and where necessary, log transformation was performed. Because the values of HoMA- IR and serum insulin and glucose were skewed, they were log transformed for the analysis, and their geometric mean (geometric standard error of the mean) was presented.

Comparisons of the distribution of the allele and genotype frequencies were performed using the Chi-square test (χ 2). The Chi-square

test was also used to examine whether genotype frequencies for each polymorphism satisfied the Hardy-Weinberg equilibrium. Differences between genotype groups in biochemical parameters were assessed by ANOVA or analysis of covariance when appropriate. The odds ratios (OR) given with the respective 95% confidence intervals (95% CI) were estimated in the alleles and genotypes with a statistically significant difference among study groups. Data were analyzed using Statistical Package for the Social Sciences, version 15.0, SPSS Inc, Chicago, Illinois, USA, and the difference was considered significant at P< 0.05.

Results

Table selected Ш presents the characteristics of cases (age range, 19-42 years) and controls (age range, 19-44 years). There were no significant differences between case and control groups in age, Body mass index (BMI), insulin resistance, and serum level of insulin. However, the cases had significantly lower serum levels of glucose compared with the controls (p=0.023) and the effect of PCOS on the variable remained significant after adjustment for age and BMI (p=0.019).

Table III shows the genotype and allele distributions for the *Fokl, Bsml, Apal, Tru9l*, and *Taql* polymorphisms of the *VDR* gene, the *Drall* polymorphism of the *PTH* gene, and the *Nsil* and *Pmll* polymorphisms of the *INSR* gene in women with PCOS and controls. None of the genotype frequency distributions deviated significantly from the Hardy-Weinberg equilibrium (Table III). No significant

differences were observed for the *Fokl, Tru9l, Taql, Drall, Nsil,* and *Pmll* polymorphisms in either genotype or allele frequencies between the cases with PCOS and controls. However, analysis of the Bsml and Apal polymorphisms revealed significant differences between women with PCOS and the control women (Table III).

After adjustment for age and BMI, the *VDR* "Bb" genotype and "Aa" genotype were significantly under transmitted to the patients (p=0.016; OR=0. 250; 95% CI= 0.081-0.769, and p=0.017; OR= 0.260; 95% CI= 0.086-0.788, respectively).

After adjustment for age and BMI in the controls, women carrying the VDR "AA" or "BB" genotype had lower serum glucose levels as compared with individuals in the "Aa + aa" (n= 8, n= 27, respectively, 120.64 ± 1.09 versus 91.74 ± 1.04, P= 0.007) or "Bb+bb" (n= 5, n= 30, respectively; 119.58 ± 1.10 versus 1.05, P= 0.037) genotypes. 94.43 ± Furthermore, in the controls, serum level of insulin and insulin resistance in the women with the INSR "pp" genotype compared with those with the "PP + Pp" genotypes were lower and the difference remained significant after adjustment for age and BMI (n= 18, n= 17, respectively; insulin 9.28 ± 1.33 versus 21.05 ± 1.23, p= 0.036; HoMA- IR 2.15 ± 1.35 versus 5.30 ± 1.26 , p= 0.041). In addition, in the women with PCOS, insulin levels was lower in the women with the INSR Nsil "NN" genotype compared with those with the "Nn + nn" genotypes and the difference remained significant after adjustment for age and BMI (n= 18, n= 17, respectively; 13.25 ± 1.13 versus 18.61 ± 1.14, p= 0.045).

TADIE II. CHINCAI AND DIOCHENNCAI VANADIES OF THE STUDIED PROUD	Table II.	Clinical a	and biochemical	variables of	the studied	groups
--	-----------	------------	-----------------	--------------	-------------	--------

	Control group (n=35)	Case group (n=35)	$\mathbf{P_1}^{\mathbf{b}}$	$\mathbf{P_2}^{c}$
Age (years)	30.29 (1.17)	28.43 (0.80)	0.196	-
BMI (kg/m ²)	26.30 (0.63)	26.98 (0.83)	0.513	-
$HoMA - IR^d$	3.33 (1.22)	3.33 (1.11)	0.999	0.848
Serum glucose (mg/dl) ^d	97.67 (1.04)	86.35 (1.03)	0.023	0.019
$Serum\ insulin^{e}(\mu IU/ml)^{d}$	13.81 (1.21)	15.63 (1.10)	0.558	0.687

^a Data are presented as mean \pm standared error (SE). Geometric mean (geometric standard error of the mean) is presented for HoMA– IR and serum glucose and insulin, ^bP₁ is the unadjusted p values (One-way ANOVA test), ^cP₂ is the adjusted p-values for age and body mass index (ANCOVA test), ^d Significance was tested on log-transformed values.

Table III. Association between genotypes and alleles of VDR, PTH, and INSR gene polymorphisms and risk of PCOS

Gene/SNP	Control group (n=35) ^a	Case group (n=35) ^a	P-value
Genotype and allele-wise comparisons			
VDR/FokI			
FF: Ff: ff	24 (68.6): 10 (28.6): 1 (2.8)	16 (45.7): 17 (48.6): 2 (5.7)	0.153
F: f	58 (82.9): 12 (17.1)	49 (70.0): 30 (21.0)	0.073
VDR/Bsml ^b			
BB: Bb: bb	5 (14.3): 23 (65.7): 7 (20.0)	10 (28.6): 12 (34.3): 13 (37.1)	0.031
B: b	33 (47.1): 37 (52.9)	32 (45.7): 38 (54.3)	0.865
VDR/Apa1 ^c			
AA: Aa: aa	8 (22.9): 21 (60.0): 6 (17.1)	15 (42.9): 11 (31.4) : 9 (25.7)	0.054
A: a	37 (52.9): 33 (47.1)	41 (58.6): 29 (41.4)	0.496
VDR/Tru9I			
RR: Rr: rr	27 (77.1): 8 (22.9): 0 (0.0)	28 (80.0): 6 (17.1): 1(2.9)	0.521
R: r	62 (88.6): 8 (11.4)	62 (88.6): 8 (11.4)	1.000
VDR/ TaqI			
TT: Tt: tt	15 (42.9): 16 (45.7): 4 (11.1)	15 (42.9): 14 (40.0): 6 (17.1)	0.766
T: t	46 (65.7): 24 (34.3)	44 (62.9): 26 (37.1)	0.724
PTH/Drall			
DD: Dd: dd	0 (0.0): 16 (45.7) : 19 (54.3)	1 (2.9): 9 (25.7): 25 (71.4)	0.151
D: d	19 (27.1): 51 (72.9)	11 (15.7): 59(84.3)	0.099
INSR/NsiI			
NN: Nn: nn	21 (60.0) : 12 (34.3) : 2 (5.7)	18 (51.4): 12 (34.3): 5 (14.3)	0.468
N: n	54 (77.1): 16 (22.9)	48 (68.6): 22 (31.4)	0.254
INSR/PmlI			
PP: Pp: pp	1 (2.9) : 16 (45.7) : 18 (51.4)	0 (0.0): 14 (40.0): 21 (60.0)	0.506
P: p	18 (25.7): 52 (74.3)	14 (20.0): 56 (80.0)	0.424

Data are presented as number (%), ^b chi-square test for genotype-wise comparisons: BB/Bb + bb (p= 0.152; OR= 2.400, 95% CI= 0.725-7.949), Bb/BB + bb (p= 0.010; OR= 0.272, 95% CI= 0.101-730), and bb/ BB + Bb (p= 0.117; OR= 2.364, 95% CI= 0.807-6.927) test models, respectively. ^c chi-square test for genotype-wise comparisons: AA/Aa+aa (p=0.079; OR=2.531, 95% CI=0.899-7.124), Aa/AA+aa (p=0.018; OR=0.306, 95% CI=0.114-0.817), and aa/AA+Aa (p= 0.079; OR= 1.673, 95% CI= 0.524-5.341) test models, respectively.

VDR: Vitamin D receptor; PTH: Parathyroid hormone; INSR: Insulin receptor.

Discussion

Based on our results, significant associations between *VDR* gene *Bsml* and *Apal* polymorphisms and risk of PCOS were found. The *Bsml* "Bb" or Apal "Aa" genotypes appeared to be a marker of decreased PCOS susceptibility.

Furthermore, in the women with PCOS the *INSR Nsil* "NN" genotype appeared to confer an increased risk for serum insulin level, which our knowledge has not been reported previously.

Also, the *INSR Pmll* "pp" genotype appeared to confer a decreased risk for serum insulin levels and insulin resistance in the controls. In addition, the *VDR Bsml* "BB" or *Apal* "AA" genotypes appeared to confer a

decreased risk for serum glucose levels in the control women.

In recent years, genes involved in insulin signaling pathway have been suggested as candidate genes for PCOS. Our finding is in line with the other studies showing no significant association between the *INSR* gene polymorphisms and the risk of PCOS (16, 17).

However, positive associations also have been reported (15). Inconsistent results may be due to false positive results, population differences in allele frequencies, differences in disease definition, genotyped markers, statistical methods, and the variation in environmental, particularly nutritional factors. Furthermore, the *Nsil* and *Pmll* RFLPs at exon8 and exon 17 are "synonymous"

polymorphisms meaning that they do not alter the amino acid sequence of the INSR gene and may only be associated with risk of PCOS populations through in some linkage disequilibrium with another functional variant. In the present study, a significant association between INSR gene Nsil polymorphism and serum levels of insulin was also observed that is in agreement with the previous reports (21); to our knowledge, ours is the first study that demonstrates the association in women with PCOS. Furthermore, the results are in line with the previous studies (22, 23) showing significant associations between the INSR *Pmll* polymorphism and serum levels of insulin and insulin resistance. However, the the mechanisms behind molecular genotype/phenotype associations were observed for the INSR gene polymorphisms remain unexplained.

In this study, significant associations between VDR Bsml and Apal gene polymorphisms and risk of PCOS were observed that are in agreement with our previous report (13), to our knowledge, the present study is the first to show that VDR gene Bsml polymorphism is associated with risk of PCOS. In our previous study, the Apal "Aa" genotype was a marker of decreased PCOS susceptibility, whereas "aa" genotype was associated with an increased risk for PCOS (13). Furthermore, in the present study, significant associations between these two SNPs and serum levels of glucose in the controls were observed that they were in agreement with the previous reports (24). The Bsml and Apal polymorphisms are located in intron 8 at the 3' end of the VDR gene. The 3' untranslated region of genes is known to be involved in the regulation of gene expression, especially through regulation of mRNA stability. Some studies have suggested that, the Bsml and Apal polymorphisms are associated with differences in VDR mRNA expression level and stability (25). Furthermore, Whitfield et al (26) have shown that the "b" allele of Bsml polymorphism appears to be more active than the "B" allele. On the other hand, in accordance with this study it has been reported that fasting plasma glucose and prevalence of glucose intolerance were significantly lower in individuals with Apal "AA" genotype compared with those with "aa" genotype, (24) and the effect of VDR Apal polymorphism on insulin secretion has

been reported (27). In addition, our previous study showed that the "aa" genotype and the "bb" genotype conferred a decreased risk of obesity in women with PCOS (13). Finally, insulin secretion is a calcium-dependent process and 1, 25 (OH) 2 D-VDR complex stimulates the expression of insulin receptor gene (28).

Accordingly, it is possible that VDR gene polymorphisms through affecting the insulin signaling pathway play a role in pathogenesis PCOS. However, because of these polymorphisms are largely nonfunctional, it seems that linkage disequilibrium with another unknown functional variant of the VDR gene is to assumed explain the associations observed.

Furthermore, it is possible that *VDR* gene polymorphisms through affecting PTH- vitamin D axis play a role in pathogenesis of PCOS. Consistent with the view, it has been shown the associations between the *VDR* gene polymorphisms and serum levels of PTH (11) and 25 (OH) D (12), and vitamin D-VDR complex inhibit the secretion and synthesis of PTH (29).

The present study has several limitations. A limitation of the present study is its sample size. Considering the small sample size, the genotype differences may be attributable strictly to chance. Another potential limitation is our lack of information on serum levels of vitamin D and PTH, which could modify the effects observed here. Accordingly, it could not completely rule out the possibility of chance findings. Nevertheless, the possibility of true finding should not be excluded.

In conclusion, *VDR* gene variants appeared to be markers of decreased PCOS susceptibility. Our findings also suggested that INSR gene variant could affect serum level of insulin in women with PCOS. However, further studies are needed to confirm the findings and clarify the biological mechanisms by which these polymorphisms influence the PCOS risk.

Acknowledgments

The authors thank all the patients and healthy blood donors for providing blood samples. This work was supported by a grant from Gastroenterology and Liver Diseases Research Center, Shahid Beheshti University of Medical Sciences.

Conflict of interest

The authors declared that they have no conflict of interest.

References

- 1. Azziz R, Woods KS, Reyna R, Key TJ, Knochenhauer ES, Yildiz BO. The prevalence and features of the polycystic ovary syndrome in an unselected population. *J Clin Endocrinol Metab* 2004; 89: 2745-2749.
- Moran LJ, Misso ML, Wild RA, Norman RJ. Impaired glucose tolerance, type 2 diabetes and metabolic syndrome in polycystic ovary syndrome: A systematic review and meta-analysis. *Hum Reprod Update* 2010; 16: 347-363.
- 3. Kurabayashi T, Yahata T, Quan J, Tanaka K. Association of polymorphisms in the beta2 and beta3 adrenoceptor gene with polycystic ovary syndrome. *J Reprod Med* 2006; 51: 389-393.
- Cankaya S, Demir B, Aksakal SE, Dilbaz B, Demirtas C, Goktolga U. Insulin resistance and its relationship with high molecular weight adiponectin in adolescents with polycystic ovary syndrome and a maternal history of polycystic ovary syndrome. *Fertil Steril* 2014; 102: 826-830.
- Panidis D, Balaris C, Farmakiotis D, Rousso D, Kourtis A, Balaris V, et al. Serum parathyroid hormone concentrations are increased in women with polycystic ovary syndrome. *Clin Chem* 2005; 51: 1691-1697.
- Hahn S, Haselhorst U, Tan S, Quadbeck B, Schmidt M, Roesler S, et al. Low serum 25-hydroxyvitamin D concentrations are associated with insulin resistance and obesity in women with polycystic ovary syndrome. *Exp Clin Endocr Diab* 2006; 114: 577-583.
- Mahmoudi T, Gourabi H, Ashrafi M, Yazdi RS, Ezabadi Z. Calciotropic hormones, insulin resistance and the polycystic vary syndrome. *Fertil Steril* 2010; 93: 1208-1214.
- Grundberg E, Brandstrom H, Ribom EL, Ljunggren O, Mallmin H, Kindmark A. Genetic variation in the human vitamin D receptor is associated with muscle strength, fat mass and body weight in Swedish women. *Eur J Endocrinol* 2004; 150: 323-328.
- Wehr E, Trummer O, Giuliani A, Gruber H, Pieber TR, Obermayer-Pietsch B. Vitamin D-associated polymorphisms are related to insulin resistance and vitamin D deficiency in polycystic ovary syndrome. *Eur J Endocrinol* 2011; 164: 741-749.
- 10. Ranjzad F, Mahban A, Irani Shemirani A, Mahmoudi T, Vahedi M, Nikzamir A, et al. Influence of gene variants related to calcium homeostasis on biochemical parameters of women with polycystic ovary syndrome. *J Assist Reprod Genet* 2011; 28: 225-232.
- 11. Zofkova II, Zajickova K, Hill M. Serum parathyroid hormone levels are associated with Fokl polymorphism of the vitamin D receptor gene in untreated postmenopausal women. *Eur J Intern Med* 2003; 14: 232-236.
- Lin MW, Tsai SJ, Chou PY, Huang MF, Sun HS, Wu MH. Vitamin D receptor 1a promotor -1521 G/C and -1012 A/G polymorphisms in polycystic ovary

syndrome. Taiwan J Obstet Gynecol 2012; 51: 565-571.

- Mahmoudi T. Genetic variation in the vitamin D receptor and polycystic ovary syndrome risk. *Fertil Steril* 2009; 92: 1381-1383.
- 14. Gohda T, Shou I, Fukui M, Funabiki K, Horikoshi S, Shirato I, et al. Parathyroid hormonr gene polymorphism and secondary hyperparathyroidism in hemodialysis patients. *Am J Kidney Dis* 2002; 39: 1255-1260.
- 15. Goodarzi MO, Louwers YV, Taylor KD, Jones MR, Cui J, Kwon S, et al. Replication of association of a novel insulin receptor gene polymorphism with polycystic ovary syndrome. *Fertil Steril* 2011; 95: 1736-1741.
- 16. Lee EJ, Yoo KJ, Kim SJ, Lee SH, Chan KY, Baek KH. Single nucleotide polymorphism in exon 17 of the insulin receptor gene is not associated with polycystic ovary syndrome in a Korean population. *Fertil Steril* 2006; 86: 380-384.
- 17. Ranjzad F, Mahmoudi T, Irani Shemirani A, Mahban A, Vahedi M, Ashrafi M, et al. A common variant in the adiponectin gene and polycystic ovary syndrome risk. *Mol Biol Rep* 2012; 39: 2313-2319.
- Ferriman D, Gallwey JD: Clinical assessment of body hair growth in women. J Clin Endocrinol Metab 1961; 21: 1440–1447.
- Zawadski JK, Dunaif A. Diagnostic criteria for polycystic ovary syndrome: towards a rational approach. In: Dunaif A, Givens JR, Haseltine FP, Meriiam GR, Ed., Polycystic ovary syndrome. In: Hershman SM, Series Ed., Current Issues in Endocrinology and Metabolism. Blackwell, Boston, MA, USA, 1992; 377-384.
- 20. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and β -cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985; 28: 412-419.
- Flores-Martínez SE, Islas-Andrade S, Machorro-Lazo MV. DNA polymorphism analysis of candidate genes for type 2 diabetes mellitus in a Mexican ethnic group. Ann Genet 2004; 47: 339-348.
- 22. Urbanek M, Legro RS, Driscoll D, Strauss JF, Dunaif A, Spielman RS. Searching for the polycystic ovary syndrome genes. *J Pediatr Endocrinol Metab* 2000; 13: 1311-1313.
- 23. Mukherjee S, Shaikh N, Khavale S, Shinde G, Meherji P, Shah N, et al. Genetic variation in exon 17 of INSR is associated with insulin resistance and hyperandrogenemia among lean Indian women with polycystic ovary syndrome. *Eur J Endocrinol* 2009; 160: 855-862.
- 24. Oh JY, Barrett-Connor E. Association between vitamin D receptor polymorphism and type 2 diabetes or metabolic syndrome in community-dwelling older adults: the Rancho Bernardo study. *Metabolism* 2002; 51: 356-359.
- 25. Fang Y, van Meurs JB, Alesio A, Jhamai M, Zhao H, Rivadeneira F, et al. Promoter and 3'-untranslatedregion haplotypes in the vitamin d receptor gene predispose to osteoporotic fracture: the rotterdam study. *Am J Hum Genet* 2005; 77: 807-823.
- 26. Whitfield GK, Remus LS, Jurutka PW, Zitzer H, Oza AK, Hope T, et al. Are human vitamin D receptor gene polymorphisms functionally significant? In:

Norman AW, Bouillon R, Thomasset M. Vitamin D Endocrine System: structural, Biological Genetic and Clinical Aspects. California, USA: University of California, Riverside; 2000; 817-824.

- 27. Apparao KB, Lovely LP, Gui Y, Lininger RA, Lessey BA. Elevated endometrial androgen receptor expression in women with polycystic ovarian syndrome. *Biol Reprod* 2002; 66: 297-304.
- Maestro B, Campion J, Davila N, Calle C. Stimulation by 1, 25-dihydroxyvitamin D3 of insulin receptor expression and insulin responsiveness for glucose transport in U-937 human promonocytic cells. *Endocr J* 2000; 47: 383-391.
 Darwish H, DeLuca HF. Vitamin D-regulated gene
- Darwish H, DeLuca HF. Vitamin D-regulated gene expression. *Crit Rev Eukaryot Gene Expr* 1993; 3: 89-116.