Evaluating the effect of insulin sensitizers metformin and pioglitazone alone and in combination on women with polycystic ovary syndrome: An RCT

Seyed Mojtaba Sohrevardi¹ Pharm. D., Fahime Nosouhi¹ Pharm.,D., Saeed Hossein Khalilzade² M.D., Parichehr Kafaie³ M.D., Mojgan Karimi-Zarchi^{4,5} M.D., Iman Halvaei⁶ Ph.D., Mehdi Mohsenzadeh⁷ Ph.D.

- 1. Faculty of Pharmacy, Shahid Sadoughi University of Medical Sciences and Health Services, Yazd, Iran.
- 2. Division of Endocrinology, Department of Internal Medicine, Shahid Sadoughi University of Medical Sciences and Health Services, Yazd, Iran.
- 3. Department of Dermatology, Shahid Sadoughi University of Medical Sciences and Health Services, Yazd, Iran.
- 4. Departments of Obstetrics and Gynecology, Shahid Sadoughi University of Medical Sciences and Health Services, Yazd, Iran.
- 5. Recurrent Abortion Research Center, Research and Clinical Center for Infertility, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- 6. Department of Anatomical Sciences, Faculty of Medical Sciences, Tarbiat Modares University, Tehran, Iran.
- 7. Research and Clinical Center for Infertility, Shahid Sadoughi University of Medical Sciences, Yazd. Iran.

Mojgan Karimi-zarchi and Fahime Nosouhi are equal corresponding authors.

Corresponding Author:

Mojgan Karimi-Zarchi, Shahid Sadoughi Hosppital, Ebne Sina Blvd., Yazd. Iran.

Tel: (+98) 35 38224705

Email: drkarimi2001@yahoo.com Fahime Nosouhi, Shahid Sadoughi University of Medical Sciences and Health Services, Shohadaye Gomnam Blvd., Prof. Hesabi Blvd., Yazd. Iran.

Tel: (+98) 9133543417 Email: F_nosouhi_pharmacy@yahoo.com

Received: 30 April 2016 Revised: 1 September 2016 Accepted: 10 October 2016

Abstract

Background: Insulin resistance and hyperinsulinemia may play a role in pathogenesis of PCOS. One of the common therapeutic methods is using insulinsensitizing drugs such as metformin and thiazolidinediones.

Objective: The purpose was to determine the effect of metformin and pioglitazone on clinical, hormonal and metabolic parameters in women with PCOS.

Materials and Methods: Eighty four women randomly received one of the following for 3 months: metformin (n=28) (500 mg three times a day), pioglitazone (30 mg daily) (n=28) and combination of both metformin and pioglitazone (n=28) (30 mg/day pioglitazone plus 500 mg metformin three times a day). Hormonal profile, fasting serum insulin, body weight, body mass index, menstrual status and waist to hip ratio were evaluated before and after treatment.

Results: Metformin and pioglitazone and combination therapy induced favorable changes in fasting serum insulin, HOMA-IR index, QUICKI, fasting glucose to insulin ratio in women with PCOS. Body weight, BMI, and waist to hip ratio increased significantly after treatment with pioglitazone but the data were similar after administration of metformin or combination therapy. Total testosterone level decreased significantly only after treatment with metformin. After 3 months in patients who received pioglitazone or combination therapy, menstrual cycles became regular in 71.4% and 73.9% respectively. While menstrual improvement happened only in 36.4% of the patients treated with metformin.

Conclusion: These findings suggest that insulin-sensitizing drugs induce beneficial effect in insulin resistance and menstrual cyclicity but only metformin ameliorated hyperandrogenemia in women with PCOS. Treatment with combination of metformin and pioglitazone did not show more benefit than monotherapy with each drug alone.

Keywords: Polycystic ovary syndrome, Metformin, Pioglitazone, Insulin resistance.

Registration ID in IRCT: IRCT201403101836N7

Introduction

olycystic ovary syndrome (PCOS) is one of the most common endocrinological disorders diagnosed by detection of at least two of the following criteria; polycystic ovaries on ultrasound, menstrual irregularities,

hyperandrogenism based on biochemical and/or clinical manifestation (1). Insulin resistance and hyperinsulinemia may play an important role in pathogenesis of PCOS and overweight in PCOS women (2, 3). It has been reported that insulin resistance and consequent hyperinsulinemia play a prominent role in hyperandrogenemia through

direct way, induction of androgen production by theca cells, and indirect ways like elevation luteinizing hormone (LH) secretion. decreasing insulin-like growth factor (IGF) reducing binding protein. and hepatic synthesis of sex hormone binding globulin (SHBG) and consequently free androgen level augmentation occurs (4-6). Hyperinsulinemia and insulin resistance in PCOS patients appears simulate aggravate or hyperandrogenism because of gonadotropin secretory disturbance (2).

It has been postulated that abnormalities in insulin action (both insulin resistance and hyperinsulinemia), contribute to important comorbidities including: glucose intolerance, type 2 diabetes mellitus (DM2), dyslipidemia, increased plasma triglycerides and decreased plasma high-density lipoprotein cholesterol (HDL-C), hypertension, and atherosclerosis (7, 8). Likewise, insulin resistance has adverse effect on fertility and pregnancy rate (9, 10). PCOS is also associated with increased risk of endometrial cancer (11).

Since there is a relationship between PCOS and insulin resistance, administration of drugs to ameliorate insulin sensitivity and ovarian activity is suggested to treat this syndrome (12). Various clinical studies have focused on two different types of insulinsensitizing drugs for treatment of women with PCOS. Metformin and thiazolidinediones (troglitazone, rosialitazone. pioglitazone) are standard medications for treatment of DM2 via decreasing insulin resistance, improving insulin sensitivity in peripheral tissues, reducing free androgen levels and facilitating normal regular menses and pregnancy (13-16).

Metformin inhibits hepatic glucose and production lipogenesis, and also increases peripheral glucose uptake and fatty acid oxidation Pioglitazone, as a peroxisome proliferatoractivated receptor y agonist, decreases insulin resistance in liver and peripheral tissue and has anti-arteriosclerotic and anti-inflammatory effects. Moreover. probably pioglitazone decreases cardiovascular risk factors (17-19). have Many investigations shown metformin reduces insulin resistance and endocrine parameters in women with PCOS and is considered the most used drug to treat PCOS patients (20-25).

Although these two groups of insulin sensitizers have been shown to be effective, but they do not entirely restitute normal reproductive phenotype. Many women with PCOS do not response optimally to insulin sensitization even at the highest doses. Metformin and TZDs are also used as a combination therapy due to their "add-on effect" on insulin resistance; therefore it seems that combination therapy is required to treat all features of PCOS.

So far, few investigations have performed on treatment of PCOS with pioglitazone and also few studies have compared these insulin sensitizers alone or in combination in patients with PCOS (26, 27). Until now, it has not been proved which group of insulin sensitizer drugs is more efficient and suitable for initiation of PCOS treatment. Another question is whether combination therapy with two different types of insulin-sensitizer drugs is more potent than a single agent therapy (27). Also the effects of pioglitazone and metformin alone and in combination in three groups have not been reported in Middle Eastern women with PCOS.

The main goal of the current study was to compare the clinical, hormonal and metabolic change after treatment with metformin and pioglitazone. The specific goals were to distinguish whether the combination of the pioglitazone and metformin might be more effective in improvement menstrual regularity, hyperandrogenemia and prevention of long-term health hazard than using each drug alone.

Materials and methods

Subjects

Eighty four women with PCOS, aged 18-40 years old, with irregular menses and infertility and/or clinical sign of hyperandrogenism (hirsutism and acne) were recruited from Department of Gynecology, Shahid Sadoughi Hospital Yazd, Iran. between April 2014 to May 2015. Written informed consent was obtained from all participants. The research protocol was approved by the Ethics Committee of the Shahid Sadoughi University of Medical Sciences.

All patients had spontaneous inception of puberty and normal sexual growth and did not take oral contraceptive pills or anti-androgens. Also none of patients had received any drugs affecting hormonal profile and lipoprotein or glucose metabolism, or appetite at least three month before the onset of study.

The diagnosis of PCOS was based on at least two of the three following abnormalities according to the 2003 Rotterdam Consensus Conference: 1) Oligomenorrhoea amenorrhea; Oligomenorrhea was defined as menstrual cycle with an interval of ≥35 days. Secondary amenorrhea was defined as the absence of mense for at least six months. Irregular menstrual cycles was defined as cycle with changes of the intermenstrual interval of more than ±4 days and a period of ≤35 days (28). Regular menstrual was defined as cycles with the minimum and maximum length change of less than four days and cycle length ranging from 21-35 days, Manifestation of hyperandrogenemia such as hirsutism (Ferriman-Gallwey Score ≥8), acne and/or serum total testosterone greater than Polycystic 70 ng/dL, 3) ovaries bγ ultrasonography; confirmed by presence of 12 or more subcapsular follicles measuring 2-9 mm in diameter at least in one ovary or increased ovarian volume (>10 cm³) (29-31).

Patients with thyroid dysfunction, hyperprolactinemia, Cushing's syndrome, any androgen-secreting tumors, DM plasma glucose >7 mmol/L), nonclassical 21hydroxylase deficiency, autoimmune disease, central nervous system disease, significant hypertension, past hysterectomy, abnormal liver or kidney functions or active liver disease (ALT >2.5 the upper limit of normal range), and known heart disease, pregnancy and lactation were excluded from the study. Lost to follow-up, diagnosis of pregnancy and elevated levels of serum transaminases were exclusion criteria during the study.

Study design

Eighty four PCOS women were randomly assigned to one of the following groups during three months: group 1 consisted 28 patient received 500 mg metformin (t.i.d) with meals, group 2 consisted 28 subject received 30 mg pioglitazone (daily) and group 3 consisted of 28 subjects, received combination of pioglitazone 30 mg (daily) and metformin 500 mg (t.i.d). To reduce gastrointestinal side effect of metformin in the first wk of therapy, the dose of metformin was increased stepwise, from 500 mg with the evening meal for two days to 500 mg twice a day for two

days and thereafter 500mg three times per day. Before initiation of the study, all patients were allocated to three different groups in sequences provided by computer program generating random number. All subjects received a sealed envelope including the number 1 or 2 or 3 which was corresponding to metformin, pioglitazone and combination therapy, respectively. In our study because commercially accessible pills were applied, there was no blinding after randomization, therefore, investigator and subjects could be aware of the actual treatment.

Before the study, cortisol levels for all patients tested to rule out Cushing's syndrome and its impact on irregular menstruation, at 8 am To evaluate the drags effect on BMI, height and weight, waist-to-hip ratio for all patients was measured before and after the study .All subjects were on unrestricted diet and were advised not to change their physical exercise patterns during the study period. During the study, monthly follow-up visits were performed for all patients to evaluate menstrual status, side effects of the drugs and patients compliance. Compliance monitored by the self-reporting of the patients. Menstrual period length and cyclicity were evaluated by a self-filled menstrual calendar.

All potentially fertile patients were advised to use barrier method of contraception during the study if pregnancy was not desired and they were cautiously instructed to stop using the drug by confirmation of pregnancy. At the beginning of the study and after three months of treatment, in the early follicular phase (day 3-5) of spontaneous bleeding or progestininduced withdrawal bleeding, venous blood samples were collected after a 10-12h overnight fast to measure luteinizing hormone (LH), follicle-stimulating hormone (FSH), fast glucose, fast insulin, dehydroepiandrosterone sulfate (DHEA-S), total testosterone (T), total cholesterol (TC), triglycerides (TG), highdensity lipoprotein (HDL) cholesterol, lowdensity lipoprotein (LDL) cholesterol, alanine aminotransferase (ALT), aminotransferase (AST) and complete blood count. Thyroid-stimulating hormone (TSH), prolactin (PRL), 17-hydroxyprogesterone (170HPG) and plasma cortisol evaluated in order to exclude other disorder with similar feature to PCOS.

Also height, weight, body mass index (BMI), waist to hip ratio, Ferriman-Gallwey

score, acne score and menstrual history were recorded for all participants. Acne was assessed in four scores; 0= no acne; 1= slight acne on the face only; 2= moderate acne on the face only; 3= severe acne on the face and back or chest (32). All examinations were performed by the same practitioner.

Women were instructed to record a menstrual diary by a self-filled menstrual calendar during the study period to evaluate menstrual period length and cyclicity. The waist and hip circumferences were measured with a soft tape measure pursuant to the World Health Organization criteria. To measure plasma cortisol, blood sample was obtained at 8 A.M. and it was in the normal range for all subjects. In all patients, congenital adrenal hyperplasia was excluded by measurement of morning follicular phase 17-hydroxyprogesterone greater than 2 ng/ml.

At the base line of study (0 week) and after 3 months of treatment, women of all groups underwent ultrasonography in order to measure the number and size of follicles and ovarian size; Volume measurements were made by using the formula for the volume of an ellipsoid: 0.523 x length xwidth x thickness (33). In this investigation, insulin resistance was evaluated by different methods consisted of the homeostasis model assessment (HOMA), the quantitative insulin sensitivity check index (QUICKI) and fasting glucose to fasting insulin ratio (34, 35). HOMA was calculated in the following way:

fasting serum insulin
$$\left(\mu \frac{u}{ml}\right) \times \text{fasting plasma glucose}\left(\frac{mmol}{1}\right)$$

The QUICKI was calculated using the formula:

$$\frac{1}{\left[\log(fasting\ serum\ insulin\ \left(\frac{\mu U}{ml}\right) + \log(fasting\ serum\ glucose\ \left(\frac{mg}{dl}\right)\right]}$$

Laboratory assay

Nonheparinized venous blood samples were centrifuged, and serum was stored at -20°C until analysis. Fasting serum insulin was determined using chemiluminescence immunoassay (Insulin AccuLite CLIA Kit, USA). Serum fasting glucose was determined using a glucose oxidase method and using BT-3000 PLUS analyzer, Italy. The intra- and inter-assay coefficients of variation were ≤1.74% and ≤1.19%, respectively.

Testosterone and DHEAS was measured using chemiluminescence immunoassay by

DiaSorin *LIAISON*® For Analyzer. testosterone the intraand inter-assay coefficients of variation were ≤5%% and ≤10%, respectively. LH, FSH and PRL were chemiluminescence determined by (AccuLite **CLIA** immunoassay Kit. Monobindlnc, Lake Forest, CA92630, USA). For LH the intra- and inter-assay coefficients variation were ≤7.9% and ≤9.5%, respectively. For PRL the intra- and interassav coefficients variation of were ≤4.3%and≤9.8%, respectively.

Cortisol was measured by radioimmunoassay (RIA kit) and TSH was measured using immunoradiometric assay (RIAKEY® TSH IRMA Tube) with DREAM GAMMA-10 analyzer and the intra- and interassay coefficients of variation were ≤5.1% and ≤8.21%, respectively. 170HP was measured by enzyme immunoassay (AccuBind ELISA kit, Monobindlnc, Lake Forest, CA92630, USA). The intra- and inter-assay coefficients of variation were ≤8.5 and ≤8%, respectively. Spectrophotometry was used to determination of HDL, TG, total cholesterol, ALT and AST by using BT-3000 PLUS analyzer, Italy. The intra- and inter-assay coefficients of variation of lipids profile were less than 5%.

Statistical analysis

All outcome measures were represented as mean±SD. In this study, Wilk-Shapiro test was performed to assess the normality of the distribution of variables. Student's two-tailed paired t-test or the Wilcoxon signed rank test were used to compare mean variables at baseline to mean values after treatment within groups. One way ANOVA and post-hoc Bonferroni, were carried out to compare means between three groups. Chi-square and Fisher's exact test were used to evaluate menstrual pattern. Acne score and hirsutism score data had skewed distribution, so Kruskal-Wallis test was used for comparison of differences between groups. These data were reported as median and quartile range.

Because there was significant difference at the baseline in fasting glucose between three groups, we used analysis of covariance (ANCOVA) for comparison fasting glucose before and after treatment. A two tailed p<0.05 was considered statistically significant. In this study the SPSS statistical software package (version 22.0 for Windows; SPSS, Inc. Chicago, IL) was used. In RCTs, it is

necessary to mention how the authors calculated the sample size.

Results

Baseline characteristics

In the metformin group, two subjects withdrew from the study because of severe gastrointestinal side effects. One woman was lost to follow up, and three became pregnant. In the pioglitazone group, seven women became pregnant and in the combination group two women were lost to follow up, two patients withdrew from the study because of intolerable gastric discomfort and one became The clinical, metabolic. pregnant. hormonal measures at the baseline evaluation did not show significant differences between women who became pregnant or were lost to follow-up (Figure 1).

Sixty six women (age 29.04±5.93 years old) with BMI of 27.80±3.89 (kg/m²) finished the trial and were included in final statistical analysis. Pioglitazone was well tolerated and no patients discontinued pioglitazone because of side-effects. During the study period, no raise in serum transaminases were detected. There were no significant differences at the baseline in metabolic, clinical and hormonal measures between the three groups (ANOVA test) but baseline fasting glucose was higher in the metformin group than in other groups (p=0.014) and consequently baseline HOMA-IR index was higher in metformin group (p=0.04) (Table I).

In metformin group, 11 (50%) and 5 (22.7%) subjects, in pioglitazone group, 8 (40%) and 6 (30%) and in combination group 12 (52.2%) and 8 (34.8%) were overweight and obese, respectively. All patients had appearing polycystic ovaries on ultrasonography. **Before** treatment, 11 patients were amenorrheic (4 in metformin group, 4 in pioglitazone, 3 in combination group); 29 women were oligomenorrheic (10 in metformin group, 8 in pioglitazone group, 11 in combination group); 14 women had irregular menses and 12 women had regular menstrual cycle.

All women were affected by clinical and/or biochemical hyperandrogenism. An F-G score ≥8 was detected in 56 patients, 86.3% in metformin group, 76.1% in pioglitazone group and 91.3% in combination group. Acne was

observed in 72.7% of patients and hyperandrogenemia (defined as total testosterone >70 ng/ml) in 36.3% of subjects. Fasting hyperinsulinemia (>16 U/ml) was detected in 30.3%.

Post-treatment characteristics Metabolic parameters

Weight were increased significantly only in pioglitazone group (1.5 kg, p<0.001). Likewise, BMI was significantly higher in the pioglitazone group (p<0.001) after 3-month treatment but BMI and weight remained metformin unchanged in group combination therapy. Waist to hip ratio was increased significantly (p=0.001)after treatment pioglitazone and remained unchanged after metformin treatment or combination therapy. None of the women had DM at the baseline.

Treatment with metformin resulted in significant decreases in fasting glucose concentration (p=0.012),fasting (p=0.003), and the HOMA-IR index (p=0.001). QUICKI (p=0.001) and fasting glucose to insulin ratio (p=0.008)was increased significantly after metformin treatment. The treatment with pioglitazone decreased fasting concentration (p=0.018),glucose insulin (p=0.015),and HOMA-IR (p=0.011). Pioglitazone treatment increased significantly QUICKI (p=0.007) and fasting glucose to insulin ratio (p=0.026).

After treatment with combination metformin and pioglitazone, fasting insulin (p=0.026) and HOMA-IR index (p=0.013) decreased significantly. QUICKI (p=0.009) and fasting glucose to insulin ratio (p=0.03) were significantly increased. Fasting glucose showed decreasing trend but the difference was not significant. There were no differences for fasting insulin levels, insulin sensitivity indices (HOMA, QUICKI and fasting glucose to insulin ratio) between three groups after treatment (p=0.144, p=0.084, p=0.33 and p=0.82, respectively). Serum concentrations of TG, TC, LDL and HDL cholesterol, were similar between groups at the baseline, within the normal ranges and remained unchanged after treatment.

Hormonal parameters

At the baseline, the hormonal parameters were similar in all groups. LH level as well as the ratio of LH to FSH were decreased

significantly (p=0.002) but there was no significant change in the FSH levels after treatment with metformin. Pioglitazone and combination therapy had similar effects on hormones, significant pituitary causing increases in the FSH level (p=0.041 and p=0.005, respectively) and no significant change in the LH level was observed. The ratio of LH to FSH significantly decreased in both pioglitazone and combination group (p=0.007 and p=0.003, respectively). There was a decrease in serum total testosterone (p=0.018) after treatment with metformin but no significant differences were observed after pioglitazone and combination therapy. After treatment, subject who received metformin or pioglitazone or combination therapy did not indicate significant change in final serum DHEAS.

Menstrual pattern

During three months of administration of metformin and pioglitazone, 11 pregnancies were occurred (3 pregnancies on metformin, 7 on pioglitazone and 1 on combination therapy). This suggests that these drugs, particularly pioglitazone, result in rapid induction of regular menstrual cycles and ovulation in a subset of these women. There was a striking amelioration in menstrual cyclicity among the patients who were received pioglitazone or combination therapy.

In patients with menstrual disturbance treated with pioglitazone and combination

therapy, menstrual cycles became regular in 71.4% and 73.9% respectively. While improvement happened in 36.4% of the patients treated with metformin. There was a significant variation between metformin group and combination therapy in regularity of menstrual cycles (p=0.031)

After three months of treatment pioglitazone group, 6 of 8 oligomenorrheic and 1 of 4 amenorrheic women achieved regular cycles, and 3 women with irregular menses attained regular cycles. Three women still had amenorrhea. In combination therapy, 9 of 11 oligomenorrheic and 1 of 3 amenorrheic subjects became eumenorrhiec, and 3 of 5 women with irregular menstrual achieved regular menses. Two remained in amenorrhea. In metformin group, 1 of 10 women who had oligomenorrhea and 3 of 5 irregular menses achieved regular cycles and 1 of 4 amenorreheic became oligomenorrheic. 11 women who were oligoor amenorrheic at baseline reported no improvement in menstrual pattern.

Hyperandrogenism clinical manifestations

In our study, the administration of metformin, pioglitazone and combination therapy resulted in a significant decrease in the acne score by 38.9% (p=0.002), 68.4% (p<0.001) and 76.1%9 (p=0.001) respectively. No significant decrease was observed in the hirsutism score during the course of the study in three groups.

Table I. Baseline characteristics of women with PCOS

Characteristics	Metformin group (N=22)	Pioglitazone group (N=21)	Combination group (N=23)	p-value	
Age	28.72 ± 6.31	27.52 ± 5.01	30.73 ± 6.15	0.19	
Weight (kg)	71.3 ± 11.2	70.6 ± 13.2	72.6 ± 9.4	0.82	
BMI (kg/m^2)	27.5 ± 3.6	27.2 ± 4.7	28.5 ± 3.2	0.54	
Waist-to-hip ratio	0.8 ± 0.03	0.8 ± 0.05	0.8 ± 0.04	0.17	
Fasting glucose (mmol/l)	5.4 ± 0.5	5 ± 0.2	5.1 ± 0.5	0.014	
Fasting insulin (µU/ml)	17.1 ± 10.8	12.4 ± 7.1	11.7 ± 7.5	0.08	
HOMA-IR	4.2 ± 2.7	2.8 ± 1.6	2.7 ± 1.8	0.04	
QUICKI	0.3 ± 0.03	0.3 ± 0.02	0.3 ± 0.03	0.05	
Fasting G/I ratio (mg/dL per µIU / mL)	7.8 ± 5.3	9.5 ± 4.7	10.6 ± 6.3	0.25	
LDL cholesterol (mg/dl)	131.8 ± 26.7	113.8 ± 54.8	126.9 ± 33.5	0.28	
HDL cholesterol (mg/dl)	51.1 ± 12.1	51.1 ± 12.5	52.2 ± 12.2	0.94	
Serum cholesterol (mg/dl)	215 ± 36.5	185.6 ± 51.6	196 ± 36.6	0.07	
Serum triglyceride (mg/dl)	140.3 ± 58.2	111.6 ± 76	131 ± 59.3	0.34	
DHEAS (mg/l)	1.5 ± 0.7	1.4 ± 0.7	1.4 ± 0.7	0.79	
Total testosterone (µg/l)	0.7 ± 0.2	0.7 ± 0.1	0.6 ± 0.2	0.58	
FSH (IU/L)	5.2 ± 1.7	4.7 ± 1.2	5.6 ± 2.2	0.34	
LH(IÙ/L)	7.9 ± 2.3	7.6 ± 2.9	6.8 ± 2.7	0.37	
LH/FSH	1.6 ± 0.6	1.6 ± 0.8	1.4 ± 0.7	0.48	

Note: Plus-minus values are means \pm SD. To convert values for fasting glucose to mmol/L, multiply by 0.05551.

BMI: Body Mass Index

QUICKI: Quantitative insulin check index DHEA-S: Dehydroepiandrosteronesulphate

LH: Luteinizing hormone $\mu U/ml$: micro Units per milliliter $\mu g/l$, microgram per liter

HOMA-IR, Homeostatic model assessment of insulin resistance

G/I: Fasting glucose to fasting insulin ratio

FSH: Follicle stimulating hormone

mmol/l: milimol per liter mg/dL: microgram per deciliter

Table II. Clinical characteristics and serum hormone concentrations in women with the polycystic ovary syndrome after

administration of insulin-sensitizing drugs for three months

	Metformin		Pioglitazone		Combination					
Variable	Pre	Post	p-value	Pre	Post	p-	Pre	Post	p-value	p- value
	treatment	treatment	_	treatment	treatment	value	treatment	treatment	_	varue
Weight(kg)	71.3±11.2	71±12.8	NS	70.6±13.2	72.1±13.1	< 0.001	72.6±9.4	71.5±9.8	NS	NS
BMI(kg/m ²)	27.5±3.6	27.4 ± 4.4	NS	27.2 ± 4.7	27.8 ± 4.7	< 0.001	28.5±3.2	28±3.4	NS	NS
Waist-to-hip ratio	0.83 ± 0.031	0.83 ± 0.036	NS	0.80 ± 0.055	0.81 ± 0.055	0.001	0.82 ± 0.045	0.82 ± 0.049	NS	NS
Fasting glucose (mmol/l)	5.4±0.5	5.1±0.3	0.012	5±0.2	4.8±0.2	0.018	5.1±0.5	5±0.4	NS	0.02
Fasting insulin (µU/ml)	17.1±10.8	10.3±5.6	0.003	12.4±7.1	8.6±4.9	0.015	11.7±7.5	7.3±4.3	0.026	NS
HOMA-IR	4.2 ± 2.7	2.3±1.2	0.001	2.8 ± 1.6	1.8±1	0.011	2.7 ± 1.8	1.6 ± 0.8	0.013	NS
QUICKI	0.31 ± 0.03	0.34 ± 0.03	0.001	0.33 ± 0.02	0.35 ± 0.03	0.007	0.33 ± 0.03	0.36 ± 0.03	0.009	NS
Fasting G/I ratio (mg/dL per µIU/mL)	7.8±5.3	14.6±12.4	0.008	9.5±4.7	13.9±8.9	0.026	10.6±6.3	15.7±7.5	0.03	NS
LDL cholesterol (mg/dl)	131.8±26.7	127±28	NS	113.8±54.8	110±52.6	NS	126.9±33.5	124±29.8	NS	NS
HDL cholesterol (mg/dl)	51.1±12.1	52.9±11	NS	51.1±12.5	52.8±12.5	NS	52.2±12.2	55.7±13.2	NS	NS
Serum cholesterol (mg/dl)	215±36.5	209.7±37.2	NS	185.6±51.6	177.7±53.3	NS	196±36.6	197±35.6	NS	NS
Serum triglyceride (mg/dl)	140.3±58.2	147±78.9	NS	111.6±76	100.9±61.8	NS	131±59.3	114.3±78.5	NS	NS
DHEAS (mg/l)	1.5 ± 0.7	1.4 ± 0.5	NS	1.4 ± 0.7	1.6 ± 0.6	NS	1.4 ± 0.7	1.5±0.8	NS	NS
Total testosterone (µg/l)	0.7±0.2	0.6 ± 0.18	0.018	0.7 ± 0.1	0.7±0.2	NS	0.6±0.2	0.6±0.1	NS	NS
FSH (IU/L)	5.2±1.7	5.5±1.4	NS	4.7 ± 1.2	5.6 ± 2.1	0.041	5.6 ± 2.2	7±2.2	0.005	0.01
LH(IU/L)	7.9 ± 2.3	6.3 ± 2.4	0.002	7.6 ± 2.9	6.3 ± 2.4	NS	6.8 ± 2.7	6.4 ± 2.2	NS	NS
LH/FSH	1.6 ± 0.6	1.1 ± 0.4	0.002	1.6 ± 0.8	1.1 ± 0.4	0.07	1.4 ± 0.7	1±0.5	0.003	NS

Note: Plus-minus values are means \pm SD. To convert values for fasting glucose to mmol/L, multiply by 0.05551.

BMI: Body Mass Index

QUICKI: Quantitative insulin check index

DHEA-S: Dehydroepiandrosterone sulphate

LH: Luteinizing hormone

μU/ml: micro Units per milliliter

μg/l: microgram per liter

HOMA-IR: Homeostatic model assessment of insulin resistance.

G/I: Fasting glucose to fasting insulin ratio

FSH: Follicle stimulating hormone.

mmol/l: milimol per liter

mg/dL: microgram per deciliter

NS; Non-significant




Figure 1. Chart of the study from recruitment to completion after three months of treatment. LTFU=Lost to follow-up WDFS=Withdrew from the study




Figure 2. Serum total testosterone in women with PCOS at baseline and after three months of treatment. *= p<0.05(± SD)

Discussion

To our knowledge, the clinical, hormonal and metabolic change after treatment with metformin and pioglitazone has not been investigated in Middle Eastern women. In this study, we evaluated the efficacy of a combined therapy with metformin and pioglitazone as compared to metformin or pioglitazone monotherapy on the clinical and endocrine-metabolic alterations of PCOS patients. Specifically, we hypothesized that combined treatment would be more potent in treatment of symptoms and biochemical abnormalities characteristic of PCOS. compared to single-agent therapy. Metformin therapy has been shown to have beneficial effects on ovarian function, clinical feature. hormonal and metabolic parameter in women with PCOS (23, 36-37). It was shown that pioglitazone ameliorates the hormonal and metabolic consequence of PCOS (38-40). This trial purposed to contrast and compare hormonal, metabolic and clinical alterations following treatment with metformin and pioglitazone alone and in combination in women with PCOS.

This study illustrated insulin-sensitizing drugs exerted multiple beneficial effects on PCOS women with or without resistance. investigation Our showed ameliorating in insulin resistance and hyperinsulinemia after administration pioglitazone and metformin for a period of three months in PCOS women. We observed decrease of 44.5%, 33.3% and 41% in the HOMA-IR index after three months treatment with metformin, pioglitazone and therapy, respectively, combination decreases of 39.7%, 30.6% and 37.2% in fasting insulin levels after administration of metformin, pioglitazone and combination therapy, respectively. These variations of improvement in insulin resistance and insulin levels can be ascribed to the effects of metformin or pioglitazone. With regard to the role of insulin resistance as a component of metabolic syndrome and its relation to cardiovascular disease, this decrease can be useful in these patients.

We found that treatment with pioglitazone improved insulin action and insulin resistance despite an increase in body weight, waist to hip ratio and unfavorable BMI, which is in agreement with some previous studies using

pioglitazone in PCOS women (41-43) or other TZDs in patients with DM2 and insulin resistance (44-45). Improvement in insulin sensitivity despite increase in body weight could be explained by the advantageous transference from visceral to subcutaneous fat and the concurrent amelioration insulin sensitivity of hepatic and peripheral tissue induced by TZDs (46-48). The reduction of lipid profile did not show significance difference probably because of the short duration of the study. Three women on metformin (10.7%), seven on pioglitazone (25%) and 3.5% on combination therapy became pregnant during the study.

In this study, it appears that insulin sensitizing drugs might improve menstrual regularity compared to the baseline nontreated state, although this result was consistent with previous reports in women with PCOS taking either pioglitazone or metformin (21, 32, 38, 49). We found combination therapy as well as administration of pioglitazone for three months is more effective than metformin alone in ameliorating menstrual cyclicity. This finding is in line with Glueck et al who evaluated the effect of pioglitazone plus metformin diet on women were non-optimally responsive who metformin diet alone (39). There was significant difference in body weight after treatment with pioglitazone and no significant change after combination therapy was seen; therefore it may be concluded that the improvement of menstrual cycles was not operated via the body weight loss.

Both in vitro and in vivo studies showed that hyperandrogenism in PCOS women might be a result of hyperinsulinemia from peripheral insulin resistance (48, 50, 51). In the present study, we observed that metformin compared to pioglitazone or combination therapy resulted in a significant decrease in the levels of total testosterone. Our findings were in contrast with Ortega-Gonzalez et al who assessed responses of serum androgen treatment with metformin pioglitazone in PCOS women (38). Also, our data were not in agreement with Legro et al who examined the effect of metformin and rosialitazone. member а thiazolidinedione family like pioglitazone, on ovarian function (52). Another study showed that metformin was more effective in reducing testosterone levels but rosiglitazone had a

better effect on decreasing serum insulin and insulin resistance (53).

The lack of change after treatment with pioglitazone in total testosterone levels in the present study could be elucidated in two ways: 1) more effect of metformin on the ovarian cells and preventing androgen production; 2) probably inadequate dose of pioglitazone was given. It is not clearly known whether the profitable effect of TZDs in women with PCOS is due to a direct effect on ovarian function or adrenal gland or is due to reduction in insulin resistance. However, according to the results of this study, it appears that the effect of these drugs in PCOS is due to improved insulin resistance.

In this study a higher pregnancy rate was occurred in the pioglitazone compared to the metformin group, nevertheless our sample size is small for any ultimate conclusions. It is necessary to do more studies on the safety of TZDs and metformin in women who may achieve pregnancy during treatment with these drugs. Metformin is classified as a pregnancy category B drug and seems to be safe during pregnancy. Nawaz et al reported that in PCOS women, the continued use of metformin during pregnancy decreases the rates of abortion, gestational diabetes and fetal growth limitation (54). Rosiglitazone and pioglitazone are classified as pregnancy category C drugs, so PCOS women treated these drugs should be contraceptive before treatment and cautiously instructed to stop using the drug instantly after confirmation of pregnancy (55, 56). However two reported cases of women who taking the rosiglitazone during pregnancy have shown no infants malformation (57, 58).

Unexpectedly, in contrast our to hypothesis, our data showed that combination therapy may have little advantages compared to treatment with each drug alone. Similar findings have been reported in women with PCOS treated with metformin and other TZDs. Baillargeon et al reported that combined administration of metformin and rosiglitazone do not illustrate noteworthy benefits over using alone in normal insulin sensitivity nonobese women with **PCOS** (27).Thev observed a considerable amelioration ovulatory frequencies on metformin in comparison with rosiglitazone, the combination was not more potent. After treatment unbound testosterone levels were

comparable between three groups, and there was also no significant change in area under insulin curve and fasting insulin levels after administration of rosiglitazone compared to but there was a noticeable placebo. improvement after metformin or combination therapy. Huang et al showed metformin can be amend health-related quality of life of women with polycystic ovary syndrome by meliorate psychological disorders due to acne, hair loss and infertility problems, especially for hyperandrogenic patients who are overweight (59). Ghaneei et al reported, the outcomes demonstrated that cabergoline can be utilized as a protected organization for PCOS patients with hyperprolactinemia to enhance menstrual cycles (60).

Legro et al reported no beneficial effects of combined rosiglitazone and metformin therapy over single-agent therapy. They showed more improvements with rosiglitazone compared to free testosterone metformin on levels. hyperinsulinemia, and trending toward elevated ovulation rate (52). Glueck et al reported that when pioglitazone was added to metformin therapy, in women with PCOS who failed to respond optimally to metformin, insulin resistance, fasting glucose, insulin level, and DHEAS significantly decreased and HDL cholesterol and sex hormone-binding globulin striking increased and menstrual cyclicity was improved (39). There are some pitfalls in our study. The primary flaws are the small sample size, short time of drug administration and restrictions to follow-up the subjects more than three months, so further large scale randomized trial with longer duration of drug use and longer follow-up period are required to assess effectiveness of the metformin and pioglitazone in women with PCOS.

summary, we found that insulinsensitizing drugs have favorable influence on menstrual cyclicity improvement and insulin resistance for women with PCOS. But effective metformin was more than pioglitazonein to improve hyperandrogenemia, serum insulin levels, and insulin-resistance; pioglitazone was more efficient in regulating the menstrual cycle and clinical sign of hyperandrogenism (acne) and the adjunct of pioglitazone to metformin may not have further benefit compared to treatment with using drugs alone. This study might recommend that pioglitazone is a better choice in less-insulin-resistant women with PCOS who are seeking fertility or improvement in clinical signs of hyperandrogenism and modest increase in testosterone levels while in cases with predominant hyperandrogenemia and insulin resistance, metformin should be applied.

Acknowledgements

This research did not receive any specific grant from any funding agency in the public, commercial or not-for-profit sector.

Conflicts of interest

The authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

References

- ESHRE TR, Group A-SPCW. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. Fertil Steril 2004: 81: 19-25.
- Dunaif A. Insulin Resistance and the Polycystic Ovary Syndrome: Mechanism and Implications for Pathogenesis 1. Endocrine Rev 1997; 18: 774-800.
- Stepto NK, Cassar S, Joham AE, Hutchison SK, Harrison CL, Goldstein RF, et al. Women with polycystic ovary syndrome have intrinsic insulin resistance on euglycaemic–hyperinsulaemic clamp. Hum Reprod 2013: des463.
- 4. Willis D, Mason H, Gilling-Smith C, Franks S. Modulation by insulin of follicle-stimulating hormone and luteinizing hormone actions in human granulosa cells of normal and polycystic ovaries. *J Clin Endocrinol Metab* 1996; 81: 302-309.
- Tsilchorozidou T, Overton C, Conway GS. The pathophysiology of polycystic ovary syndrome. *Clin Endocrinol* 2004; 60: 1-17.
- Nestler JE, Jakubowicz DJ, Falcon de Vargas A, Brik C, Quintero N, Medina F. Insulin Stimulates Testosterone Biosynthesis by Human Thecal Cells from Women with Polycystic Ovary Syndrome by Activating Its Own Receptor and Using Inositolglycan Mediators as the Signal Transduction System 1. J Clin Endocrinol Metab 1998; 83: 2001-2005.
- Ehrmann DA, Liljenquist DR, Kasza K, Azziz R, Legro RS, Ghazzi MN. Prevalence and predictors of the metabolic syndrome in women with polycystic ovary syndrome. J Clin Endocrinol Metab 2006; 91: 48-53.
- Coviello AD, Legro RS, Dunaif A. Adolescent girls with polycystic ovary syndrome have an increased risk of the metabolic syndrome associated with increasing androgen levels independent of obesity and insulin resistance. *J Clin Endocrinol Metab* 2006; 91: 492-497.

- Glueck C, Phillips H, Cameron D, Sieve-Smith L, Wang P. Continuing metformin throughout pregnancy in women with polycystic ovary syndrome appears to safely reduce first-trimester spontaneous abortion: a pilot study. Fertil Steril 2001; 75: 46-52.
- 10. Pasquali R, Gambineri A, Biscotti D, Vicennati V, Gagliardi L, Colitta D, et al. Effect of long-term treatment with metformin added to hypocaloric diet on body composition, fat distribution, and androgen and insulin levels in abdominally obese women with and without the polycystic ovary syndrome. *J Clin Endocrinol Metab* 2000; 85: 2767-2774.
- Barry JA, Azizia MM, Hardiman PJ. Risk of endometrial, ovarian and breast cancer in women with polycystic ovary syndrome: a systematic review and meta-analysis. *Hum Reprod Update* 2014; 20: 748-758.
- 12. Tang T, Norman RJ, Balen AH, Lord JM. Insulin-sensitising drugs (metformin, troglitazone, rosiglitazone, pioglitazone, D-chiro-inositol) for polycystic ovary syndrome. Cochrane Database Syst Rev 2003; CD003053
- Dunaif A. Drug insight: insulin-sensitizing drugs in the treatment of polycystic ovary syndrome-a reappraisal. Nat Rev Endocrinol 2008; 4: 272-283.
- Wang P, Streicher P, Glueck CJ. Treatment of polycystic ovary syndrome with insulin-lowering agents. Exp Opin Pharm 2002; 3: 1177-1189.
- Glueck CJ, Wang P, Goldenberg N, Sieve-Smith L. Pregnancy outcomes among women with polycystic ovary syndrome treated with metformin. *Hum Reprod* 2002; 17: 2858-2864.
- Badawy A, Elnashar A. Treatment options for polycystic ovary syndrome. Int J Women's Health 2011; 3: 25.
- 17. Stadtmauer LA, Wong BC, Oehninger S. Should patients with polycystic ovary syndrome be treated with metformin? Benefits of insulin sensitizing drugs in polycystic ovary syndrome-beyond ovulation induction. *Hum Reprod* 2002; 17: 3016-3026.
- Vitek W, Alur S, Hoeger KM. Off-label drug use in the treatment of polycystic ovary syndrome. Fertil Steril 2015; 103: 605-611.
- 19. Ishibashi M, Egashira K, Hiasa K-i, Inoue S, Ni W, Zhao Q, et al. Antiinflammatory and antiarteriosclerotic effects of pioglitazone. *Hypertension* 2002; 40: 687-693.
- 20. Velazquez E, Mendoza S, Hamer T, Sosa F, Glueck C. Metformin therapy in polycystic ovary syndrome reduces hyperinsulinemia, insulin resistance, hyperandrogenemia, and systolic blood pressure, while facilitating normal menses and pregnancy. *Metabolism* 1994; 43: 647-654.
- 21. Velázquez E, Acosta A, Mendoza SG. Menstrual cyclicity after metformin therapy in polycystic ovary syndrome. *Obstet Gynecol* 1997; 90: 392-395.
- 22. Diamanti-Kandarakis E, Kouli C, Tsianateli T, Bergiele A. Therapeutic effects of metformin on insulin resistance and hyperandrogenism in polycystic ovary syndrome. *Eur J Endocrinol* 1998; 138: 269-274.
- 23. Moghetti P, Castello R, Negri C, Tosi F, Perrone F, Caputo M, et al. Metformin Effects on Clinical Features, Endocrine and Metabolic Profiles, and Insulin Sensitivity in Polycystic Ovary Syndrome: A Randomized, Double-Blind, Placebo-Controlled 6-Month Trial, followed by Open, Long-Term Clinical

- Evaluation 1. J Clin Endocrinol Metab 2000; 85: 139-146
- 24. Harborne L, Fleming R, Lyall H, Norman J, Sattar N. Descriptive review of the evidence for the use of metformin in polycystic ovary syndrome. *Lancet* 2003; 361: 1894-1901.
- Lord JM, Flight IH, Norman RJ. Metformin in polycystic ovary syndrome: systematic review and meta-analysis. BMJ 2003; 327: 951.
- Stumvoll M, Häring H-U. Glitazones: clinical effects and molecular mechanisms. Ann Med 2002; 34: 217-224.
- 27. 27. Baillargeon J-P, Jakubowicz DJ, Iuorno MJ, Jakubowicz S, Nestler JE. Effects of metformin and rosiglitazone, alone and in combination, in nonobese women with polycystic ovary syndrome and normal indices of insulin sensitivity. Fertil Steril 2004; 82: 893-902.
- Homburg R. Polycystic ovary syndrome-from gynaecological curiosity to multisystem endocrinopathy. Hum Reprod 1996; 11: 29-39.
- Polson D, Wadsworth J, Adams J, Franks S. Polycystic ovaries-a common finding in normal women. *Lancet* 1988; 331: 870-872.
- Ferriman D, Gallwey J. Clinical assessment of body hair growth in women. J Clin Endocrinol Metab 1961; 21: 1440-1447.
- Balen AH, Laven JS, Tan SL, Dewailly D. Ultrasound assessment of the polycystic ovary: international consensus definitions. *Hum Reprod Update* 2003; 9: 505-514.
- Kolodziejczyk B, Duleba AJ, Spaczynski RZ, Pawelczyk L. Metformin therapy decreases hyperandrogenism and hyperinsulinemia in women with polycystic ovary syndrome. Fertil Steril 2000; 73: 1149-1154.
- 33. Robert Y, Dubrulle F, Gaillandre L, Ardaens Y, Thomas-Desrousseaux P, Lemaitre L, et al. Ultrasound assessment of ovarian stroma hypertrophy in hyperandrogenism and ovulation disorders: visual analysis versus computerized quantification. *Fertil Steril* 1995; 64: 307-312.
- 34. Matthews D, Hosker J, Rudenski A, Naylor B, Treacher D, Turner R. Homeostasis model assessment: insulin resistance and β-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* 1985; 28: 412-419.
- 35. 35. Katz A, Nambi SS, Mather K, Baron AD, Follmann DA, Sullivan G, et al. Quantitative insulin sensitivity check index: a simple, accurate method for assessing insulin sensitivity in humans. *J Clin Endocrinol Metab* 2000; 85: 2402-2410.
- 36. Fleming R, Hopkinson ZE, Wallace AM, Greer IA, Sattar N. Ovarian function and metabolic factors in women with oligomenorrhea treated with metformin in a randomized double blind placebo-controlled trial. *J Clin Endocrinol Metab* 2002; 87: 569-574.
- 37. Kocak M, Caliskan E, Simsir C, Haberal A. Metformin therapy improves ovulatory rates, cervical scores, and pregnancy rates in clomiphene citrate-resistant women with polycystic ovary syndrome. *Fertil Steril* 2002; 77: 101-106.
- 38. Ortega-Gonzalez C, Luna S, Hernandez L, Crespo G, Aguayo P, Arteaga-Troncoso G, et al. Responses of serum androgen and insulin resistance to metformin and pioglitazone in obese, insulin-resistant

- women with polycystic ovary syndrome. *J Clin Endocrinol Metab* 2005; 90: 1360-1365.
- Glueck CJ, Moreira A, Goldenberg N, Sieve L, Wang P. Pioglitazone and metformin in obese women with polycystic ovary syndrome not optimally responsive to metformin. *Hum Reprod* 2003; 18: 1618-1625.
- 40. Romualdi D, Guido M, Ciampelli M, Giuliani M, Leoni F, Perri C, et al. Selective effects of pioglitazone on insulin and androgen abnormalities in normo-and hyperinsulinaemic obese patients with polycystic ovary syndrome. *Hum Reprod* 2003; 18: 1210-1218.
- 41. Aroda VR, Ciaraldi TP, Burke P, Mudaliar S, Clopton P, Phillips S, et al. Metabolic and hormonal changes induced by pioglitazone in polycystic ovary syndrome: a randomized, placebo-controlled clinical trial. *J Clin Endocrinol Metab* 2009; 94: 469-476.
- 42. Stabile G, Borrielli I, Artenisio AC, Bruno LM, Benvenga S, Giunta L, et al. Effects of the insulin sensitizer pioglitazone on menstrual irregularity, insulin resistance and hyperandrogenism in young women with polycystic ovary syndrome. *J Pediatric Adolescent Gynecol* 2014; 27: 177-182.
- 43. Naka KK, Kalantaridou SN, Kravariti M, Bechlioulis A, Kazakos N, Calis KA, et al. Effect of the insulin sensitizers metformin and pioglitazone on endothelial function in young women with polycystic ovary syndrome: a prospective randomized study. *Fertil Steril* 2011; 95: 203-209.
- 44. Pavo I, Jermendy Gr, Varkonyi TT, Kerenyi Z, Gyimesi A, Shoustov S, et al. Effect of pioglitazone compared with metformin on glycemic control and indicators of insulin sensitivity in recently diagnosed patients with type 2 diabetes. J Clin Endocrinol Metab 2003; 88: 1637-1645.
- 45. Carey DG, Cowin GJ, Galloway GJ, Jones NP, Richards JC, Biswas N, et al. Effect of rosiglitazone on insulin sensitivity and body composition in type 2 diabetic patients. *Obes Res* 2002; 10: 1008-1015.
- 46. Fonseca V. Effect of thiazolidinediones on body weight in patients with diabetes mellitus. Am J Med 2003; 115: 42-48.
- 47. Kelly IE, Han TS, Walsh K, Lean M. Effects of a thiazolidinedione compound on body fat and fat distribution of patients with type 2 diabetes. *Diabetes Care* 1999; 22: 288-293.
- Akazawa S, Sun F, Ito M, Kawasaki E, Eguchi K. Efficacy of troglitazone on body fat distribution in type 2 diabetes. *Diabetes Care* 2000; 23: 1067-1071.
- 49. Brettenthaler N, De Geyter C, Huber PR, Keller U. Effect of the insulin sensitizer pioglitazone on insulin resistance, hyperandrogenism, and ovulatory dysfunction in women with polycystic ovary syndrome. J Clin Endocrinol Metab 2004; 89: 3835-3840.
- Qu J, Wang Y, Wu X, Gao L, Hou L, Erkkola R. Insulin resistance directly contributes to androgenic potential within ovarian theca cells. *Fertil Steril* 2009; 91: 1990-1997.
- 51. Garmes HM, Tambascia MA, Zantut-Wittmann DE. Endocrine-metabolic effects of the treatment with pioglitazonein obese patients with polycystic ovary syndrome. *Gynecol Endocrinol* 2005; 21: 317-323.
- 52. Legro RS, Zaino RJ, Demers LM, Kunselman AR, Gnatuk CL, Williams NI, et al. The effects of metformin and rosiglitazone, alone and in

- combination, on the ovary and endometrium in polycystic ovary syndrome. *Am J Obstet Gynecol* 2007; 196: 402.
- 53. Mitkov M, Pehlivanov B, Terzieva D. Metformin versus rosiglitazone in the treatment of polycystic ovary syndrome. *Eur J Obstet Gynecol Reprod Biol* 2006; 126: 93-98.
- 54. Nawaz FH, Khalid R, Naru T, Rizvi J. Does continuous use of metformin throughout pregnancy improve pregnancy outcomes in women with polycystic ovarian syndrome? J Obstet Gynaecol Res 2008; 34: 832-837.
- 55. Ghazeeri G, Kutteh WH, Bryer-Ash M, Haas D, Ke RW. Effect of rosiglitazone on spontaneous and clomiphene citrate—induced ovulation in women with polycystic ovary syndrome. Fertil Steril 2003; 79: 562-566.
- 56. Cataldo NA, Abbasi F, McLaughlin TL, Lamendola C, Reaven GM. Improvement in insulin sensitivity followed by ovulation and pregnancy in a woman with polycystic ovary syndrome who was treated

- with rosiglitazone. Fertil Steril 2001; 76: 1057-1059.
- 57. Yaris F, Yaris E, Kadioglu M, Ulku C, Kesim M, Kalyoncu NI. Normal pregnancy outcome following inadvertent exposure to rosiglitazone, gliclazide, and atorvastatin in a diabetic and hypertensive woman. *Reprod Toxicol* 2004; 18: 619-621.
- 58. Kalyoncu NI, Yaris F, Ulku C, Kadioglu M, Kesim M, Unsal M, et al. A case of rosiglitazone exposure in the second trimester of pregnancy. *Reprod Toxicol* 2005; 19: 563-564.
- 59. Huang T, Chen PC, Wu MH, Lin CY. Metformin improved health-related quality of life in ethnic Chinese women with polycystic ovary syndrome. Health Qual Life Outcomes 2016; 14: 119.
- 60. Ghaneei A, Jowkar A, Hasani Ghavam MR, Ghaneei ME. Cabergoline plus metformin therapy effects on menstrual irregularity and androgen system in polycystic ovary syndrome women with hyperprolactinemia. *Iran J Reprod Med* 2015; 13: 93-100.