

Maternal Anemia during pregnancy and infant low birth weight: A systematic review and Meta-analysis

Shoboo Rahmati¹ B.Sc., Ali Delpishe² Ph.D., Milad Azami¹ M.D., Mohammed Reza Hafezi Ahmadi³ M.D., Kurosh Sayehmiri⁴ Ph.D.

1. Ilam University of Medical Sciences, Ilam, Iran.
2. Department of Epidemiology, Ilam University of Medical Sciences, Ilam, Iran.
3. Department of Pathobiology, Ilam University of Medical Sciences, Ilam, Iran.
4. Department of Biostatistics, Ilam University of Medical Sciences, Ilam, Iran.

Corresponding Author:

Kurosh Sayehmiri, Psychosocial Injuries Research Center, Ilam University of Medical Sciences, Ilam, Iran. P.O. Box: 69391-77143
Email: Sayehmiri@razi.tums.ac.ir
Tel: (+98) 091834107824

Received: 17 May 2016

Revised: 18 October 2016

Accepted: 4 December 2016

Abstract

Background: Infant low birth weight is one of the major problems in different societies. Different reports have provided different results regarding the relationship between maternal anemia and infant low birth weight in different months of pregnancy.

Objective: The aim of this study was to determine the relationship between maternal anemia during pregnancy and infant low birth weight.

Materials and Methods: This systematic review was conducted using related keywords in national (Sid, Iran.doc, Iran medex and Magiran) and international (PubMed, Science Direct, Cochrane, Medline, Web of Science, Scopus, Springer, Embase, Google scholar) databases. Relative risks and confidence intervals were extracted from each study. The results were combined using random-effects model for meta-analysis. The I^2 index was also used to measure heterogeneity between the studies.

Results: Overall, 17 studies with a total sample size of 245407 entered the final meta-analysis and demonstrated that the relative risk for maternal anemia in the first, second and third trimester of pregnancy were 1.26 (95% CI: 1.03-1.55), 0.97 (95% CI: 0.57-1.65), and 1.21 (95% CI: 0.84-1.76), respectively. The relationship between maternal anemia and infant low birth weight in the first trimester of pregnancy was significant.

Conclusion: Maternal anemia, especially during the first trimester of pregnancy, can be considered as a risk factor for pregnancy outcomes. Therefore, one needs to take the necessary steps to cure this disease in order to reduce the incidence of infant low birth weight.

Key words: Anemia, Hemoglobin, Low birth weight, Pregnancy, Pregnant women, Meta-analysis.

Introduction

Low birth weight (LBW) is a major problem in different societies. On average, 7.7% of infants weigh less than 2500 gr. Infants with birth weight less than 2500 gr are known as LBW infants, regardless of gestational age. According to a study, 10% of infants born in the United States are LBW infants (2). Several maternal factors such as age, anemia, etc. may affect LBW of the infants. Therefore, anemia during pregnancy can be considered as one of the major causes of infant LBW (1, 2). According to the data collected from the Center for Disease Control in 1989, if hemoglobin level in

the first and the second trimester of pregnancy is less than 11 g/dL and is less than 10.5 g/dL in the third trimester, this condition is considered anemia (3). According to the recommendations of World Health Organization (WHO) in 1972, a pregnant woman is diagnosed with anemia when the hemoglobin level is below 110 g/l. Based on this report, more than 50% of women who do not take supplements suffer anemia (4). Statistics show that 14-62% of women in developing countries, and 16-29% in developed countries are suffering from anemia (5-7). LBW infants are more vulnerable and exposed to different health problems and complications compared with

infants with normal weight. Based on other reports, there is a direct relationship between Low levels of hemoglobin during pregnancy and high birth weight (8, 9). Considering the high prevalence of anemia among pregnant women aged 17-45, contradictory results regarding the effect of maternal anemia on birth weight and absence of a definite conclusion in obstetrics and genealogy books, we decided to investigate this topic using meta-analysis methods (10, 11). These methods are able to put together all the facts, thus presenting a more clear and accurate picture of the problem to provide a scientifically proved advice for pregnant women (13, 14).

Materials and Methods

Data sources

This study was a systematic review of the literature and methods based on PRISMA guideline from the beginning of 1990 until April 2017 (15). Valid keywords were used for searching in national (Sid, Iran.doc, Iran medex and Magiran) and international (PubMed, Science Direct, Cochrane, Medline, Web of Science, Scopus, Springer, Embase, Google scholar) databases. In order to maximize the comprehensiveness of the study, general Persian keywords and all possible combinations (maternal anemia and LBW) were used in the search. We used English equivalents such as "anemia", "LBW" and "pregnancy" in international databases and the "either/or" structure was applied for further searching; references were also examined to find further necessary sources on the subject.

Study selection (Inclusion and exclusion criteria)

STROBE, which is a standard and famous international checklist, was used to determine the quality of the study (17). This checklist contains 22 sections, covering various

aspects of the study including methodology, sampling, variables measurement, statistical analysis, modification of confounders and the reliability and validity of the tools used to evaluate the study objectives. 15.5 was considered the minimum attainable score. Finally, studies that attained the minimum score of the STROBE checklist entered the study and their data were analyzed using meta-analysis method.

Study selection

First, researchers prepared a list of topics and abstracts of each study at the beginning of study and studies that covered the relationship between maternal anemia and LBW based on pregnancy month entered the main study. Cohort, case-control and cross-sectional that evaluated the relationship between maternal anemia and LBW were analyzed in the present study. Relative risk with 95% confidence interval was used to assess the significance of the study.

Data extraction

Data extraction was performed by two researchers independently and data extraction form (containing name of the author, year of publication, country, continent, number of participants, trimester and LBW pregnancy outcomes) was used to minimize bias and error in collecting data. In cases of necessity, ambiguity of article information and some certain questions, we contacted the authors by E-mail. The data extracted by the two researchers was compared and discussed in case of possible discrepancies; the dispute was shared with a third person and eventual consensus was reached through re-examination of the issue.

Risk of bias assessment

One of the risks of bias assessment was the fact that data extraction was conducted by two researchers. There was also the risk of dispute.

Outcomes of interest

LBW defined as an infant born at less than 2,500 gr. And hemoglobin level in the first and the second trimester of pregnancy is less than 11 g/dL and is less than 10.5 g/dL in the third trimester, this condition is considered anemia (3).

Statistical analysis

In present study, the extracted data included study design, type of the study, name of the author, year of publication, continent, number of participants, trimester and the outcome of pregnancy (LBW). The level of hemoglobin was classified in three groups of <14, 11-13, 11> g/dL, while <11 was considered maternal anemia. LBW, as an outcome of pregnancy, was analyzed when the infant weighed less than 2500 grams. The effect of hemoglobin was evaluated using relative risk index (RRI) in three different phases (first, second, and third trimester) of pregnancy based on the weight of the infant. I² index and Q test were used to assess the heterogeneity. Considering the heterogeneity of the studies, random effects model was used to combine different results. STATA Ver 3.2 was used for data analysis and p<0.05 was considered significant for all analyses.

The minimum and maximum relative risks (RR) or odds ratio (OR) at upper and lower limits of 95% confidence intervals were extracted for each study. In studies, for which this index was not extracted (articles without the OR to use the formula), the index was calculated using $OR = ad/bc$ formula and OR or RR logarithm were used to symmetrize the effect size. Error standard was calculated using the following formula: $SE = 1/n \text{ (upper 95\% CI / lower 95\% CI) } / (2 \times 1.96)$. If only upper and lower limit had been reported in studies, $OR = ad/bc$ was used. Meta-regression was used to investigate the relationship between the year of study, effect size and causes of heterogeneity. Beggs Funnel Plot was used to examine the publication bias. Sensitivity

analysis was used to assess the effect of each study on the overall result.

To analyze sensitivity, the effect of each article was excluded and the result was combined with other studies. This shows whether exclusion of this study has significant effect on the overall result or not. In the case of our study, the effect was not statistically significant.

Results

Overall, 2454 relevant articles were identified by searching databases. Studies that did not yield any helpful and relevant information were excluded; 76% of them were cohort studies with an age range of less than 30 years old. 70% of articles were conducted in Asia. (Diagram 1, Table I)

The relationship between maternal anemia in the first trimester of pregnancy and LBW

By type of study

As illustrated in Figure 1 and table II, there were 9 cohort studies in which the relationship between maternal anemia in the first trimester and LBW was investigated. After combining cohort studies using random effects model, a no significant relationship was observed between maternal anemia in the first trimester of pregnancy and LBW (relative risk, 1.20 [95% CI: 0.98 to 1.46]) (8, 18, 19, 22, 23, 24, 28,29,33). It seems that Levy's study has estimated this relationship with a relatively high RR (8). There were 2 case-control studies about the relationship between maternal anemia and LBW indicating a significant relationship with a RR of 2.40 (95% CI: 1.62-3.55) (11, 12).

It seems that Hamalainen's study has estimated this relationship with relatively high risk (12). And There were 1 cross-sectional studies about the relationship between maternal anemia and LBW indicating a no significant relationship with a RR of 0.72 (95%

CI: 0.52-0.99) (30). However, the total combination of cohort, case-control and cross-sectional studies using random effects model showed a significant relationship between maternal anemia and LBW in the first trimester with a RR of 1.26 (95% CI: 1.03-1.55) (8, 10, 11, 18, 19, 22-24, 28-30, 33).

As illustrated in Figure 1, I^2 index showed that the results of 9 cohort studies were heterogeneous and this heterogeneity was significant (I^2 Index=69.8%, $p=0.001$). The heterogeneity of 2 case-control studies was 0.0% and this heterogeneity was not significant ($p=0.481$). On each article was measured individually and it was found that their weight is calculated when the data is combined. Among cohort studies, Levy's study had the highest weight because of large sample size (8). Kidnato's study had the highest weight among case-control studies (11).

By type of continent

Table II and Figure 2 shows the data analysis based on the continent; about 10 studies were conducted in Asia, 1 in Europe and 1 in Africa. RR, confidence interval, and P value were measured through random effects model, that combining Asian studies using a random effects model and this relationship was no significant (relative risk, 1.13 [95% CI: 0.93 to 1.37]). The data related to European studies (relative risk, 3.14 [95% CI: 1.35 to 7.29]) and African studies 2.23 [95% CI: 1.44 to 3.46]) are shown in Table II and Figure 2 separately. Levy's study in Asia had the highest weight (8).

By year of publication

According to Figure 3 and Table III, there is not significant relationship between publication year and logarithm size and the regression equation was estimated to be $y=41.52-0.02x$. Since p-value was found to be 0.43 for the slope of the line.

By Begg's funnel plot

Figure 4 was used to find the direction of publication bias, according to which the publication direction effect was significant ($p=0.02$).

By Sensitivity analysis

Sensitivity analysis showed that the elimination of a study can either change the results or does not affect the final direction at all and it shows the influence of studies in meta-analysis research. Sometimes, excluding a study can change the overall effect of meta-analysis significantly. In this study, as shown in the figure 5, excluding a study had no effect on the overall results.

The relationship between maternal anemia and LBW in the second trimester at birth

According figure 6 data analysis, 4 studies investigated this relationship in the meta-analysis with a RR of 0.97 (95% CI: 0.57-1.66), while this relationship was not significant. I^2 index for both studies was measured to be 59.3%, showing a homogenous relationship between the 4 studies. The homogeneity was significant ($p=0.0061$) (10, 20, 29, 30).

Relationship between maternal anemia and LBW in the third trimester

Of 8 studies that discussed this issue, only 4 studies showed a significant relationship (9, 29, 31, 32). Then, we combined studies using a random effects model and no significant relationship was observed between maternal anemia in the third trimester of pregnancy and LBW (relative risk, 1.21 [95% CI: 0.84 to 1.76] (9, 10, 16, 20, 29-32). Yildiz's study had the highest weight (31). I^2 index between this studies was 72.8%, which shows a significant heterogeneous relationship between them ($p=0.001$) (Figure 7).

Table I. Characteristics of studies that entered meta-analysis

Author-publication year-reference	Study design	Sample size	Mean age	Country	Trimester	RR (lbw)	CI-	Confounding factors
Levy-2005 (8)	Cohort	153396	28.3±5.9	Israel	First	1.10	1.10-1.20	Ethnicity, maternal age, placental problems, caesarean delivery, and no vertex presentation
Bodeau-livinec-2011 (9)	Clinical trail	1508	<30	Benin	Third	2/8	1.40-5.6	-
Hamalain-2003 (10)	Case-control	22799	28.9±5.9	Finland	First Secondary Third	3.14 0.61 1.28	1.35-7.28 0.25-1.48 0.63-2.62	No specify
Kidnato-2009 (11)	Case-control	1721	24	Tanzania	First	2.23	1.45-3.5	No specify
Knotterus -1990(16)	Cohort	796	<30	Netherland	Third	0.86	0.35-2.03	-
Zhou-1998 (18)	Cohort	829	25.5±3.8	China	First	2.96	1.15-7.62	No specify
Abeyseena-2010 (19)	Cohort	817	26.4±5.5	Srilanka	First	0.56	0.07-4.74	No specify
Chang-2003 (20)	Cohort	918	16.1±1.1	USA	Secondary Third	1.48 0.70	0.55-3.98 0.42-1.25	Parity, BMI, smoking, preeclampsia, and antenatal care
Monawar hosian-2006 (22)	Cohort	350	NA	Bangladesh	First	2	1.18-3.40	No specify
Xiong-1996 (23)	Cohort	16936	25±2.8	China	First	0.79	0.55-1.12	Hospital stay, maternal age, maternal education, parity, gestational age at the first prenatal visit, BMI, hypertensive disorder in pregnancy, vaginal bleeding, and prior spontaneous abortion
Kumar-2010 (24)	Cohort	2027	24.6±3.82	India	First	1.02	0.83-1.25	Maternal age, parity, maternal height, maternal weight, BMI, and gestational age
Sekhvat-2011 (28)	Cohort	1842	-	Iran	First	1.40	1.15-2.06	-
Huang -2015 (29)	Cohort	500	-	China	First Secondary Third	0.61 1.62 2.07	0.32-1.15 0.89-2.90 1.08-3.97	-
Yang-2017 (30)	Cross-sectional	7375	-	china	First Secondary Third	0.72 0.67 0.47	0.50-0.95 0.42-0.98 0.24-0.93	-
Yildiz- 2014 (31)	Cohort	28600	-	turkey	Third	1.08	1.05-1.11	-
Nair - 2016 (32)	Cohort	1007	-	India	Third	6.19	1.44-26.71	-
Chikjwa-2015 (33)	Cohort	1986	-	China	First	2	1.30-3.10	-

Median

LBW: low birth weight

RR: Relative risk

CI: Confidence interval

Table II. The relationship between maternal anemia and LBW in the first trimester based on the continent and study type

Variable	Number of studies	OR	95%CI	I ² %	P (heterogeneity)
Type of study					
Cohort	9	1.20	0.98-1.46	69.8	0.001
Case-control	2	2.40	1.62-3.55	0.0	0.481
Cross-sectional	1	0.72	0.52-0.99	-	-
Mixture of studies	12	1.26	1.03-1.55	77.5	0.000
Continent					
Asia	10	1.13	0.93-1.37	72.9	0.000
Africa	1	2.23	1.44-3.46	0	-
Europe	1	3.14	1.35-7.29	0	-
Aggregate of continents	12	1.26	1.03-1.55	77.5	0.000

OR: odds ratio

CI: confidence interval

P: p-value

Table III. The relationship between maternal anemia and LBW in the first trimester by continent based on year of publication and sample size

		Coefficients	STD. Err*	P-value
Model 1	Date of Publication	-0.02	0.025	0.43
	Constant	41.52	50.3	0.42

*standard error of coefficients. P-value computed using T statistics




Diagram 1. Entry procedures of studies into meta-analysis study




Figure 1. The relationship between maternal anemia and LBW in the first trimester based on the type of study




Figure 2. The relationship between maternal anemia and LBW in the first trimester based on continent




Figure 3. The relationship between maternal anemia and LBW in the first trimester based on the year of publication




Figure 4. Begg's funnel plot for relationship of maternal anemia with low birth weight in the first trimester of pregnancy




Figure 5. Sensitivity analysis for relationship between maternal anemia with low birth weight in the first trimester of pregnancy




Figure 6. The relationship between maternal anemia and LBW in the secondary trimester of pregnancy




Figure 7. The relationship between maternal anemia and LBW in the third trimester of pregnancy

Discussion

The present study is the first systematic review and meta-analysis research about the relationship between maternal anemia and LBW. Different studies with different values of relative risk and confidence interval were previously conducted (8-11, 16, 18-20, 22-24, 28-33). Various maternal criteria such as anemia, age, intervals between pregnancies, physical tumor factor and weight loss were included in this study. The present study investigated the relationship between maternal anemia and infant LBW.

According to the results, the relationship between maternal anemia and LBW in the first

trimester of the pregnancy is significant. maternal anemia causes LBW in the first trimester of the pregnancy. However, this relationship was not significant for the second and third trimesters of the pregnancy. Nowadays, maternal anemia is considered as a public health problem in the world, especially in developing countries. For example, Statistics show that 14-62% of women in developing countries, and 16-29% in developed countries are suffering from anemia (5-7). That the analysis also showed that this relationship is commonly observed in Asia; i.e. LBW is widely seen in Asian countries compared to developed countries according to a study by Lee *et al* (21, 25).

Sukrat and Haider's review article, which analyzed the relationship between density of hemoglobin and outcomes of pregnancy in 2013, showed that hemoglobin less than 11 g/dl increases LBW risk in the first trimester, which is consistent with the present study (26, 27). The present results of meta-regression analysis for the publication year of study showed that there is no statistically significant relationship between publication year of study with the effect size ($p=0.43$).

Publication bias, which was mostly observed in meta-analysis and review studies, was due to the fact that published articles did not represent the all researches carried out in that specific area. Actually, articles with significant results, high quality precise design and extensive sample size had higher chances of publications. Another reason is that both native and English languages are used and allowed in meta-analysis studies. According to the results, bias had played a key role in this type of publications. Usually, studies which are conducted according to the norms of relationship assessment increase the chance of bias because articles with positive results are more likely to get published and they are more widely searched.

Limitations

1. Exclusion of the relevant and new studies that were not eligible based on inclusion criteria (in terms of Epidemiology)
2. Inability of databases to search using combined keywords
3. Some studies were excluded due to low quality and insufficient proofs
4. Exclusion of some studies due to ambiguity in the month of pregnancy
5. Different definitions for anemia in different studies
6. Exclusion of some studies and articles due to improper direction of publications and lack of causative relationship.

Conclusion

Considering the relationship between maternal anemia and LBW in the first trimester of pregnancy, anemia can be used

as a preventive factor for infant LBW. Therefore, we need to cure maternal anemia to prevent or cure LBW.

Acknowledgements

Hereby, the authors wish to express their deepest sense of gratitude and indebtedness to Research Council of Ilam University of Medical Sciences (Grant Number: 910597) for their financial support.

Conflict of interest

There is no conflict of interest.

References

1. Ramazanali F, Vahid Dastjerdi M, Beigi A, Moini A. [The relationship between maternal HCT levels, birth weight and risk of low birth weight]. *Iran J Pediatr* 2006; 16: 447-454. (In Persian)
2. Cunningham FG, Leveno KJ, Bloom SL, Hauth JC, Gilstrap III L, Wenstrom KD. Williams's obstetrics. 22nd Ed. Philadelphia, McGraw-Hill; 2005:1143-1167.
3. Steer P, Alam MA, Wadsworth J, Welch A. Relation between maternal haemoglobin concentration and birth weight in different ethnic groups. *BMJ* 1995; 310: 489-491.
4. Mirzaie F, Eftekhari N, Goldozeian S, Mahdavinia J. Prevalence of anemia risk factors in pregnant women in Kerman, Iran. *Iran J Reprod Med* 2012; 8: 67-69.
5. Shobeiri F, Begum K, Nazar M. A prospective study of maternal hemoglobin status of Indian women during pregnancy and pregnancy outcome. *Nutr Res* 2006; 24: 209-213.
6. Perry IJ, Beevers DG, Whincup PH, Bareford D. Predictors of ratio of placental weight to fetal weight in multiethnic community. *BMJ* 1995; 310: 436-439.
7. Levy A, Fraser D, Katz M, Mazor M, Shiener E. Maternal anemia during pregnancy is a independent risk factor for low birth weight and preterm delivery. *Eur J Obstet Gynecol Reprod Biol* 2005; 122: 182-186.
8. Salimi S, Nakhostin B, Alijahan J, Hazrati S. Relationship Hemoglobin and mother bmi in pregnancy with low birth weight. *J Ardebil Univ Med Sci* 2012; 15: 14-20.
9. Bodeau-Livinec F, Briand V, Berger J, Xiong X, Massougbdji A, Day K, Cot M. Maternal Anemia in Benin: Prevalence, Risk Factors, and Association with Low Birth Weight. *Am Soc Trop Med Hygiene* 2011; 85: 414-420.
10. Nasiri Amiri F, Ahmadi M. Relationship between mother hematocrit and pregnancy result. *J Gorgan Univ Med Sci* 2007; 9: 29-34.
11. Kidanto HL, Mogren I, Lindmark G, Massawe S, Nystrom L. Risks for preterm delivery and low birth weight are independently increased by severity of maternal anaemia. Department of Clinical Science, Obstetrics and Gynaecology, Umeå University. 2009;

- 99: 98-102 .
12. Hamalainen H, Hakkarainen K, Heinonen S. Anemia in the first but not in the second or third trimester is a risk factor for low birth weight. *Clin Nutr* 2003; 22: 271-275.
 13. Faraji R, Mirbolok F, Sharemi S.H, Asgharnia M, Afshani JO, Gholamzadeh M. Relationship between maternal hemoglobin concentration and BMI during the first trimester in primiparous women and her neonatuls birth weight. *Iran Surg J* 2010; 18: 62-68.
 14. Garshasbi A, Fallah N. Maternal hematocrite level and risk of low birth weight and preterm Delivery. *Tehran Univ Med J* 2006; 64: 87-94.
 15. Bondevik GT, Lie RT, Ulstein M, Kvale G. Maternal hematological status and risk of low birth weight and preterm delivery in Nepal. *Acta Obstet Gynecol Scand* 2001; 80: 402-408.
 16. Morrone A, Nosotti L, Piombo L, Scardella P, Spada R, Pitidis A. Iron deficiency anaemia prevalence in a population of immigrated women in Italy. *Eur J Public Health* 2010; 22: 256-262.
 17. Knotterus JA, Delgado LR, Knipschid PG, Essed GG, Smirt F. Hamatologic parameters and pregnancy outcome. *J Clin Epidemiol* 1990; 43: 461-466.
 18. Zhou LM, Yang WW, Hua JZ, Deng CHQ, Tao X, Rebecca J. Relation of Hemoglobin Measured at Different Times in Pregnancy to Preterm Birth and Low Birth Weight in Shanghai, China. *Am J Epidemiol* 1998; 148: 998-1006.
 19. Abeyseena CH, Jayawardana P. Maternal haemoglobin level at booking visit and its effect on adverse pregnancy outcome. *Aust N Z J Obstet Gynaecol* 2010; 50: 423-427.
 20. Chang SHCH, Brien K, Nathanson M, Mancini J, Frank R. Hemoglobin Concentrations Influence Birth Outcomes in Pregnant African-American Adolescents. *J Nutr* 2003; 133: 2348-2355.
 21. Lee HS, Kim MS, Kim MH, Kim YJ, Kim WY. Iron status and its association with pregnancy outcome in Korean pregnant women. *Eur J Clin Nutr* 2006; 60: 1130-1135.
 22. MonawarHosain M, Chatterjee N, Begum A, Saha S.CH. Factors Associated with Low Birthweight in Rural Bangladesh. *J Trop Pediatr* 2005; 52: 87-91.
 23. Xiong X, Buekens P, Alexander S, Wollast E, "The relationship between anemia during pregnancy and birth outcome." *Arch Public Health* 1996; 53 (Suppl.): 136.
 24. Kumar A, Chaudhary K, Prasad S. Maternal indicators and obstetric outcome in the north Indian population: a hospital- based study. *J Postgraduate Med* 2010; 56: 192-195.
 25. Rahman MM, Abe SK, Rahman MS, Kanda M, Narita S, Bilano V, et al. Maternal anemia and risk of adverse birth and health outcomes in low-and middle-income countries: systematic review and meta-analysis. *Am J Clin Nutr* 2016; 103: 495-504.
 26. Sukrat B, Wilasrusmee C, Siribumrungwong B, McEvoy M, Okascharoen C, Attia J, Thakkinstian A. Hemoglobin concentration and pregnancy outcomes: a systematic review and meta-analysis. *BioMed Res Int* 2013; 2013: 1-9.
 27. Haider BA, Olofin I, Wang M, Spiegelman D, Ezzati M, Fawzi WW. Anaemia, prenatal iron use, and risk of adverse pregnancy outcomes: systematic review and meta-analysis. *BMJ* 2013; 346: 1-19.