Triage by cervical length sonographic measurements for targeted therapy in threatened preterm labor: A double blind randomized clinical trial

Homeira Vafaei¹ M.D., Neda Rahimirad¹ M.D., Seyedeh Marjan Hosseini² M.D., Maryam Kasraeian¹ M.D., Nasrin Asadi¹ M.D., Hadi Raeisi Shahraki³ M.D., Khadijeh Bazrafshan M.Sc.⁴

- 1. Maternal-Fetal Medicine Research Center, Perinatology Ward, Shiraz University of Medical Sciences, Shiraz, Iran.
- Department of Obstetrics and Gynecology, Shiraz University of Medical Sciences, Shiraz, Iran.
- 3. Department of Biostatistics, School of Medicine, Shiraz University of Medical Sciences, Shiraz, Iran.
- 4. Maternal-Fetal Medicine Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

Corresponding Author:

Homeira Vafaei, Maternal-Fetal Medicine Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.

Email: Vafaeih@gmail.com **Tel:** (+98) 9173430210

Received: 28 September 2016 Revised: 1 July 2017 Accepted: 17 October 2017

Abstract

Background: Preterm labor and birth are associated with several neonatal complications including respiratory distress syndrome and intraventricular hemorrhage. Differentiating true and false labor pain is a dilemma to obstetricians.

Objective: To elucidate the role of cervical length measurement in prediction of birth in pregnant women with threatened preterm labor.

Materials and Methods: In this double blind randomized clinical trial, 120 women with gestational age <34 wk who presented painful uterine contractions randomly assigned to undergo measurement of cervical length. Patients were registered in the hospital and a unit number was given. Based on the unit numbers, patients were randomly assigned to two groups using a computerized random digit generator. All participants were managed accordingly (n=65) or to receive tocolysis as planned (n=55). Tocolysis was prescribed when cervical length was <15 mm while those with cervical length ≥15 mm were managed expectantly. Delivery within 7 days of the presentation was the primary outcome.

Results: This RCT showed in case group, 78.9% of patient with cervical length <15 mm were delivered within 7 days and only 21.1% of them maintained their pregnancy. Of those with cervical length ≥ 15 mm, only 15.2% were delivered within the study period and the rest (84.8%) maintained their pregnancy (p<0.001).

Conclusion: "Our results indicate that in women who presented preterm labor symptoms, cervical length measurement will result in decreased unnecessary tocolytic treatment. Women with cervical length ≥15mm should not receive tocolysis, however, withholding corticosteroid therapy in these patients needs further evidence.

Key words: Cervical length, Preterm labor, Tocolysis, Ultrasonography. **Registration ID in IRCT: IRCT2017092317034N5**

Introduction

reterm labor defined as occurrence of uterine contractions accompanied by cervical changes (dilation and effacement) before 37 wk of gestation is considered ,estimated prevalence of preterm birth is 5-10% of all pregnancies (1). Preterm birth is increased short term complications ascribed to immaturity of multiple organ systems &long term adverse effects on health in whole life (2). Vasodilatory effects of progesterone on the vessels of fetus and uterus, seems be effective in the prevention of preterm birth (3).

However, the drug of choice is yet to be identified because of complications associated with most of them (3, 4). Several criteria have been introduced for diagnosis of preterm

labor. According to Creasy, preterm labor is uterine contractions as contractions per 20 min), cervical dilation 2≥ cm in a nulliparous and 3≥ cm in a multipara) and cervical effacement (>80%) or uterine contractions and cervical change However, preterm labor is usually overdiagnosed and unnecessary tocolytic and corticosteroid treatment are administered (6). Several other methods have been used for diagnosis of preterm labor including transvaginal ultrasonography. Measurement using cervical length transvaginal ultrasonography has been shown to predict preterm birth in those presenting with threatened preterm labor (7, 8). A systematic review done in 2008 showed cervical length measurement using transvaginal has ultrasonography high accuracy

predicting spontaneous preterm birth in high risk asymptomatic population (9).

Alfirevic and his coworkers found that those women with threatened preterm labor who has cervical length more than 15 mm should not receive any tocolysis because the preterm birth will not occur in these patients (10). In another study by Tsoi and his colleagues, 99% of women with cervical length ≥15 mm did not deliver within 7 days without receiving tocolytic therapy (11).

According to these observations and in order to elucidate the role of cervical length measurement in prediction of preterm birth in those with threatened preterm labor, we performed this randomized clinical trial.

Materials and methods

This randomized clinical trial was performed in Hafez hospital, Shiraz University of Medical Sciences, Shiraz, Iran during November 2010 to February 2015.

All women with live, singleton pregnancies who presented true uterine contractions before 34 wk of gestation age (at least 4 contractions in 20 min interval) that were selected to receive tocolytic and steroid therapy were included. Cervical effacement was not considered as a criterion for inclusion or exclusion in this trial. Women with cervical dilatation >5 cm, polyhydramnios (amniotic fluid index greater than 24 cm), macrosomia (estimated fetal weight >90th centile), multiple pregnancy, suspected intrauterine infection, rupture of membranes, vaginal bleeding (more than bloody show), major medical disorders in mother, contraindicated for vaginal delivery (e.g. placenta previa), non-reassuring fetal heart monitoring, fetal anomaly, any contraindication for administer of betadrugs sympathomimetic and previous treatment with tocolytics/corticosteroid in this pregnancy were excluded. No other tests including fibronectin were used for diagnosis of preterm labor. Transvaginal sonography for cervical length measurement was done by an expert obstetrician."

Study protocol

Patients were selected by labor residents at the time of admission. A thorough history and physical examination was performed by the residents including the digital vaginal examination. The findings including the fetal presentation, cervical dilatation, effacement

and past medical history were recorded in the structured questionnaire. A total of 120 patients with preterm labor were enrolled in this study being randomized to two groups including: 65 patients undergoing cervical length measurement by transvaginal ultrasonography for further management (case group) and 55 patients who received tocolytic and corticosteroid therapy (control group).

Those who were assigned to the case underwent transvaginal group ultrasonography for measuring the cervical length by one expert obstetrician. Neither residents nor sonographers were not aware of this clinical trial. Those with cervical length mm were received tocolvtic <15 corticosteroid therapy according hospital protocol. Expectantly management was performed when cervical length was ≥15 mm, but with persistent uterine contractions more than 4 hr, another cervical measurement was performed by the same operator. If the cervical length was become <15 mm tocolytic and antenatal corticosteroids were given.

Patients were followed for 7 days after enrollment in the study and the uterine contractions were screened. If the pain and uterine contractions were resolved, the patient was discharged and followed by the phone. All the participants were asked to record their uterine contraction in a diary manner. All the patients who had not delivered within the first 7 days after enrollment in the study, underwent uterine contraction check-up. The primary outcome of study was proportion of the patients who maintained their pregnancy at least 7 days after randomization. All the patients were observed precisely in the labor suite.

Ethical consideration

The study protocol was approved by the institutional review board and Ethics Committee of Shiraz University of Medical Sciences, Shiraz, Iran (Ct-88-5026, 1388-12-27). All the participants patients in case & control groups gave their informed written consent.

Statistical analysis

Descriptive statistics were reported as mean±SD for quantitative or frequency (%) for qualitative variables. Association between two categorical variables was assessed using chisquare or Fisher's exact test where

appropriate. Normality of all the quantitative variables was approved in both groups via Shapiro-Wilk test (P>0.05 in all variables) therefore independent t-test was used to examine the difference between two groups.

All the statistical analysis performed in SPSS (Statistical Package for the Social Sciences, version 18.0, SPSS Inc, Chicago, Illinois, USA) and p<0.05 was considered significant.




Figure 1. CONSORT flow diagram of the participants.

Results

The mean age of the participants was 28.4±4.3 (range 18-36 yrs.). The mean age of the patients and controls was found to be 24.8±4.6 and 24.81±4.1 yrs. respectively (p=0.978). Table I compares the demographic characteristics of cases and controls. The cervical dilation was significantly higher in the controls compared to the case group (p=0.003). While the control group had significantly higher effacement compared to case group (50.91±22.7% vs. 59.61±13.6%; p=0.015).

Table II summarizes the pregnancy characteristics of two study groups. Cervical length in all of the 65 patients in case group was measured with a cut-off value of 15 mm. The mean cervical length in the cases was 21.4±7.9 mm (range 8-37 mm). 46 patients (70.7% of case group) had cervical length ≥15 mm, and only 19 patients (29.3% of case group) had cervical length <15 mm and received tocolytic therapy.

Among 46 patients in case group with the cervical length ≥15 mm, uterine contractions persisted in 6 patient (13%) and cervical length measurement was repeated for them after 4 hr. In these 6 patients, mean cervical length was 20.6±5.4 mm (range 16-28 The mean cervical length decreased significantly in those with persistent contraction (p=0.008); however, none of them had cervical length <15 mm and thus did not receive therapy.

Of those 19 (29.3%) patients who had cervical length <15mm, 15 (78.9%) delivered during the study period and only 4 (21.1%) maintained their pregnancy. From 46 patients with cervical length ≥15 mm, only 7 (15.2%) delivered within the study period and the rest (84.8%) maintained their pregnancy. The difference was significant at the level of p<0.001. Overall, 22 patients (0.33%) in case group delivered during the study period, compared to 13 patients (28%) in control group, the difference was not significant (p=0.31).

Table I. Baseline characteristics of patients in two study groups

Variables	Case	Control	n volue		
variables	group(n=65)	group (n=45)	p-value		
Age (yr)*	24.8 ± 4.6	24.8 ± 4.1	0.978†		
BMI(Kg/m²)*	25.7 ± 2.4	25.3 ± 2.4	0.324†		
Gravidity**					
Prim gravid	24 (36.9)	26 (47.3)	0.69744		
Multigravid	41 (63)	29 (52)	0.687††		
Parity**					
Nulliparous	32 (49.2)	32 (58.2)	0.608††		
Multiparous	33 (50.8)	23 (41.8)			
Previous history of abortion**					
No abortion	49 (75.4)	43 (78.2)			
One abortion	13 (20.0)	10 (18.2)	0.928††		
Two abortion	3 (4.6)	2 (3.6)			
Previous history of stillbirth**					
No stillbirth	59 (90.8)	53 (96.4)	0.10044		
One still birth	6 (9.2)	2 (3.6)	0.198††		

^{*} Data presented as mean±SD; ** Data presented as n (%).

[†]Independent *t*-test

^{††}Chi-square test

BMI: Body mass index

Table II. Pregnancy characteristics in two study groups

	Case group (n= 65)	Control group(n= 55)	p-value
Gestational age by LMP (wk)*	30.7 ± 1.6	30.9 ± 1.9	0.586
Gestational age by sonography (wk)*	30.9 ± 1.6	30.9 ± 1.9	0.901
Cervical dilatation (Cm)*	1.6 ± 0.7	1.9 ± 0.5	0.003
Cervical effacement (%)*	50.91 ± 22.7	59.61 ± 13.6	0.015
delivery during the study period**	22 (33)	13 (28)	0.31

^{*} Data presented as mean±SD; ** Data presented as n (%). Independent *t*-test

Table III. Relation of cervical length and duration of

pregnancy

Cervical length	Total	Delivery within 7days	Maintaining pregnancy	p-value
≥15 mm <15 mm	46 (70.7) 19 (29.3)	7 (15.2) 15 (78.9)	39 (84.8) 4 (21.1)	< 0.001
<13 IIIII	19 (29.3)	13 (78.9)	4 (21.1)	

Data presented as n (%) Chi-square test

Discussion

We found that approximately 80% of patients who has cervical length <15 mm will deliver within a 7-day period despite receiving tocolytic therapy and this means that preterm birth is inevitable when the cervical canal is less than 15 mm in length. So, prescribing tocolytic agents and corticosteroids to this group seems to be reasonable and will provide the patient with a favorable result of neonate lung maturity and decreased prevalence of respiratory distress syndrome.

In another group, those with cervical length ≥15 mm, 85% maintained their pregnancy without tocolytic any need for and avoiding corticosteroid thus therapy, unnecessary treatment. These results are in agreement with several previous studies that have shown the beneficial advantages of cervical length measurement in threatened preterm labor and eligibility of selecting the patient for treatment with tocolytic agent and corticosteroids (10, 11).

In this regards, Alfirevic and coworkers performed a randomized clinical trial in order to test the supposition of managing threatened preterm labor based on measuring of cervical length by ultrasonography can decrease the number of women who receive inadvisable treatment. They included 41 patients with threatened preterm labor who were assigned to receive tocolytic and corticosteroid therapy. They divided their patients into two groups to undergo cervical length measurement (n=21) or to receive conventional treatment (n=20). The treatment was avoided in 14 women who had cervical length ≥15 mm while the other 7 patients received tocolytic and corticosteroids. prescription of corticosteroids significantly higher in control group compared to ultrasonography group (14% vs. 90%; relative risk (RR) 0.16; 95% CI, 0.05-0.39). Prevention of tocolytic therapy was also significant between two groups (33.3% vs. 100%; RR 0.3; 95% CI, 0.15-0.54). None of the neonates were born without receiving a full course of corticosteroids (10).

In another study by Tsoi and coworkers, the efficacy of cervical length measurement for distinguishing between true and false labor was examined. They included 216 singleton pregnancies at 24-36 (mean, 32) wk of gestation referring with threatened preterm Delivery within 7 days labor. presentation was considered as the primary outcome. Of 173 patients with cervical length ≥15mm, only 1 delivered within the 7dayperiod. Of 43 patients with a cervical length <15mm, 16 delivered within the 7 days of which 6/14 (42%) were treated with tocolytics and 10/29 (35%) were managed expectantly. They showed that cervical length was the only predictor of preterm birth (OR=101, 95% CI 12-800, p < 0.0001) independent of other demographic factors including ethnicity, parity, maternal age, gestational age, body mass index, history of previous preterm delivery, smoking, frequency of contractions or use of any tocolytics (11).

Fuchs and colleagues also tried to investigate the role of cervical length measurement in differentiating false and true labor pains they included 253 singleton pregnancies presenting with painful uterine

contractions before 34 wk of gestation. Cervical length was measured and the cut-off of 15mm was considered the reference for prescribing treatment. Delivery within 7 days of the presentation was considered the main outcome. 21/253 (8.3%) delivered within the study period was reversely correlated with cervical length. Only 4 (1.8%) of 217 patients with cervical length ≥15 mm delivered within 7 days while 47.2% of those with a cervical length <15 mm delivered within 7 days' despite receiving tocolytic therapy. The delivery in labor patients was threatened preterm attributed to cervical length, frequency of contractions at presentation, history preterm delivery and vaginal previous bleeding. It was not found to have correlation with ethnicity, maternal age, parity, smoking or using tocolysis, antibiotics or steroids. It is corroborated by univariate as well as multivariate analyses that cervical length is a significant independent predictor of delivery within 7 days in this population (12).

All these studies, like ours, show that measuring cervical length in those presenting with threatened preterm labor is a powerful tool for differentiating false and true labor pains and using this modality will decrease the unnecessary treatment. However, a question rises here and that is whether this approach will result in widespread of transvaginal ultrasonography. Withholding corticosteroid therapy in those with cervical length ≥15 mm will also result in babies born with inadequate corticosteroid therapy. However, it was shown that this risk is unlikely to be greater than 2-3% (14).

Prospective cohort following patients till birth is needed in order to clarify this issue. In conclusion, measurement of cervical length in those with threatened preterm labor will result in reducing inappropriate tocolytic treatment. Although in women with cervical length ≥15 mm should not administer tocolysis, the prevention of using corticosteroid therapy in these patients needs further evidence.

Acknowledgments

This work was supported by vice-chancellor of research, Shiraz University of Medical Sciences, Shiraz, Iran.

Conflict of interest

The authors declare that there is no conflict of interests regarding the publication of this paper.

References

- Martin JA; Centers for Disease Control and Prevention (CDC). Preterm births-united states, 2007. MMWR 2011; 60 (Suppl.): 78-79.
- Min J, Watson HA, Hezelgrave NL, Seed PT, Shennan A. Ability of preterm surveillance clinic triage risk of preterm birth: a prospective cohort study. Ultrasound Obstet Gynecol 2016; 48: 38-42.
- 3. Vafaei H, Zamanpour T, Raeisi-Shahraki H. Preterm birth prevention: effect of vaginal progesterone administration on blood flow impedance in uterinefetal circulation by doppler sonography. *Glob J Health Sci* 2015; 8: 172-178.
- Creasy RK. Preterm birth prevention: where are we? *Am J Obstet Gynecol* 1993; 168: 1223-1230.
- King J, Flenady V. Prophylactic antibiotics for inhibiting preterm labor with intact membranes. Cochrane Database Syst Rev 2002; 4: CD000246.
- Leitich H, Brunbauer M, Kaider A, Egarter C, Husslein P. Cervical length and dilatation of the internal cervical os detected by vaginal ultrasonography as markers for preterm delivery: A systematic review. Am J Obstet Gynecol 1999; 181: 1465-1472.
- Vendittelli F, Volum'enie JL. Transvaginal ultrasonography examination of the uterine cervix in hospitalized women undergoing preterm labor. Eur J Obstet Gynecol Reprod Biol 2000; 90: 3-11.
- Honest H, Bachmann LM, Coomarasamy A, Gupta JK, Kleijnen J, Khan KS. Accuracy of cervical transvaginal. sonography in predicting preterm birth: a systematic review. *Ultrasound Obstet Gynecol* 2003; 22: 305-322.
- Crane JM, Hutchens D. Transvaginal sonographic measurement of cervical length to predict preterm birth in asymptomatic women at increased risk: a systematic review. *Ultrasound Obstet Gynecol* 2008; 31: 579-587.
- Alfirevic Z, Allen-Coward H, Molina F, Vinuesa CP, Nicolaides K. Targeted therapy for threatened preterm labor based on sonographic measurement of the cervical length: a randomized controlled trial. *Ultrasound Obstet Gynecol* 2007; 29: 47-50.
- 11. Tsoi E, Akmal S, Rane S, Otigbab C, Nicolaides KH. Ultrasound assessment of cervical length in

[DOR: 20.1001.1.24764108.2017.15.11.3.1]

- threatened preterm labor. *Ultrasound Obstet Gynecol* 2003; 21: 552-555.
- 12. Fuchs IB, Henrich W, Osthues K, Dudenhausen JW. Sonographic cervical length in singleton pregnancies

with intact membranes presenting with threatened preterm labor. *Ultrasound Obstet Gynecol* 2004; 24: 554-557.